

Sentinel-2 images for the monitoring of dissolved contaminants on Sicily's South-east coast

Anselme Muzirafuti¹, Giovanni Randazzo², Stefania Lanza², Antonio Crupi³, Maria Cascio¹, Giovanni Barreca¹, and Francesco Gregorio¹

¹Interreg Italia – Malta – Progetto : Pocket Beach Management & Remote Surveillance System (BESS)

²Università degli Studi di Messina, Dipartimento di Scienze Matematiche e Informatiche, Scienze Fisiche e Scienze della Terra, Messina; GEOLOGIS s.r.; Interreg Italia – Malta – Progetto : Pocket Beach Management & Remote Surveillance System (BESS)

³GEOLOGIS s.r.; Interreg Italia – Malta – Progetto : Pocket Beach Management & Remote Surveillance System (BESS)

November 23, 2022

Abstract

Coastal environment pollution is one of major challenge facing coastal area development. A large number of Sicilian population lives in low laying area where, among other economy activities, tourist and intensive agriculture are two of most important sources of revenue. However, its beaches are threatened by natural and anthropic phenomena ranging from sea level rise, coastal erosion to marine pollution. In efforts to manage the influx of harmful substances reaching Sicily's south-east coast, we present methodological approach integrating land use\land cover and normalized difference suspended sediment index mapping techniques to identify the origin of dissolved sediments observed in coastal water especial during rainy season. In this paper we used 12 freely-available Copernicus sentinel-2 images acquired during 2018. Water color analysis on satellite images shows that the contaminants reach the sea through river runoff and when arrived in relatively calm water in particularly in pocket beaches they stay in near shoreline for a while however they can also be exported further offshore when they reach the sea in high dynamic water where currents and waves play a role in their distribution. These sediments are sometimes erosion products from agricultural fields or contaminated soils containing organic or chemical fertilizers used for crop protection or for vegetation growth. The monitoring of sediments can help characterize and determine the fate of these products for a better management of this coastal area.

Sentinel-2 images for the monitoring of dissolved contaminants on Sicily's south-east coast



Anselme Muzirafuti¹, Giovanni Randazzo², Stefania Lanza³, Antonio Crupi⁴, Maria Cascio⁵, Giovanni Barreca⁶, and Francesco Gregorio⁷
 Università degli Studi di Messina, Dipartimento di Scienze Matematiche e Informatiche, Scienze Fisiche e Scienze della Terra, Messina, Italy (1 muzansel@gmail.com, 2 grandazzo@unimme.it, 3 lanzas@unime.it, 4 antoniocrupi5@hotmail.it, 5 maria.cascio86@gmail.com, 6 giovanni_barreca@hotmail.it, 7 gregoriofrancesco92@gmail.com)



1. Introduction

Coastal environments are facing numerous challenges due to urban development, anthropic activities, seas level rise, Population growth, marine pollution, among others. These problems affect not only economic development but also marine ecosystem functioning, Devlin 2015. During the period of high precipitation the coast of Sicily receives amount of sediments reaching near shore water from the land through river runoffs. These sediments are the results of erosion of agricultural fields or high Slope bare land and they are one of main sources of pollutants and contaminants entering in marine ecosystem, Catania 2015, Devlin 2015. Recent studies conducted in coastal areas showed that the influx of sediments from river runoff contain harmful substances, Waterhouse 2012, adsorbed, dissolved or suspended in water. These sediment-plumes can be monitored in coastal waters using satellite image and determine the origin and their destination by observing their dispersion in two dimensions on the sea surface. As traditional accurate water quality monitoring requires intensive field water, Liu 2017, Sentinel-2 has an advantage of providing multi-temporal water quality information on large area at very low cost, Caballero 2018. The aim of this paper is to use various indices algorithms (colored dissolved organic matter (CDOM), normalized difference chlorophyll index (NDCI), normalized suspended material index (NSMI) and land cover/ land use techniques to estimate the composition, the shapes and the destination of sediments- plume observed in the water of the South-east coast of Sicily. The proposed procedure is the combination of qualitative and quantitative interpretation for the dynamics of sediments from the land to the deep sea. Due to the high dynamics of Ionian Sea, this work contribute to the effort of monitoring marine pollution where freely available satellite images can be used to identify marine sites for in-situ measurements.. We intend to provide answers regarding the fate and the origin of dissolved pollutants introduced in coastal environment such as these pollutants destined to remain in one location or spread far away and their impact on coastal environment inhabitants.

2. Material and methods

2.1. Area of study

The area of study is located on the South-east coast of Sicily, Figure 1, stretching from Taormina in the North to Syracuse in the South by passing on the coast of Catania. It is an important touristic site destination with antique theatre and monuments, Etna Mountain and beautiful beaches. Economically this region relies mainly on tourism and agricultural activities. Two main rivers Alcantara near Taormina and Simeto near Catania, the amount of sediments brought by these rivers from land to the sea play a big role in beach nourishment, Lanza 2011, but also contaminate coastal environment by bringing pollutants from industries and agricultural activities all along their courses, Guglielmo, 2013. The area is the site of high Dynamic currents shown in Figure 2, characteristic of wave-dominated Mediterranean Sea, Longhitano 2007, which influence the distribution of these sediments





Table 1. Sentinel-2 images used


Acquisition Time	Acquisition date	Sen2Cor atmospheric correction	Cloud cover %	Spatial resolution of the bands used	Spectral bands used
09:40:19	12-09-2018	Yes	0.29	10 meters	Blue (B2), Green (B3), Red (B4), Near-infrared (B8),
09:50:31	30-09-2018	Yes	2.42		
09:40:31	07-10-2018	Yes	1.67		
09:50:31	20-10-2018	Yes	31.92		
09:50:59	25-10-2018	Yes	1.01		
09:51:21	30-10-2018	Yes	0.51		
09:52:21	09-11-2018	Yes	2.46		
09:43:09	21-11-2018	Yes	0.12		
09:43:51	06-12-2018	Yes	0.09		
52:21:02	09-11-2018	Yes	2.46		
09:44:01	16-12-2018	Yes	15.25		
09:54:19	24-12-2018	yes	0.33		

2.2 Data used

High resolution Sentinel-2 images, Table 1, acquired in the area have spectral bands capable of providing information on the quality of water. In ENVI software these images were processed specially by analyzing spectral indices on offshore, Table 2, and supervised classification on the land to determine the land cover.

Spectral indices	Formula and mathematical equation used
Colored dissolved organic matter (CDOM)	CDOM = (B3-B4)/(B3+B4)
Normalized suspended material index (NSMI)	NSMI = (B4+B3-B2)/(B4+B3+B2)
Normalized difference chlorophyll index (NDCI)	NDCI = (B5-B4)/(B5+B4)

3. Results



Figures 4. RGB natural color composite Sentinel-2 images and maps of different water indices

4. Conclusion

This study demonstrates the potential of Sentinel-2 images in monitoring and managing the quality of water in coastal environment. By analyzing sediments plumes brought in sea water by Simeto and Alcantara rivers we were able to estimate the destination of dissolved organic matter, suspended material and some bounded nutrients (such as Nitrogen or Phosphorus) leaving terrestrial environment especially in agricultural and industrial areas. When arrived in the sea their fate is influenced by sea currents and waves.

The influx of pollutant sediments in coastal area is still the major problem threatening marine ecosystem habits and its socio-economic development. In coastal environment such as East-coast of Sicily with economic benefits rely on tourism and agricultural activities, this problem has to be studied and addressed accordingly.

5. References

- H. Liu, Q. Li, T. Shi, S. Hu, G. Wu, and Q. Zhou, 2017. Application of Sentinel-2 MSI Images to retrieve Suspended particule Matter concentrations in Poyang Lake. Remote Sens. 2017, 9, 761; doi:10.3390/rs9070761.
- I. Caballero, F. Steinmetz and G. Navarro, 2018. Evaluation of the First Year of Operational Sentinel-2A Data for Retrieval of Suspended Solids in Medium- to High-Turbidity Waters. Remote Sens. 2018, 10, 982; doi:10.3390/rs10070982.
- L. Guglielmo A., F. Azzaro B., C. Baviera A., A. Bergamasco B., S. N. Bisset C., C. Brugnano A., G. Caruso B., F. Decembrini B., A. L. Garey D., A. Granata A., C. Gugliandolo A., V. Lentini A., M. A. Lo Gullo A., T. L. Maugeri A., M. Panseera A., F. Raimondo A., L. P. Rodriguez Valdes A., L. A. Smock D., A. Spano A., P. Trifilo A., J. K. Vick C., D. R. Young C., G. Zagami A., J. C. Zinner Cand R. Minutoi, 2013. Multidisciplinary ecological assessment of the Alcantara River (Sicily, Italy) using bioindicators. Marine and Freshwater Research.
- S. Lanza and G. Randazzo, 2010. A project for a protected re-nourishment of the coastline of giardini naxos, the touristic beach in the neighbouring area of taormina (north - eastern sicily). Littoral 2010, 05001 (2011) doi:10.1051/litt/201105001.
- Michelle J. D., Caroline Petus, Eduardo da Silva, Dieter Tracey, Nicholas H. Wolff, Jane Waterhouse 1 and Jon Brodie, 2015. Water quality and river plume monitoring in the great barrier reef: an overview of methods based on ocean colour satellite data. remote sens. 2015, 7, 12909-12941; doi:10.3390/rs71012909.
- S. Longhitano, A. Coella, 2007. Geomorphology, sedimentology and recent evolution of the anthropogenically modified Simeto River delta system (eastern Sicily, Italy). Sedimentary Research 194 (2007) 195-221.
- V. Catania , S. Santisi , G. Signa d , S. Vizzini , Antonio Mazzola , Simone Cappello , Michael M. Yakimov , Paola Quatrini, 2015. Intrinsic bioremediation potential of a chronically polluted marine coastal area. Marine Pollution Bulletin 99 (2015) 138-149.
- Waterhouse, J., Brodie, J., Lewis, S., Mitchell, A. Quantifying the sources of pollutants in the Great Barrier Reef catchments and the relative risk to reef ecosystems. Mar. Pollut. Bull. 2012, 65, 394-406.

Contact

Anselme Muzirafuti, Giovanni Randazzo, Stefania Lanza, Antonio Crupi, Maria Cascio, Giovanni Barreca, and Francesco Gregorio
 Università degli Studi di Messina
 Dipartimento Scienze Matematiche e Informatiche, Scienze Fisiche e Scienze della Terra (MIFT) ,Via F. Stagno d'Alcontres, 31 98166 – Sant'Agata di Messina
 ITALY.
 E-mail muzansel@gmail.com