

Mechanics of mostly non-deforming effusive eruptions: a simple model to explain dome effusion and improved constraints on the VEI 5 2008-2009 Chaitén rhyolitic eruption (Southern Andes, Chile)

Francisco Delgado¹ and Sergey Samsonov²

¹Institut de Physique du Globe de Paris

²Canada Centre for Remote Sensi

November 24, 2022

Abstract

Simple models of fluid and solid mechanics predict that the depressurization of a shallow reservoir that occurs during large effusive eruptions produces exponential trends in time series of both pressure drop and extruded volume. These models are attractive due to their simplicity and because they can explain geodetic and extruded volume data recorded at several volcanoes like at St. Helens and Cordón Caulle, regardless of their magma compositions. However, several lava and dome-forming eruptions like at Redoubt, Hekla and Santiaguito volcanoes do not show clear ground deformation coeval to lava and dome effusion despite the extrusion of at least 0.1 km³ DRE of magma. This apparent paradox can be explained by a variety of factors including deep magma sources and highly compressible magmas that leave no geodetic footprint. Here we explore the role of magma buoyancy with a reanalysis of ALOS-1, TerraSAR-X and RADARSAT-2 InSAR ground deformation and Pleiades DEM data of the VEI 5 Plinian and dome forming rhyolitic eruption of Chaitén volcano in 2008-2009. We show that almost all of the recorded ground deformation occurred during the first three weeks of the eruption, which implies that the extrusion of a rhyolitic dome (~0.8 km³ DRE) did not result in significant depressurization of a magma reservoir, despite the clear exponential trends in the extrusion data. Instead, we show that the exponential trend in the time series of extruded volume can be explained by magma ascending due its buoyancy instead of its overpressure. These results imply that ground deformation alone is not always indicative of the temporal evolution of an eruption and urges for the acquisition of denser time series of DEM data to calculate time-lapse extrusion rates.

Mechanics of mostly non-deforming effusive eruptions: a simple model to explain dome effusion and improved constraints on the VEI 5 2008-2009 Chaitén rhyolitic eruption (Southern Andes, Chile)

Mechanics of mostly non-deforming effusive eruptions: a simple model to explain dome effusion and improved constraints on the VEI 5 2008-2009 Chaitén rhyolitic eruption (Southern Andes, Chile)
 Francisco Delgado (1), Sergey Samsonov (2)
 1 Univ. de Paris, Institut de Physique du Globe de Paris, CNRS, Paris, France, Universidad de Chile, 2 Canada Centre for Mapping and Earth Observation, Natural Resources Canada

Introduction
 Analytic models of effusive eruptions predict that the series of general deformation and associated volume show exponential trends, in response to reservoir depressurization (Bardouin and Burg, 2011). These models explain general and individual volume data recorded at several volcanoes like at St. Helens (Bardouin and Burg, 2011) and Cordón Caulle (Delgado et al., 2020), despite their different magma compositions.

The 2008-2009 Chaitén eruption
 Chaitén is a rhyolite volcano in the Southern Volcanic Zone (SVZ) of the Andes of southern Chile (Fig. 1). The volcano erupted on May 2 2008 with a VEI 5 eruption. This eruption sets the first rhyolite eruption with instrumental observations and the first historical eruption of this volcano (Mayer and Lara, 2011).

INSAR and DEM data
 We use InSAR data from the ALOS, TerraSAR-X (TSX) and COSMO-SkyMed 2 (CSM2) satellites and stereo optical Pleiades data. We calculate InSAR line series for the ALOS-1 data and a 2 engine DEM (DEM) to calculate the volume of the rhyolite dome.

Kinematic source model
 We model the interregimes with a dilatant and a deflating interregime (stable volume) (produced by the level of dilatant (black lines) and deflating (grey lines) models, and difference between dilatant and deflating data (light blue lines)).

A physical model for dome effusion without reservoir depressurization
 The decreasing exponential trend in the calculated volume and the observed lack of deformation during the dome effusion cannot be explained by a model in which the dome effusion is followed by magma release to the reservoir (e.g., Siegel 2011, Arnold et al., 2017). The magma flow rate is a result of gas by the Porous flow (Jaeger and Tok, 1980).

$$Q = \frac{\sigma^2}{4\eta L} (\Delta P + (\rho_m - \rho_f)gL)$$

$$\sigma^2 \propto \Delta P$$
 and the density difference decreases exponentially then:

$$Q = \frac{\sigma^2}{4\eta L} (\rho_m - \rho_f) e^{-t/\tau} L$$

Conclusions and perspectives

- Some effusive eruptions display both of general deformation despite the existence of large volumes of lava and domes. Our model can explain the exponential trends observed during these events.
- Most of the SVZ Chaitén volume was calculated due to the magma buoyancy cost by its overpressure.
- This analysis why some effusive eruptions display clear reservoir depressurization while others do not and the role of magma buoyancy must be explained in other non-deforming systems.
- To work here influence on the absence of general deformation, we used volume DEM data with a better temporal resolution than what is currently available with TerraSAR-X, Sentinel-1, etc.

Source volume changes
 $\Delta V_+ = 3.04 \text{ km}^3$
 $\Delta V_- = 3.23 \text{ km}^3$
 $\Delta V = 3.23 \text{ km}^3$

Magma Compressibility
Explosive phase
 Magma compressibility β can be calculated from the ratio between the calculated volume ΔV and the rate of volume change \dot{V} (Borisov et al., 2008)

Navigation: LIVE SESSION | CHAT INFO | AUTHOR INFORMATION | ABSTRACT | REFERENCES | CONTACT AUTHOR | PRINT | GET POSTER

Francisco Delgado (1), Sergey Samsonov (2)

1 Université de Paris, Institut de Physique du Globe de Paris, CNRS, Paris, France. Universidad de Chile.
 2 Canada Centre for Mapping and Earth Observation, Natural Resources Canada

PRESENTED AT:



INTRODUCTION

Analytic models of effusive eruptions predict that time series of ground deformation and extruded volume show exponential trends, in response to reservoir depressurization (Anderson and Segall, 2011). These models explain geodetic and extruded volume data recorded at several volcanoes like at St. Helens (Anderson and Segall, 2011) and Cordón Caulle (Delgado et al., 2019), despite their different magma compositions.





Figure 1. Time series during the 2009 Redoubt eruption Top. Displacement (Grapenthin et al., 2013). Bottom. Extruded dome volume (Diefenbach et al., 2013).

However, eruptions like those at Redoubt 2009 (Grappenthin et al., 2013; Diefenbach et al., 2013) and Hekla 2000 (Hautmann et al., 2017; Pedersen et al., 2018) show little deformation during most of their effusive phases (Figure 1), despite the extrusion of at least 0.1 km^3 DRE of magma (Figure 1).

Here we explore the role of magma buoyancy in driving dome effusion with no ground deformation. We focus in the VEI 5 Plinian and dome forming rhyolitic eruption of Chaitén volcano in 2008-2009 with InSAR and DEM data.

THE 2008-2009 CHAITÉN ERUPTION

Chaitén is a rhyolitic caldera in the Southern Volcanic Zone (SVZ) of the Andes of southern Chile (Fig. 2). The volcano erupted on May 2 2008 with a VEI 5 eruption. This eruption was the first rhyolitic eruption with instrumental observations and the first historical eruption of this volcano (Major and Lara, 2013).







Figure 2. Top. Location of Chaitén (red triangle) in the SVZ. All the volcanoes with red and green triangles erupted during 2008 and 2015 with VEI 4-5 eruptions and significant ground deformation (Delgado et al., 2017; Delgado et al., 2019). Bottom . InSAR observations of these three eruptions

The eruption had two stages

- Explosive stage with a Plinian column during 9 days (Fig 2b), $\Delta V = 0.5\text{-}1 \text{ km}^3$ DRE (Alfano et al., 2011)
- Hybrid explosive - effusive stage from day 9 to the end of the eruption (Fig 2d) $\Delta V = 0.8 \text{ km}^3$ (Pallister et al., 2013).




Figure 3. The Chaiten eruption. b) Explosive phase, d) effusive phase (Major and Lara, 2003).

Magma was stored at depths of ~ 5 km below the surface from experimental petrology (Castro and Dingwell, 2009) and by a large dipping sill at depths of 10-20 km (Wicks et al., 2011).

INSAR AND DEM DATA

We use InSAR data from the ALOS, TerraSAR-X (TSX) and RADARSAT-2 (RS2) missions and stereo optical Pléiades data. We calculate InSAR time series for the ALOS-1 data and a 2 m/pixel DEM to calculate the volume of the rhyolitic dome.




Figure 4. Deformation during the eruption. Wrapped interferograms from pairs of SAR images (a-c), unwrapped interferograms from InSAR time series (e-h), and elevation difference between a post-eruptive Pléiades DEM and the pre-eruptive SRTM DEM (dDEM).

a-c) 0 cm during 0.7-2.5 months of the eruption.

e-f) ~37 cm of line-of-sight increase during 0-2.5 months (ascending orbit).

g) ~10 cm of subsidence during 2.5-8.6 months (ascending orbit).

h) ~45 cm of mostly subsidence during the whole eruption (descending orbit).

d) 500 m of dome height, with a volume of 1 km^3




Figure 5. Left axis shows time series of line-of-sight deformation (U_{LOS}) for the points of the same color in Fig. 3. TSX and RS2 indicate that these interferograms record no deformation given the extent of coherence loss in the data. Right axis shows the erupted volume during the effusive phase. Red points are the time series of extruded volume (Pallister et al., 2013), red dashed line is the best-fit function $\Delta V(t) = A(1 - e^{-t/\tau})$ and the orange triangle is the dome volume from the dDEM data.

InSAR indicates that 3/4 of the deformation occurred within the first three weeks of the eruption and likely during the onset of the explosive phase, with minor deformation during 2.5-8 months. The lack of deformation implies that the dome effusion was not related to a significant pressure drop in the volcano plumbing system.

KINEMATIC SOURCE MODEL

We model the Interferograms with a dike and a sill.




Figure 6. Downsampled interferograms (left column), synthetic interferograms (middle columns) predicted by the best fit dike (black lines) and sill (grey lines) models, and difference between data and synthetic data (right columns).

Source volume changes

$$\Delta V_{dike} = -0.04 \text{ km}^3$$

$$\Delta V_{sill} = -0.15 \text{ km}^3$$

$$\Delta V_r = -0.19 \text{ km}^3$$

Magma Compressibility

Explosive phase

Magma compressibility β_m can be calculated from the ratio r_v between the extruded volume ΔV_e and the source volume change ΔV_r (Mastin et al., 2008).

$$r_v = \frac{\Delta V_e}{\Delta V_r} = -\frac{\rho_m}{\rho_e} \left(1 + \frac{\beta_m}{\beta_c} \right)$$

For the explosive phase

$$\Delta V_e = 0.5-1 \text{ km}^3$$

$$\Delta V_r = -0.19 \text{ km}^3$$

$$\beta_m = 9 \times 10^{-10} - 6.4 \times 10^{-9} \text{ Pa}^{-1}$$

This is similar to other large eruptions (Kilbride et al., 2016) and is larger than for the VEI 5, 2011-2012 Cordon Caulle rhyolitic eruption (Jay et al., 2014; Delgado et al., 2019, Fig 2), with

$$\beta_m = 1-2 \times 10^{-10} \text{ Pa}^{-1}.$$

Effusive phase

The extrusion of 1 km³ dome was not related to significant deformation and reservoir volume change, therefore β_m cannot be estimated.

A PHYSICAL MODEL FOR DOME EFFUSION WITHOUT RESERVOIR DEPRESSURISATION

The decreasing exponential trend in the extruded volume and the almost lack of deformation during the dome effusion cannot be explained by a model in which the dome extrusion is balanced by magma influx in the reservoir (e.g., Segall 2013, Arnold et al., 2017).

The magma flow rate in a conduit is given by the Poiseuille law (Jaupart and Tait, 1990).

$$Q = \frac{\pi R^4}{8\eta LV(\beta_m + \beta_c)} (\Delta P + (\rho_m - \rho_r)gL)$$

If $\Delta P=0$ and the density difference decreases exponentially, then

$$Q = \frac{\pi R^4}{8\eta LV(\beta_m + \beta_c)} (\rho_m - \rho_r) e^{-t/\tau} gL$$

Integrating with respect to the time, the extruded volume is

$$V_e = \frac{\pi R^4}{8\eta LV(\beta_m + \beta_c)} (\rho_m - \rho_r) gL \tau (1 - e^{-t/\tau})$$

Magma ascent by an exponential decrease in its density difference (magma buoyancy) also results in an exponential trend in the extruded volume, like in Fig 5.

The model can explain other time series with near exponential trends like at Redoubt 2009 (Fig 1).

CONCLUSIONS AND PERSPECTIVES

- Some effusive eruptions display lack of ground deformation despite the extrusion of large volumes of lava and domes. Our model can explain the exponential trends observed during these events.
- Most of the 1 km³ Chaiten dome was extruded due to the magma buoyancy, not by its overpressure.
- It is unclear why some effusive eruptions do display clear reservoir depressurization while others do not and the role of magma buoyancy must be explored in other non-deforming events.
- To track lava effusion in the absence of ground deformation, we need satellite dDEM data with a better temporal resolution than what is currently possible with TanDEM-X, Pléiades, SPOT, and WorldView.

Acknowledgements

CNES for a postdoctoral fellowship to F.D., Arthur Delorme (IPGP) for his help with the Pléiades processing, and Tara Shreve (IPGP) and Kyle Anderson (U.S. Geological Survey) for discussions.

ALOS-1 data were provided by the Alaska Satellite Facility. TerraSAR-X data were provided by DLR. RADARSAT-2 data were provided by CSA and MDA. Pléiades data were provided by CNES and Airbus. Pléiades and TerraSAR-X data were distributed through the CEOS Volcano Demonstrator.

AUTHOR INFORMATION

Francisco Delgado, delgado@ipgp.fr

ABSTRACT

Simple models of fluid and solid mechanics predict that the depressurization of a shallow reservoir that occurs during large effusive eruptions produces exponential trends in time series of both pressure drop and extruded volume. These models are attractive due to their simplicity and because they can explain geodetic and extruded volume data recorded at several volcanoes like at St. Helens and Cordón Caulle, regardless of their magma compositions. However, several lava and dome-forming eruptions like at Redoubt, Hekla and Santiaguito volcanoes do not show clear ground deformation coeval to lava and dome effusion despite the extrusion of at least 0.1 km³ DRE of magma. This apparent paradox can be explained by a variety of factors including deep magma sources and highly compressible magmas that leave no geodetic footprint. Here we explore the role of magma buoyancy with a reanalysis of ALOS-1, TerraSAR-X and RADARSAT-2 InSAR ground deformation and Pleiades DEM data of the VEI 5 Plinian and dome forming rhyolitic eruption of Chaitén volcano in 2008-2009. We show that almost all of the recorded ground deformation occurred during the first three weeks of the eruption, which implies that the extrusion of a rhyolitic dome (~0.8 km³ DRE) did not result in significant depressurization of a magma reservoir, despite the clear exponential trends in the extrusion data. Instead, we show that the exponential trend in the time series of extruded volume can be explained by magma ascending due its buoyancy instead of its overpressure. These results imply that ground deformation alone is not always indicative of the temporal evolution of an eruption and urges for the acquisition of denser time series of DEM data to calculate time-lapse extrusion rates.

REFERENCES

- Alfano, F., C. Bonadonna, A. C. Volentik, C. B. Connor, S. F. Watt, D. M. Pyle, and L. J. Connor (2011), Tephra stratigraphy and eruptive volume of the May, 2008, Chaiten eruption, Chile, *Bulletin of Volcanology*, doi:10.1007/s00445-010-0428-x.
- Anderson, K., and P. Segall (2011), Physics-based models of ground deformation and extrusion rate at effusively erupting volcanoes, *Journal of Geophysical Research-Solid Earth*, 116, doi:10.1029/2010jb007939.
- Arnold, D. W. D., J. Biggs, K. Anderson, S. V. Vargas, G. Wadge, S. K. Ebmeier, M. F. Naranjo, and P. Mothes (2017), Decaying lava extrusion rate at El Reventador Volcano, Ecuador measured using high-resolution satellite radar, *Journal of Geophysical Research: Solid Earth*, pp. n/a–n/a, doi:10.1002/2017JB014580.
- Castro, J. M., and D. B. Dingwell (2009), Rapid ascent of rhyolitic magma at Chaiten volcano, Chile, *Nature*, doi:10.1038/nature08458.
- Delgado, F., M.E. Pritchard, S. Ebmeier, P. Gonzalez and L. Lara (2017), Recent unrest (2002–2015) imaged by space geodesy at the highest risk Chilean volcanoes: Villarrica, Llaima, and Calbuco (Southern Andes), *Journal of Volcanology and Geothermal Research*, 524, 115,736, doi:10.1016/J.EPSL.2019.115736.
- Delgado, F., J. Kubanek, K. Anderson, P. Lundgren, and M. Pritchard (2019), Physicochemical models of effusive rhyolitic eruptions constrained with InSAR and DEM data: A case study of the 2011-2012 Cordón Caulle eruption, *Earth and Planetary Science Letters*, 344, 270-288, doi:10.1016/j.jvolgeores.2017.05.020.
- Diefenbach, A. K., K. F. Bull, R. L. Wessels, and R. G. McGimsey (2013), Photogrammetric monitoring of lava dome growth during the 2009 eruption of Redoubt Volcano, *Journal of Volcanology and Geothermal Research*, 259, 308–316, doi:10.1016/j.jvolgeores.2011.12.009.
- Grapenthin, R., J. T. Freymueller, and A. M. Kaufman (2013), Geodetic observations during the 2009 eruption of Redoubt Volcano, Alaska, *Journal of Volcanology and Geothermal Research*, doi:10.1016/j.jvolgeores.2012.04.021.
- Hautmann, S., I. S. Sacks, A. T. Linde, and M. J. Roberts (2017), Magma buoyancy and volatile ascent driving autocyclic eruptivity at Hekla Volcano (Iceland), *Geochemistry, Geophysics, Geosystems*, doi:10.1002/2017GC007061.
- Jaupart, C., and S. Tait (1990), Dynamics of eruptive phenomena, *Reviews in Mineralogy*, 24, 213–238.
- Jay, J., F. Costa, M. Pritchard, L. Lara, B. Singer, and J. Herrin (2014), Locating magma reservoirs using InSAR and petrology before and during the 2011-2012 Cordón Caulle silicic eruption, *Earth and Planetary Science Letters*, 403, 463, doi:10.1016/j.epsl.2014.07.021.
- Kilbride, B. M. C., M. Edmonds, and J. Biggs (2016), Observing eruptions of gas-rich compressible magmas from space, *Nature Communications*, doi:10.1038/ncomms13744
- Major, J. J., and L. E. Lara (2013), Overview of Chaiten Volcano, Chile, and its 2008-2009 eruption, *Andean Geology*, 40(2), 196–215, doi:10.5027/andgeoV40n2-a01.
- Mastin, L. G., E. Roeloffs, N. M. Beeler, and J. E. Quick (2008), Constraints on the size, overpressure, and volatile content of the Mount St. Helens magma system from geodetic and dome-growth measurements during the 2004–2006+ eruption, in *A volcano rekindled; the renewed eruption of Mount St. Helens, 2004- 2006*, vol. 1750, edited by D. R. Sherrod, W. E. Scott, and P. H. Stauffer, p. 856, U.S. Geological Survey Professional Paper.
- Pallister, J. S., A. K. Diefenbach, W. C. Burton, J. Munoz, J. P. Griswold, L. E. Lara, J. B. Lowenstern, and C. E. Valenzuela (2013a), The Chaiten rhyolite lava dome: Eruption sequence, lava dome volumes, rapid effusion rates and source of the rhyolite magma, *Andean Geology*, 40 (2), 277–294, doi:10.5027/andgeoV40n2-a06.
- Pedersen, G. B. M., J. M. C. Belart, E. Magnússon, O. K. Vilmundardóttir, F. Kizel, F. S. Sigurmundsson, G. Gísladóttir, and J. A. Benediktsson (2018), Hekla Volcano, Iceland, in the 20th Century: Lava Volumes, Production Rates, and Effusion Rates, *Geophysical Research Letters*, 45(4), 1805–1813, doi:10.1002/2017GL076887.

Segall, P. (2013), Volcano deformation and eruption forecasting, *Remote Sensing of Volcanoes and Volcanic Processes: Integrating Observation and Modelling*, 380, 85–106, doi:10.1144/sp380.4.

Wicks, C., J. C. de la Llera, L. E. Lara, and J. Lowenstern (2011), The role of dyking and fault control in the rapid onset of eruption at Chaiten volcano, Chile, *Nature*, 478(7369), 374–+, doi:10.1038/nature10541.