

Morphological phenotyping of hooked hairs in *Phaseolus Vulgaris*

Ankita Roy¹, Addison Bralick², Peter Pietrzyk¹, and Alexander Bucksch³

¹Department of Plant Biology

²Department of Chemistry, University of Georgia

³Department of Plant Biology, University of Georgia

November 21, 2022

Abstract

We quantify the shape of hooked hairs which is a newly observed phenotype of epidermal cell extensions [1] in the common bean genotype L88-57 (*Phaseolus vulgaris*). The hooked hairs emerge below-ground before the root hairs and have a distinct ‘hooking’ morphology. We generated a dataset capturing their full distribution under the microscope within 3-5 days of germination. We quantify their shape by a novel computational pipeline that can automatically phenotype morphology. Our phenotyping pipeline quantifies traits like length, curvature, perimeter, area, and ‘hooking.’ Our objective is to quantify the response of hooked hairs to nutrient-stress to determine their function in common bean during early development. We used the pipeline for analyzing our dataset of hydroponically grown beans and observed statistically significant responses compared to the control for length, curvature, perimeter, and area to nitrogen ($p < 0.001^{**}$) and phosphorus ($p < 0.001^{**}$) stress treatments. The calculation of ‘hooking’ for our dataset is still ongoing. We are simultaneously developing a landmark-free method for the two-dimensional shape analysis of our dataset and believe that our phenotyping efforts will enable the high-throughput analysis of morphological root hair traits for any plant species.

Hosted file

essoar.10508329.1.docx available at <https://authorea.com/users/243173/articles/597331-morphological-phenotyping-of-hooked-hairs-in-phaseolus-vulgaris>

The morphological phenotyping of hooked hairs in *Phaseolus Vulgaris*

Ankita Roy¹, Addison Bralick², Peter Pietrzyk¹ Alexander Bucksch^{1,3,4}

Plant Biology¹, Chemistry², Warnell School of Forestry and Natural Resources³,
Inst. Of Bioinformatics⁴

We quantify the shape of hooked hairs which is a newly observed phenotype of epidermal cell extensions [1] in the common bean genotype L88-57 (*Phaseolus vulgaris*). The hooked hairs emerge below-ground before the root hairs and have a distinct 'hooking' morphology. We generated a dataset capturing their full distribution under the microscope within 3-5 days of germination. We quantify their shape by a novel computational pipeline that can automatically phenotype morphology. Our phenotyping pipeline quantifies traits like length, curvature, perimeter, area, and 'hooking.' Our objective is to quantify their response to nutrient-stress to determine the function of hooked hairs in common bean during early development. We used the pipeline for analyzing our dataset of hydroponically grown beans and observed statistically significant responses compared to the control for length, curvature, perimeter, and area to nitrogen ($p < 0.001^{**}$) and phosphorus ($p < 0.001^{**}$) stress treatments. The calculation of 'hooking' for our dataset is still ongoing. We are simultaneously developing a landmark-free method for the two-dimensional shape analysis of our dataset and believe that our phenotyping efforts will enable the high-throughput analysis of morphological root hair traits for any plant species.

[1] A. Roy, A. Bucksch. Root Hairs vs. Trichomes: Not everyone is straight! Current Opinion in Plant Biology, 2021 (under revision)

Keywords: morphology, root hair, hooking, *Phaseolus vulgaris*, phenotyping, landmark-free, shape analysis, trait