

# Role of Nuclear Factor of Activated T cells 2 (NFATc2) in allergic asthma

Marielena Jakobi<sup>1</sup>, Alexander Kiefer<sup>2</sup>, Hooman Mirzakhani<sup>3</sup>, Manfred Rauh<sup>1</sup>, Theodor Zimmermann<sup>4</sup>, PARASKEVI XEPAPADAKI<sup>5</sup>, Barbara Stanic<sup>6</sup>, Mubeccel Akdis<sup>7</sup>, Nikolaos Papadopoulos<sup>8</sup>, Benjamin Raby<sup>3</sup>, Scott Weiss<sup>9</sup>, and Susetta Finotto<sup>1</sup>

<sup>1</sup>University Hospital Erlangen

<sup>2</sup>FAU-Erlangen

<sup>3</sup>Partners Harvard Medical International

<sup>4</sup>Friedrich-Alexander-Universität Erlangen-Nürnberg

<sup>5</sup>National and Kapodistrian University of Athens

<sup>6</sup>SIAF

<sup>7</sup>University of Zürich

<sup>8</sup>Second Pediatric Clinic

<sup>9</sup>Channing Division of Network Medicine

June 22, 2020

## Hosted file

12012020.NFATc12 Paper sf\_HM2\_MJ\_STW.doc available at <https://authorea.com/users/335722/articles/461533-role-of-nuclear-factor-of-activated-t-cells-2-nfatc2-in-allergic-asthma>




Figure 1




Figure 2