

Successful revascularization with percutaneous coronary intervention using a combination of the subintimal transcatheter withdrawal technique and coronary artery fenestration for spontaneous coronary artery dissection

Shintaro Matsuura¹, Kanichi Otowa¹, Michiro Maruyama¹, and Kazuo Usuda¹

¹Toyama Prefectural Central Hospital

July 7, 2021

Abstract

We present a case of successful revascularization for spontaneous coronary artery dissection (SCAD) using coronary artery fenestration followed by the subintimal transcatheter withdrawal (STRAW) technique. The combination of the STRAW technique and coronary artery fenestration with cutting balloon angioplasty could be a new treatment option for SCAD patients requiring revascularization.

Successful revascularization with percutaneous coronary intervention using a combination of the subintimal transcatheter withdrawal technique and coronary artery fenestration for spontaneous coronary artery dissection

Shintaro Matsuura, MD; Kanichi Otowa, MD, PhD; Michiro Maruyama, MD, PhD; Kazuo Usuda, MD, PhD
Department of Cardiology, Toyama Prefectural Central Hospital, Toyama, Japan

Shintaro Matsuura, Department of Cardiology, Toyama Prefectural Central Hospital, Toyama, 2-2-78 Nishinagae, Toyama-shi, Toyama-ken 930-8550, Japan.

E-mail address: *mat.s.86m23@gmail.com*

Word count: 1984 words

ABSTRACT

We present a case of successful revascularization for spontaneous coronary artery dissection (SCAD) using coronary artery fenestration followed by subintimal transcatheter withdrawal (STRAW) technique. The combination of the STRAW technique and coronary artery fenestration with cutting balloon angioplasty could be a new treatment option for SCAD patients requiring revascularization.

KEY CLINICAL MESSAGE

The combination of the STRAW technique and coronary artery fenestration using a cutting balloon could be effective in SCAD patients, especially with dissection to the distal end of the coronary artery.

KEYWORDS

Coronary Balloon Angioplasty, Percutaneous Coronary Intervention, Myocardial Infarction, Intravascular Ultrasonography, Computed Tomography Angiography

1. INTRODUCTION

Spontaneous coronary artery dissection (SCAD) has emerged as an important cause of acute coronary syndrome, particularly in young woman.¹⁻³ Although recent studies have demonstrated that conservative therapy is appropriate for the initial management of SCAD, revascularization should be considered in a patient with ongoing ischemia, high-risk anatomy, or hemodynamic instability.^{4,5} Percutaneous coronary intervention (PCI) for SCAD treatment is associated with an increased risk of complications and suboptimal outcomes; however, several case reports have shown the efficacy of coronary artery fenestration with cutting balloon angioplasty.⁶⁻⁸ Here, we present the case of a patient with SCAD whose ischemia was not corrected with coronary artery fenestration alone but was successful revascularized by the additional use of the subintimal transcatheter withdrawal (STRAW) technique.^{9,10}

2. CASE PRESENTATION

A 31-year-old woman with a history of smoking but no prior coronary artery disease was admitted to our hospital at 20 weeks of gestation for impending preterm labor. At 36 weeks of gestation, a cesarean section was performed. However, 2 days thereafter, she developed chest pain with anterior ST-segment elevation on electrocardiogram, indicative of acute myocardial infarction. Emergency coronary angiography (CAG) showed subtotal occlusion in the middle part of the left anterior descending artery (LAD), and there appeared to be a contrast delay distally. In order to confirm the cause of subtotal occlusion, an IVUS (Opti-Cross HD, Boston Scientific, Inc., Marlborough, MA, USA) examination was performed (Figure 1). IVUS demonstrated that the false lumen compressed the true lumen from the middle to distal part of the LAD. Based on the findings of the CAG and IVUS, she was diagnosed with type-1 SCAD. The IVUS catheter was advanced as far distally as possible; however, the distal end of the false lumen could not be identified. We decided to perform primary PCI because of ongoing chest pain and persistent ST-segment elevation. First, we attempted coronary artery fenestration using cutting balloon angioplasty. A 3.5-mm cutting balloon (Wolverine, Boston Scientific, Inc., Marlborough, MA, USA) was advanced into the distal LAD where the IVUS catheter could be advanced and then dilated. The IVUS findings after cutting balloon angioplasty showed the creation of some reentries; however, the true lumen at the distal LAD was still compressed by the hematoma in the false lumen. In order to make an additional fenestration at a more distal site, we attempted to advance a 2.5-mm cutting balloon (Wolverine, Boston Scientific, Inc., Marlborough, MA, USA); however, this attempt was unsuccessful. Therefore, we then applied the STRAW technique. A second guidewire (SUOH03, ASAHI INTECC CO., LTD, Aichi, Japan) was intentionally advanced into the false lumen under real-time IVUS guidance, and a microcatheter (FINECROSS GT, TERUMO Corp., Tokyo, Japan) was also advanced into the false lumen at the far distal part of the LAD. After the guidewire was removed, we aspirated blood from the tip of the microcatheter while gradually moving the microcatheter position to the proximal side; then, ST-segment elevation improved, and chest pain disappeared (Figure 2). We attempted to advance a 2.5-mm cutting balloon again, and it was able to advance more distally than before the STRAW technique; thereafter, coronary artery fenestration was performed at the distal LAD. The IVUS findings after the STRAW technique and the additional coronary artery fenestration at the distal LAD showed an enlarged true lumen and multiple reentries formations. Although there were several stenoses in the middle part of the LAD due to compression by the false lumen, coronary artery blood flow improved to Thrombolysis in Myocardial Infarction (TIMI) flow grade 2 (Figure 3,4). Considering the absence of ongoing ischemia and the patient's age and gender, we decided to complete the procedure without stenting. The post-procedural course was good and uneventful till 10 months thereafter. We followed up the patient with coronary computed tomographic angiography; the flap and the stenosis at the middle part of the LAD disappeared after 10 months of PCI (Figure 5).

3. DISCUSSION

Previous observational studies have consistently shown that PCI for SCAD treatment is associated with an increased risk of complications and suboptimal outcomes.¹⁻⁵ However, owing to ongoing ST-segment elevation, we performed primary PCI for this patient.

Some case reports have shown the efficacy of coronary artery fenestration with cutting balloon angioplasty to decompress the hematoma and avoiding longitudinal extension.⁶⁻⁸ We also attempted this technique; however, our attempt was unsuccessful. We believe the following reasons were responsible for the failure of this attempt: 1) the cutting balloon could not advance sufficiently distally because the true lumen was severely compressed by the false lumen and 2) the dissection had reached the distal end of the LAD; therefore, coronary artery fenestration could not be made at the distal end of the false lumen, and the compression of the true lumen was not released by the remaining hematoma in the false lumen. In order to overcome these problems, we used the STRAW technique.^{9,10}

The STRAW technique was reported as an option to decompress the distal true lumen and enable successful reentry for antegrade in chronic total occlusion cases. In the original method, an over-the-wire balloon was positioned in the proximal vessel at the site of origin of the dissection plane. Following balloon inflation, a 10-mL syringe was attached to the proximal lumen port on the negative pressure.¹⁰ In addition, the case report of iatrogenic coronary artery dissection from our hospital has shown that aspiration using a microcatheter could reduce the hematoma in the false lumen.¹¹ In this case, we performed microcatheter aspiration to successfully improve the ongoing ischemia. Furthermore, it became possible to deliver a cutting balloon to a more distal part of the LAD, and we succeeded in performing the coronary artery fenestration at a sufficiently distal side. We believe that aspiration using a microcatheter offers the advantage of safe advancement of the aspiration device to the distal part of the false lumen. However, if microcatheter aspiration is insufficient, it may be helpful to perform aspiration during balloon occlusion in the proximal vessel, referring to the original method.

PCI in SCAD should focus on restoring the TIMI flow grade and stabilizing the patient clinically rather than restoring normal coronary artery structure.² Moreover, systematic studies have been reported on whether stents should be used in PCI for SCAD. In this case, we decided to perform stent-less PCI because ST-segment elevation and TIMI flow grade improved, and the patient was a young woman. Observational data have shown that spontaneous healing occurs in most SCAD patients³, and in the present case, coronary computed tomographic angiography still showed healing of the SCAD lesion.

This new PCI strategy that combined the STRAW technique and coronary artery fenestration using a cutting balloon appeared effective in SCAD patients; however, a more large-scale study is required to confirm the safety and efficacy of this method.

4. CONCLUSION

A combination of the STRAW technique and coronary artery fenestration with cutting balloon angioplasty could be effective in SCAD patients, with dissection to the distal end of the coronary artery. However, a more large-scale evaluation is required to confirm the safety and efficacy of this method.

CONFLICT OF INTEREST

The authors declare that there is no conflict of interest.

AUTHOR CONTRIBUTIONS

SM: involved in management of the patient and wrote the article. KO, MM, and KU: involved in critically reviewed and revision of the manuscript. All authors: approved the final version of the case report for submission to *Clinical Case Reports*.

ETHICAL STATEMENT

Published with written consent of the patient.

REFERENCES

1. Kim ESH. 2020. Spontaneous Coronary-Artery Dissection. *N Engl J Med* . 383(24):2358-2370.

2. Hayes SN, Tweet MS, Adlam D, et al. 2020. Spontaneous Coronary Artery Dissection: JACC State-of-the-Art Review. *J Am Coll Cardiol* . 76(8):961-984.
3. Hayes SN, Kim ESH, Saw J, et al. 2018. Spontaneous Coronary Artery Dissection: Current State of the Science: A Scientific Statement From the American Heart Association. *Circulation* . 137(19):e523-e557.
4. Lettieri C, Zavalloni D, Rossini R, et al. 2015. Management and Long-Term Prognosis of Spontaneous Coronary Artery Dissection. *Am J Cardiol* . 116(1):66-73.
5. Tweet MS, Eleid MF, Best PJ, et al. 2014. Spontaneous coronary artery dissection: revascularization versus conservative therapy. *Circ Cardiovasc Interv* . 7(6):777-786.
6. Bresson D, Calcaianu M, Lawson B, Jacquemin L. 2019. Coronary artery fenestration as rescue management of intramural haematoma with luminal compression. *Catheter Cardiovasc Interv* . 94(1):E17-E19.
7. Ito T, Shintani Y, Ichihashi T, Fujita H, Ohte N. 2017. Non-atherosclerotic spontaneous coronary artery dissection revascularized by intravascular ultrasonography-guided fenestration with cutting balloon angioplasty. *Cardiovasc Interv Ther* . 32(3):241-243.
8. Motreff P, Barber-Chamoux N, Combaret N, Souteyrand G. 2015. Coronary artery fenestration guided by optical coherence tomography before stenting: new interventional option in rescue management of compressive spontaneous intramural hematoma. *Circ Cardiovasc Interv* . 8(4):e002266.
9. Wu EB, Brilakis ES, Lo S, et al. 2020. Advances in CrossBoss/Stingray use in antegrade dissection reentry from the Asia Pacific Chronic Total Occlusion Club. *Catheter Cardiovasc Interv* . 96(7):1423-1433.
10. Smith EJ, Di Mario C, Spratt JC, et al. 2015. Subintimal TRANscatheter Withdrawal (STRAW) of hematomas compressing the distal true lumen: a novel technique to facilitate distal reentry during recanalization of chronic total occlusion (CTO). *J Invasive Cardiol* . 27(1):E1-4.
11. Nagata Y, Maruyama M, Aburadani I, Hirazawa M, Mayumi T, Usuda K. 2015. A case of delayed occlusive dissection of the right coronary artery during coronary intervention of the left anterior descending artery. *Cardiovasc Interv Ther* . 30(2):155-161.

FIGURE LEGENDS

FIGURE 1 Baseline left coronary angiography and IVUS images. The angiography (left) shows subtotal occlusion in the middle part of the LAD, and there appeared to be a contrast delay distally. IVUS images from middle to distal LAD (right) demonstrate the false lumen compressing the true lumen. t = true lumen, f = false lumen.

FIGURE 2 The STRAW technique using a microcatheter. The first guidewire (arrowhead) was placed into the true lumen, and the microcatheter (arrow) advanced into the false lumen at the far distal part of the LAD. With the application of negative pressure using a syringe, the hematoma was aspirated from the tip of the microcatheter, while gradually moving the microcatheter position to the proximal part.

FIGURE 3 Final coronary angiography and IVUS images. The angiography (left) shows improvement in the coronary artery blood flow from TIMI 1 to 2. In addition, small side branches at distal LAD can be observed. IVUS images (right) demonstrate multiple connections between the true lumen and the false lumen, with true lumen enlargement (D-F). t = true lumen, f = false lumen.

FIGURE 4 Comparison of the IVUS images during each procedure (baseline, pre-STRAW, final). IVUS images recorded at approximately the same site are placed side by side. t = true lumen, f = false lumen.

FIGURE 5 Coronary computed tomographic angiography images for follow up after PCI. Each image shows multiplanar reconstructed images of LAD, (A) 1 week, (B) 2 weeks, (C) 2 months, (D) 10 months after PCI. The flap and coronary artery stenosis in the middle of the LAD (arrow) gradually disappeared.





