

Factors associated with early uptake of COVID-19 vaccination among healthcare workers in Azerbaijan, 2021

Jason Doran¹, Nabil Seyidov², Samir Mehdiyev², Girogia Gon³, Esther Kissling⁴, Trent Herdman¹, Javahir Suleymanova³, Ana Paula Couthino Rehse³, Richard Pebody³, Mark Katz³, and Gahraman Hagverdiyev²

¹UK Health Security Agency

²Centre of Public Health and Reforms of the Ministry of Health

³World Health Organization Regional Office for Europe

⁴EpiConcept

February 27, 2022

Abstract

We evaluated uptake and factors associated with COVID-19 vaccination among health workers (HWs) in Azerbaijan. Among 1,575 HWs, 73% had received at least one dose and 67% received two doses; all received CoronaVac. Factors associated with vaccination uptake included no previous COVID-19 infection, older age, belief in the vaccine's safety, previous vaccination for influenza, having patient-facing roles, and good or excellent health by self-assessment. These findings could inform strategies to increase vaccination uptake as the campaign continues.

Hosted file

Final manuscript.docx available at <https://authorea.com/users/462609/articles/558006-factors-associated-with-early-uptake-of-covid-19-vaccination-among-healthcare-workers-in-azerbaijan-2021>

Hosted file

Figures.docx available at <https://authorea.com/users/462609/articles/558006-factors-associated-with-early-uptake-of-covid-19-vaccination-among-healthcare-workers-in-azerbaijan-2021>