

Updating the dual C and O isotope – gas exchange model: A concept to understand plant responses to the environment and its implications for tree rings

Rolf Siegwolf¹, Marco Lehmann¹, Gregory Goldsmith², Olga Churakova (Sidorova)³, Cathleen Mirande-Ney⁴, Galina Timofeeva⁵, Rosemarie Weigt⁴, and Matthias Saurer¹

¹Swiss Federal Institute for Forest Snow and Landscape Research WSL

²Chapman University Schmid College of Science and Technology

³Siberian Federal University

⁴Paul Scherrer Institute Laboratory of Atmospheric Chemistry

⁵Swiss Federal Institute of Technology

November 16, 2022

Abstract

The combined study of C and O isotopes in plant organic matter has emerged as a powerful tool for understanding plant functional responses to environmental change. The approach relies on established relationships between leaf gas exchange and isotopic fractionation to derive a series of model scenarios that can be used to infer changes in photosynthetic assimilation and stomatal conductance driven by changes in environmental parameters (CO₂, water availability, air humidity, temperature, nutrients). We review the mechanistic basis for a conceptual model, in light of recently published research, and discuss where isotopic observations don't match our current understanding of plant physiological response to environment. We demonstrate that 1) the model was applied successfully in many, but not all studies, 2), while originally conceived for leaf isotopes, the model has been applied extensively to tree ring isotopes in the context of tree physiology and dendrochronology. Where isotopic observations deviate from physiologically plausible conclusions, this mismatch between gas-exchange and isotope response provides valuable insights on underlying physiological processes. Overall, we found that isotope responses can be grouped into situations of increasing resource limitation versus higher resource availability. The dual isotope model helps to interpret plant responses to a multitude of environmental factors.

Updating the dual C and O isotope – gas exchange model: A concept to understand plant responses to the environment and its implications for tree rings

Rolf T.W. Siegwolf¹(orcid.org/0000-0002-0249-0651), Marco M. Lehmann¹, Gregory R. Goldsmith³, Olga V. Churakova (Sidorova)⁴, Cathleen Mirande-Ney², Galina Timofeeva^{2,5}, Rosemarie B. Weigt², Matthias Saurer¹

¹Forest Dynamics, Swiss Federal Institute for Forest, Snow and Landscape Research WSL, 8903 Birmensdorf, Switzerland

²Ecosystem Fluxes Group, Laboratory for Atmospheric Chemistry, Paul Scherrer Institute, Forschungsstrasse 111, 5232 Villigen PSI, Switzerland

³Schmid College of Science and Technology, Chapman University, Orange, California 92866, USA

⁴Siberian Federal University, Institute of Ecology and Geography, 660041 Krasnoyarsk, Russian Federation

⁵ ETH Alumni Association, Rämistrasse 101, 8092 Zürich, Switzerland

Correspondence: Forest Dynamics, Swiss Federal Institute for Forest, Snow and Landscape Research WSL, 8903 Birmensdorf, Switzerland Email: rolf.siegwolf@wsl.ch

Abstract

The combined study of C and O isotopes in plant organic matter has emerged as a powerful tool for understanding plant functional responses to environmental change. The approach relies on established relationships between leaf gas exchange and isotopic fractionation to derive a series of model scenarios that can be used to infer changes in photosynthetic assimilation and stomatal conductance driven by changes in environmental parameters (CO₂, water availability, air humidity, temperature, nutrients). We review the mechanistic basis for a conceptual model, in light of recently published research, and discuss where isotopic observations don't match our current understanding of plant physiological response to environment. We demonstrate that 1) the model was applied successfully in many, but not all studies, 2), while originally conceived for leaf isotopes, the model has been applied extensively to tree ring isotopes in the context of tree physiology and dendrochronology. Where isotopic observations deviate from physiologically plausible conclusions, this mismatch between gas-exchange and isotope response provides valuable insights on underlying physiological processes. Overall, we found that isotope responses can be grouped into situations of increasing resource limitation versus higher resource availability. The dual isotope model helps to interpret plant responses to a multitude of environmental factors.

Introduction

Plants are subject to a wide range of environmental impacts and even withstand environmental extremes, e.g. in temperature (e.g. -60°C to + 45°C in boreal forests), water availability (e.g. prolonged months of drought in arid regions), nutrient availability (e.g. low nutrients in mountainous regions) and disturbance (e.g. storm damage, fires or land use change). Their ability to acclimate is therefore critical for their survival. To describe the nature of responses and acclimatization, detailed and comprehensive plant physiological measurements are necessary; however, the intensive nature of these measurements often limits the extent to which they can be applied across space and time. Moreover, numerous physiological measurements of plant responses to the environment (e.g. classical approaches such as plant gas exchange, light interception, sap flux measurements, etc.) cannot be applied retrospectively. Here, the analysis of stable isotope ratios of plant organic matter can provide additional insights by effectively tracing and integrating physiological processes over both time and space (Dawson & Siegwolf, 2007).

The use of stable isotopes as a proxy for understanding the relationships between environment and plant function relies on a mechanistic understanding of the drivers of isotopic fractionation (Dawson *et al.*, 2002). For instance, changes in environmental conditions and related effects on photosynthesis and transpiration result in predictable carbon and oxygen isotope ratios that are imprinted on photosynthetic assimilates (e.g. carbohydrates). Although there may be further possible isotope fractionations between primary and secondary metabolites, these isotopic “fingerprints” are ultimately reflected in the biomass. The isotopic ratio of assimilates can thus be used as a record of the interaction between the plant and its environment over the lifetime of the specific tissue (e.g. leaf, roots or wood).

In the last three decades, the application of stable carbon and oxygen isotope ratios (¹³C/¹²C and ¹⁸O/¹⁶O) in plant ecophysiological research has become common for studies of short- and long-term effects of environmental changes on vegetation (Farquhar *et al.*, 1982; Griffiths *et al.*, 1997; Ehleringer *et al.*, 2000; Dawson & Siegwolf, 2007). As each isotope ratio reflects particular physiological responses caused by specific environmental changes, the use of both carbon and oxygen isotope ratios provides highly complementary information that improves the potential for interpreting cause and effect. Consequently, the two isotope ratios have been combined and linked with leaf-level CO₂ and H₂O gas exchange in a formal conceptual model that facilitates inferences about changes in assimilation and stomatal conductance in response to the environment (Scheidegger *et al.*, 2000).

The dual carbon and oxygen isotope model formally proposed by Scheidegger *et al.*, (2000), hereafter described as the DI-model, has been subject to a diversity of applications. With these applications have come new insights into the strengths and weaknesses of the model that have led to advances in our understanding of plant function. To highlight these insights, we carried out a quantitative review of studies that have engaged the DI-model to date. Specifically, we aimed to 1) review the mechanistic basis of the model, 2) determine the nature and extent of its application to date in different fields, and 3) review studies that provide data on the mechanistic relationships between plant physiology and carbon/oxygen isotopes to strengthen our ability to interpret the model.

The dual carbon and oxygen isotope model for C3 plants

The purpose of the DI-model is to make inferences regarding the responses of net assimilation (A_{net} , Larcher, 2003) and stomatal conductance (g_s) to environmental variables based on a dual-isotope (carbon and oxygen) response. Since 1) both carbon and plant oxygen isotope ratios are modified during CO_2 and H_2O gas exchange in the same leaf, but by different processes (photosynthesis and transpiration), and 2) both carbon and oxygen are incorporated into organic matter equally, it provides a means by which to distinguish whether a change in isotope ratios is the result of a change in photosynthesis or in stomatal conductance. Current theory on carbon and oxygen isotope fractionations is summarized in Breakout Boxes 1 and 2.

In general, two conditions are compared so that it is possible to draw an arrow from condition A to condition B both with respect to the isotope and gas-exchange observations (Figure 1). The conditions A and B may represent two treatments in a controlled experiment, two functional groups (e.g. broad leaved plants and conifers) in the same ecosystem, leaf or stem samples from two positions in a canopy, one species compared between two sites, the same plant at two time points (temporal development), etc. Note that the red arrows in Figure 1 refer to a range of plant responses, indicating the plasticity of plants responding to environmental impacts within the indicated range.

First, the carbon and oxygen isotope ratios need to be measured in an organic compound (e.g. bulk leaf, cellulose, sugars, wood or root material) of the plants under investigation. Proper selection of the plant and compound is important, as it has consequences for the temporal scale at which the physiological processes are integrated. For instance, the time integrated by different compounds is usually longest for cellulose, intermediate for bulk, smaller for water-extracted compounds and shortest for isolated sucrose (Lehmann *et al.*, 2017). Second, the “input-scheme” of the DI-model, a plot of $\delta^{18}\text{O}$ data versus $\delta^{13}\text{C}$ data comparing the conditions of interest, can be drawn. Alternatively, the oxygen enrichment in the organic compound relative to source water ($\Delta^{18}\text{O}$) can be presented. This is advantageous in cases where the $\delta^{18}\text{O}$ of source water differs between treatment conditions A compared to B; given that this is not a leaf-level physiological effect, it could ultimately result in erroneous model interpretations (Roden & Siegwolf, 2012). Similarly, it is also possible to present the carbon isotope discrimination ($\Delta^{13}\text{C}$) instead of original $\delta^{13}\text{C}$ data, particularly in cases where the $\delta^{13}\text{C}$ of source CO_2 is not constant (e.g. atmospheric CO_2 enrichment studies, or retrospective analyses of organic matter, tree ring, peat bogs). Note that in the case of $\Delta^{13}\text{C}$, the ^{13}C arrows in the dual-isotope plot will be pointing inverse inclination as compared to using (δ) original data, whereas for $\Delta^{18}\text{O}$ the direction of the arrows remains unchanged. Here, we assume the presentation of $\delta^{18}\text{O}$ or $\Delta^{18}\text{O}$ vs. $\delta^{13}\text{C}$ (Figure 1).

Deducing the $A_{\text{net}} - g_s$ response from the isotope pattern can be explained as a 3-step-process: 1) a change in c_i/c_a (the ratio of leaf internal to ambient CO_2) is deduced from a change in the $\delta^{13}\text{C}$, 2) a change e.g. in the relative humidity (RH) or vapour pressure deficit (VPD) is causing a change in g_s , which is deduced from a change in the $\delta^{18}\text{O}$, and 3) the estimated c_i/c_a change is re-interpreted as a change in net photosynthesis (A_{net}) and stomatal conductance (g_s) based on the information derived in the previous steps. The first step is relatively straightforward based on the theory of carbon isotope fractionation during photosynthesis (Breakout Box 1) (Farquhar *et al.*, 1982; Farquhar *et al.*, 1989.). Thus, a significant increase in $\delta^{13}\text{C}$ (scenarios #1, 2, 8) will be interpreted as a decrease in c_i/c_a , a decrease in $\delta^{13}\text{C}$ (scenarios #4 to 6) as an increase in c_i/c_a , and no change in $\delta^{13}\text{C}$ (scenarios #3 and #7) as no change in c_i/c_a . In the second step, a change in $\delta^{18}\text{O}$ can be interpreted as a response of g_s to humidity (Breakout Box 2). This is also

relatively straightforward, given established theory on oxygen isotope fractionation. In particular, enrichment is strongly driven by drier conditions (low RH or high VPD affecting g_s , resulting in a low leaf external to internal partial vapour pressure ratio (e_a/e_i , see Breakout Box 2) and leading to higher $H_2^{18}O$ enrichment, as described by the Pécelt modified Craig-Gordon/Dongman model (Craig & Gordon, 1965; Dongmann *et al.*, 1974; Farquhar & Lloyd, 1993; Cernusak *et al.*, 2002; Kahmen *et al.*, 2008). The third step is the most critical one, resulting in the qualitative nature of the model. As an example, we discuss scenarios #4 and #6, which both indicate increasing c_i/c_a that can be caused by either increasing g_s or decreasing A_{net} . In scenario #4, we additionally know that there is higher $\delta^{18}O$ (“drier conditions”), therefore, the case of increasing g_s seems physiologically implausible and the other possibility, decreasing A_{net} , is selected. In contrast, for scenario #6, we observe lower $\delta^{18}O$ at increasing c_i/c_a and thus increasing g_s is more likely than decreasing A_{net} . This kind of reasoning directly results in the proposed A/g_s scenarios shown in the bottom row of Figure 1.

The use of the original DI-model is applicable for a large number of studies. However, care must be taken when plants respond differently to specific conditions, e.g., to increasing CO_2 , air pollutants, changes in nutrient supply, or extreme drought. This can lead to model outputs that suggest plant responses that contradict our current physiological understanding because such additional and potentially uncontrollable impacts are not considered by current C- or O-isotope fractionation models. Consequently, the DI-model will produce physiologically non-plausible results as it is strictly based on the C and O isotope fractionation and gas exchange principles (see sections 3, 5 and 6). Such discrepancies then need to be resolved, by considering the changes in fractionation mechanisms or different timing of leaf- and stem level processes.

An example for such discrepancies is cases of extreme and on-going drought. Under such conditions the isotopic signal of the leaves could not be found in the stems or other heterotrophic tissue. Since water became so limiting, no growth was possible and the assimilates were barely sufficient to maintain the metabolism. Therefore, no isotope signal representing these hot and dry conditions were evident in the wood (Sarris *et al.*, 2013; Pflug *et al.*, 2015).

Slightly deviating responses are possible and shown with the dashed red arrows in the model outputs, which reflect a certain range of physiological responses rather than a singular direction (Figure 1), as suggested also by (Grams *et al.*, 2007). Overall, the strength of the approach remains obvious: plant functional response to environmental impacts can generally be deduced using isotopes in the absence of detailed physiological measurements. Or, in the case of non-plausible results, the model facilitates the discovery of previously overlooked plant responses through a new lens.

The C and O isotopic signals during carbohydrate transfer from leaf to whole plant (Post photosynthetic fractionation)

The DI-model was originally derived from leaf isotope and gas exchange data. This raises the question as to whether or not, and to what degree, its application to other tissues (e.g. tree rings, roots, etc.) is possible? As the isotope signals of the newly formed assimilates are dampened by post-photosynthetic processes, a careful evaluation is needed to understand whether the patterns are consistent among tissues. Given the available literature (Cernusak *et al.*, 2013, Gessler *et al.*, 2014, and literature therein; Cernusak *et al.*, 2016) we focus on those factors that have the most prominent post-photosynthetic impact on the isotope signals during the carbohydrate transfer processes between leaves and whole plant.

a) Carbon isotopes: Post-photosynthetic fractionations begin immediately following the sugar formation and prior to phloem export. For carbon, kinetic and equilibrium isotope fractionations related to aldolase-catalysed reactions can cause that transitory starch and remobilized sugars at night-time are more ^{13}C -enriched than daytime sucrose (Gleixner *et al.*, 1998, Tcherkez *et al.*, 2004, Tcherkez *et al.*, 2011). However, starch related ^{13}C -fractionations can be very small (Maunoury-Danger *et al.*, 2009) and their significance can vary among species and are under debate (Bögelein *et al.*, 2019, Lehmann *et al.*, 2019a). Moreover, $\delta^{13}C$ differences among individual sugars in leaf and phloem have been observed (Rinne *et al.*, 2015; Churakova *et al.*, 2018), with sucrose being often the most ^{13}C -enriched soluble sugar. The ^{13}C -

fractionations have been related to the invertase reaction (Mauve *et al.* , 2009). In addition, tree tissues, leaves or needles often have high abundances of sugar alcohols or cyclitols, which can be isotopically different from primary sugars (Rinne *et al.* , 2015; Lehmann *et al.* , 2017). The low turnover rate of sugar alcohols can strongly dampen the $\delta^{13}\text{C}$ responses of bulk sugar fractions to changes in physiology and climate (Galiano Pérez *et al.* , 2017). Additional isotope fractionations may occur during the export of leaf sugars into the phloem and during their translocation to sink tissues (Gessler *et al.* , 2009a). Only recently it has been observed that phloem exudates can be more enriched than leaf assimilates and that the ^{13}C -enrichment increased with canopy height (Bögelein *et al.* , 2019). The isotope fractionation was explained by compartmentalisation of leaf sugars in the mesophyll, causing that the designated export sugar sucrose is more ^{13}C -enriched in the cytosol than stored sucrose in the vacuole. As a result of these fractionation processes, leaf $\delta^{13}\text{C}$ values are on average often 2 - 3 (Badeck *et al.* , 2005; Chevillat *et al.* , 2005). Subsequent carbon isotope fractionation associated with branch respiration could also contribute to a shift in isotope ratios (Ghashghaie *et al.* , 2003). Furthermore, mixing between old, stored and new freshly assimilated carbohydrates occurs during phloem transport (Boegelein *et al.* , 2019), which is partly affected by the formation of fresh carbohydrates via stem photosynthesis (Brüggemann *et al.* , 2011).

During early wood formation in spring, trees (particularly broadleaves) utilize stored assimilates from previous years, then gradually switch to fresh more ^{13}C depleted assimilates (Helle & Schleser, 2004). These stored carbohydrate compounds are generally enriched in ^{13}C (Jaeggi *et al.* , 2002) compared to fresh assimilates, especially for starch and its derivatives (Gleixner & Schmidt, 1997).

Also, when considering longer time periods, the impact of industrialization (starting ca. 1850) on the C-isotope ratio in atmospheric CO_2 becomes visible as $\delta^{13}\text{C}$ continuously decreases (Suess effect) resulting from fossil fuel burning and land use change. For the evaluation of time series (tree rings or comparing conserved plant material) this effect needs to be considered (McCarroll & Loader, 2004).

b) Oxygen Isotopes: For $\delta^{18}\text{O}$ in organic matter, the isotope ratio of source water is key. In early studies, the $\delta^{18}\text{O}$ values of precipitation were used as surrogate for source water. However, plants acquire water from a range of lateral and vertical distances with considerable $\delta^{18}\text{O}$ ($\delta^2\text{H}$) gradients depending on the soil structure (Mueller *et al.* , 2014; Sprenger *et al.* , 2016; Goldsmith *et al.* , 2019), resulting in a strongly dampened amplitude of the of the $\delta^{18}\text{O}$ variability of the source water compared to that of precipitation. Thus, the xylem water represents a mixture of water bodies extracted from the soil, with the highest $\delta^{18}\text{O}$ values generally at the soil surface and the lowest at the deepest rooting depth (Barbeta *et al.* , 2020). Further issues arise with seasonal changes in $\delta^{18}\text{O}$ of source water (Saurer *et al.* , 2016). Brinkmann *et al.* , 2018 and Allen *et al.* , 2019, showed that the seasonal origin of the isotope signature of xylem water in summer originated in large part from winter precipitation. Thus, source water varies as a function of both time and space. This temporal shift in the isotope signal needs critical consideration: The source water $\delta^{18}\text{O}$ signal represents a delayed value depending on its seasonal origin relative to $\delta^{18}\text{O}$ in precipitation and the given climatic conditions (Brinkmann *et al.* , 2018; Allen *et al.* , 2019). On the other hand, the effect of the seasonal origin could be balanced under continuous humid climate conditions, as the impact of $\delta^{18}\text{O}$ of water vapour diffusing from the ambient air into the leaf intercellular spaces could override the isotopic signal of the source water (Roden *et al.* , 2004; Lehmann *et al.* , 2019b,).

While no significant isotope fractionation is assumed during H_2O uptake and transport in the xylem along a tree trunk (Barbeta *et al.* , 2020), leaf water is generally enriched in ^{18}O in the leaves during transpiration with its magnitude depending on the atmospheric evaporative demand (VPD) and the $\delta^{18}\text{O}$ of ambient water vapour (Kahmen *et al.* , 2011; Lehmann *et al.* , 2018, Breakout Box 2). During photosynthesis, an exchange between the oxygen atoms originating from CO_2 and that of H_2O takes place imprinting the $\delta^{18}\text{O}$ values of the leaf water on the assimilates because the amount of O-atoms of water is much larger than that of assimilated CO_2 (Lehmann *et al.* , 2017; Barbour *et al.* , 2007; Barbour & Farquhar, 2000;). Thus, freshly assimilated carbohydrates always reflect both the isotopic variations of source and leaf water. Biosynthetic fractionation causes sugars to be about 27‰ water used for photosynthesis (Sternberg & DeNiro, 1983; Yakir & Deniro, 1990). During carbohydrate transport via phloem and concurrent cellulose synthesis, a partial O

exchange with unenriched xylem water occurs. This exchange, resulting in a ca. 40% O-exchange, dampens the leaf water $\delta^{18}\text{O}$ signal (Roden & Ehleringer, 1999, Barbour et al., 2007), likely the largest modification of the $\delta^{18}\text{O}$ signal in assimilates between the leaves and tree rings.

To summarize: For the $\delta^{13}\text{C}$ values, the mixing of old stored carbohydrates (starch and sugar synthesis and remobilization) likely has the strongest dampening effect, probably stronger than phloem loading and (heterotrophic) stem and root respiration. Therefore, we recommend for tree ring analyses to distinguish between early and latewood wherever possible to minimize the inclusion of isotopic signals from old carbohydrates from previous years. For the $\delta^{18}\text{O}$ values, the O-exchange during carbohydrate transport in the phloem and during tree-ring cellulose synthesis probably impacts the oxygen isotope signal the strongest, which might also be amplified by the seasonal origin of the source water (Brinkmann *et al.*, 2018; Allen et al., 2019). To minimize the post-photosynthetic fingerprints on C and O isotope ratios, it is recommended to use late wood in tree rings whenever possible. However, in spite of these modifications and dampening of the C and O isotope values during the carbohydrate transport from leaves to whole plant and cellulose synthesis, the leaf level isotope signal is largely maintained in the wood (Saurer *et al.*, 1997, Song *et al.*, 2011).

Overview of research engaging the model

We identified 261 studies citing Scheidegger et al., 2000 (ISI Web of Knowledge, September, 2018). Of these studies, we excluded review and commentary articles and identified 184 containing new plant carbon and oxygen isotope data (Appendix S1). Studies after this date were considered in the text. These studies covered a broad range of plant life forms, although the majority (89 %) comprised woody (tree or shrub) species. Most studies were performed in temperate regions, but a few were from tropical (dry or moist), polar, and/or dry regions. Gymnosperms and angiosperms from evergreen and deciduous species were similarly represented. Notably, although the DI-model was originally applied to leaf isotope ratios, studies applying the model to wood are more numerous. Only a few studies combined leaf gas exchange and wood measurements.

Paired measurements of leaf gas exchange (i.e. A_{net} and g_s) and isotopes ($\delta^{13}\text{C}$, $\delta/\Delta^{18}\text{O}$) serve as the basis for validating the DI-model to interpret plant functional response to environment; we identified 17 studies with such paired measurements (Table S1). The majority of these studies comprised broadleaf and coniferous trees of temperate and Mediterranean climate regions. Additionally, three studies were performed on grasses and one on herbs. Most of the measurements were carried out at the leaf level on field grown, mature plants, while some studies were performed under controlled conditions on saplings. Four studies compared leaf gas exchange with the isotope pattern of tree stem tissue.

Within the 17 studies with isotopes and leaf gas exchange, we identified 29 different conditions for which the model scenarios had been evaluated (Table 1S, Figure 2). In 18 of the 29 cases (62%), the gas exchange measurements were fully consistent with the observed isotope patterns, clearly validating the DI-model. In a few cases (21%), the scenarios of gas exchange and isotope patterns did clearly not match (e.g. under the influence of air pollutants, extreme droughts and various nitrogen regimes, see also section 5). However, when grouping the responses into situations of improved versus reduced resource availability, rather than individual scenarios, a consistent pattern emerged (Figure 2): Studies that resulted in improved plant resource availability (e.g. light, water, nutrients and CO_2) clustered around scenarios # 6 and 7, both with respect to isotopes and gas-exchange (Figure 2a), consistent with an increase in photosynthesis (A_{net}) and/or stomatal conductance (g_s , Figure 2b). In contrast, a limitation or reduction of resource availability (e.g. increased competition, reduced water/nutrient supply, or temperature stress) commonly reflected scenarios #2 to 4 (Figure 2a). Following the model, plants show a decrease in A_{net} and/or g_s (Figure 2 b). For both groups, these model interpretations are generally well confirmed by gas exchange measurements. As a consequence, this means that an isotope pattern can give reliable information about resource limitation also in the absence of gas-exchange data.

Nevertheless, the DI-model cannot resolve all cases, as it is based on generally accepted plant physiological gas exchange responses to the environment and the well-accepted isotope fractionation models. As indicated

by the grey arrow in Figure 2b, the original DI-model does not predict all A_{net}/g_s cases, particularly not a negative relationship between A_{net} and g_s . The two “missing” A_{net}/g_s scenarios are only observed in rare situations; for instance, a change from cold and wet to moderate and dry conditions, scenario #2 (Figure 1) might cause a decrease in g_s and increase in A_{net} . A similar pattern is observed for increasing ambient CO_2 , where A_{net} is stimulated with a constant or concomitant decline in g_s , corresponding to scenario #8 (Figure 1 and 2). Scenario #5 can be found for plants growing under very humid conditions and (opened stomata) decreasing light (decrease in A_{net} e.g. understory plants in the tropics). The DI-model can also yield non-plausible results for extreme stress conditions such as drought and air pollution. Under such conditions the assumptions of the isotope fractionation models are violated (i.e. See Section 5a and 5e), as the metabolic changes are not considered relative to known fractionation and gas exchange principles.

Model Applications – Environmental Effects on Plant Carbon and Oxygen Isotopes

Here we describe how the DI-model has been applied to infer plant functional responses to common environmental factors over time and space. We explain, which isotope responses would be expected for a certain dominant driving variable, like drought, and review the most common applications of the DI-model as found in the literature, both at the scale of the leaf and that of the whole plant, with some representative examples. A summary of the physiological responses to these environmental drivers, as reflected in the C and O isotope ratios, is given in Table 2.

a) Water Supply and Demand

Drought stress primarily affects plants through diminished soil water availability and increased atmospheric moisture demand, i.e. VPD (Grossiord *et al.*, 2020) that increases non-linearly with rising temperatures (Breshears *et al.*, 2013). Generally, stomata close with increasing drought stress to mitigate leaf turgor loss or stem hydraulic failure and desiccation. This often leads to reduced photosynthetic rates due to limited CO_2 diffusion (Flexas *et al.*, 2008). The strong stomatal response mostly results in increased $\delta^{13}\text{C}$ (Farquhar *et al.*, 1989). Similarly, $\delta^{18}\text{O}$ is expected to increase, particularly when air is getting dry. The increase in VPD (decrease of RH) also causes stomatal closure (reduced g_s) diminishing transpiration. Thus, the Péclet effect is reduced due to a reduced replenishment with H_2^{18}O depleted xylem water minimizing the dilution of the H_2^{18}O enriched leaf water (Farquhar & Lloyd, 1993; Cernusak *et al.*, 2016). Furthermore the back diffusion of ambient ^{18}O -depleted water vapour via stomata into the leaf intercellular spaces is reduced (Lehmann *et al.*, 2018), corresponding to scenario #2 for the isotopes. However, when both A_{net} and g_s are reduced to similar degrees, this may result in a constant c_i/c_a (Ehleringer & Cerling, 1995, Saurer *et al.*, 2004b) and no change in $\delta^{13}\text{C}$, as reflected in scenario #3 (Tab. 2).

Leaf-level isotope and gas exchange studies focused on functional response to water supply and demand have frequently observed scenario #2, including for black poplar plants under various VPD regimes (Rasheed *et al.*, 2015); for conifer species along a wet-dry gradient in Australia (Brodribb *et al.*, 2013), and for leaves and phloem sap in beech trees along a climatic gradient in Europe (Keitel *et al.*, 2006). Therefore, the DI-model generally works for explaining drought-related physiological responses.

However, an isotopic decoupling between leaf-level and tree rings can sometimes result in implausible isotope patterns. During hot and dry summers with low precipitation, water availability is often insufficient for plant growth. Although some CO_2 assimilation is still measurable, no tree growth occurs during these dry, but climatically relevant periods. Thus, there is a mismatch between the period of interest and the period in which the tree ring tissue of interest was formed. Therefore, the DI-model often demonstrates an increase in A_{net} and g_s (scenarios # 7 & 8, Tab 2), which is not a plausible response for hot and dry conditions (high VPD, low soil moisture; (Sarris *et al.*, 2013)). What is observed in the tree rings is actually the $\delta^{18}\text{O}$ signal from cooler, more humid springtime periods, with lower $\delta^{18}\text{O}$ values of precipitation (thus mimicking high g_s see Figure 3). Since most growth occurs during spring time, the remaining tree ring signal is not representative for a hot summer period with little or no growth (Sarris *et al.*, 2013). This has also been observed in a controlled drought experiment with *Quercus robur* and *Quercus petraea* seedlings (Pflug *et al.*, 2015).

In contrast, conditions with high air humidity (RH [?] 90%) and high stomatal conductance allow for high bidirectional water vapour fluxes from leaf-intercellular cavities to the ambient atmosphere and vice versa (Goldsmith *et al.*, 2017, Lehmann *et al.*, 2018). As a result, the isotopic leaf water enrichment is very small and often not detectable. Thus we can assume that, the $\delta^{18}\text{O}$ of leaf water is close to that of the source water (Barbour *et al.*, 2004, Roden *et al.*, 2005). This is mostly the case under long lasting rain and fog conditions, often found in tropical rain forests or given orographic precipitation and persistent fog at the mountainous tree line. Under such conditions, the variability in $\delta^{18}\text{O}$ of leaf water and consequently in cellulose tends toward zero and limits any conclusions regarding g_s (Roden & Siegwolf, 2012).

To summarize: While there may be considerable utility at the leaf scale, the application of the DI-model to tree rings for scenarios involving considerable drought stress or conditions with high humidity must be addressed carefully. Ultimately, changes in water availability and demand are generally well reflected in isotope patterns (scenario 2 & 3, Table 2), as long as plants operate under non-extreme conditions, i.e. severe drought, near saturated air humidity, or at temperature extremes (see section 5b below).

b) Temperature

Temperature plays a critical role in plant function by mediating photosynthesis and post-photosynthetic processes, water use, and ultimately growth. Increases in global temperature have been associated with decreased growth and increased tree mortality in a number of vegetation studies (Allen *et al.*, 2010). Assuming that water supply and VPD are not a limiting factor and thus the stomatal response and $\delta^{18}\text{O}$ values remain unchanged, then temperature primarily affects plant photosynthetic activity at the biochemical level. Consequently, when approaching their optimum temperature range, plants get closer to maximum photosynthesis, leading to a decrease in c_i/c_a and less negative $\delta^{13}\text{C}$ values, which is consistent with scenario #1. In contrast, low and high (non-optimal) temperatures cause a decrease in A_{net} and thus an increase in c_i/c_a and lower $\delta^{13}\text{C}$ values, reflecting scenario #5 (Tab. 2).

Outside of highly controlled experimental conditions, however, a change in the daily air temperature is probably always accompanied by a change in air humidity and thus VPD. A change in g_s due to a possible change in VPD can therefore not fully be excluded and humidity often masks potential temperature effects on gas exchange and thus on isotope fractionation, particularly under warm and dry conditions (Martin-Benito *et al.*, 2010, Moreno-Gutierrez *et al.*, 2012, Liu *et al.*, 2014). In the field, as temperature increases VPD also increases, which affects stomatal conductance much more than temperature (Grossiord *et al.*, 2020). Thus, what is often described as temperature effects is in fact a VPD effect. This may explain the highly equivocal results with both gas exchange and stable isotopes in leaf-level and tree-ring studies; most studies on the effect of temperature did not show any significant response. What further complicates the evaluation of the temperature response of the isotope fractionation is the ability of plants for photosynthetic acclimatization, i.e. that plants shift their photosynthetic temperature optimum with increasing/decreasing temperature.

Studies at tree lines may be useful for assessing temperature effects on plant isotope ratios in the field (Streit *et al.*, 2013), as water availability is not limiting (yearly precipitation > 1000 mm) and atmospheric water demand (VPD) is low. A strong temperature and CO_2 driven relationship was observed between isotope ratios and growth in conifers at the tree line of the Austrian Alps (Wieser *et al.*, 2016), reflecting scenario #6. An often-overlooked condition is low temperatures (0°C to 7°C). Although water is hardly limiting under such conditions, the fact that growth at low temperatures is very slow (Körner, 2015) results in small tree rings. Consequently, the proportion of tree-ring biomass that would contain an isotopic low temperature signal is very small compared to that of the remaining tree ring mass, which was formed under warmer conditions. Therefore, the low temperature signal is hardly visible. So far, only a few studies on low temperature effects to date have matched the predicted scenarios #1 and #5 (Tab.2).

To summarize: Since changes in temperature are mostly accompanied by changes in VPD, but also light intensities, seasonality, and nutrient availability, a single factorial assessment with regard to temperature is only possible in controlled experiments. Evaluating the impact of temperature must always include the effect

of other factors. Nevertheless, there seems to be a temperature driven impact on isotopic fractionation above 15°C and below 40°C on average (see v. Caemmerer, and Evans, 2015).

c) Light

Light drives the variations of c_i/c_a by mediating A_{net} and for variations in e_a/e_i by mediating g_s directly, and indirectly via leaf energy balance, changing the leaf temperature particularly with the given variability of responses under non-light saturated conditions (e.g. shaded plants). The theoretical $\delta^{13}\text{C}$ response is rather straightforward given the strong A_{net} effects on c_i/c_a and thus on $\delta^{13}\text{C}$ values (Ehleringer & Cerling, 1995). Therefore, the $\delta^{13}\text{C}$ would follow the light response curve for A_{net} . However, experimental evidence is somewhat controversial because light effects on carbon isotope discrimination have often been inferred from canopy height gradient studies where other factors may interfere, like VPD gradients and hydraulic constraints for tall trees (Farquhar *et al.*, 1989, Waring & Silvester, 1994). Moreover, the apparent ^{13}C fractionation increases considerably in the low light range (near light compensation point, Barbour *et al.*, 2017a, Busch *et al.*, 2020; Liu *et al.*, 2021; see also Breakout Box 1). Yet the large part of carbon acquisition occurs mostly under high light and biomass production under non-limiting growth conditions (Körner, 2015; Zweifel *et al.*; 2021). Therefore, except for understorey plants the low light impact on ^{13}C fractionation is hardly visible in tree rings. In comparison to $\delta^{13}\text{C}$, changes in $\delta^{18}\text{O}$ might be rather low or follow the inverse g_s direction of the A_{net} response given constant water supply and VPD conditions.

Studies with both gas-exchange and dual-isotope data that considered the influence of light are rare. Boegelein *et al.* (2012) and Roden & Farquhar (2012) show that “pure” light effects may be faithfully described by the DI-model Scenario 7 or 8 (Tab. 2), but that there are often overlapping abiotic effects. Studies focused on understanding the effect of thinning (Giuggiola *et al.*, 2016) using tree rings showed that changes in soil water $\delta^{18}\text{O}$ due to evaporation or changes in transpiration may complicate the application of the model, as soil water availability may become the dominant driver and lead to Scenario 1 or most likely 2 (Tab. 2).

To summarize: What might appear straightforward at first sight turns out to be more complex, as water, temperature, or nutrient regimes can confound interpretations of light as a sole driver of plant physiological response. The strongest impact of light on isotopic fractionation is found under non-saturated light conditions where A_{net} and g_s show its greatest variability (i.e. in the understory); with decreasing light intensity the fractionation of photo- and day respiration has an increasing impact on the net fractionation (Tcherkez *et al.*, 2017; Busch *et al.*, 2020), which is in support of generally lower $\delta^{13}\text{C}$ values of understorey or shaded leaves in contrast to sun lit leaves (see Breakout Box 1).

d) CO_2

Understanding the influence of increasing CO_2 concentration on plant gas exchange and carbon allocation is essential for a correct interpretation of the isotope ratios and to improving carbon cycle models (Schimel *et al.*, 2001). Tree-ring isotope studies, particularly when applied in large spatial networks, can provide unique information on how forests have responded to increases in CO_2 since the start of the industrialization (Frank *et al.*, 2015, Voelker *et al.*, 2016). Based on $\delta^{13}\text{C}$ trends over the last 100 years, presumably caused by increasing CO_2 , three types of gas-exchange responses caused by a variable g_s and A_{net} interaction have been inferred (Saurer *et al.*, 2004b): a) $c_i = \text{constant}$, i.e. the supply function (g_s) decreases and demand functions (A_{net}) stays constant; b) $c_i/c_a = \text{constant}$, g_s and A_{net} change proportionally; c) $c_a - c_i = \text{constant}$, g_s stays constant and A_{net} increases. If we assume that $\delta^{18}\text{O}$ remains constant (no stomatal conductance response) or increases (stomatal conductance decreases), we can deduce the following from the DI-model: for case a) we find either scenario #1 or #2, and for cases b) and c) we find scenarios #4 or #5 (Table 2). However, these are theoretical assumptions and applications of the DI-model to CO_2 responses are relatively rare, particularly in combination with leaf gas-exchange measurements. A 9-year study at an alpine site with a free air CO_2 enrichment (FACE) experiment observed an increase in A_{net} in two conifer species, but no stomatal response to CO_2 (no changes in $\delta^{18}\text{O}$), which is consistent with scenario #1 (Streit *et al.*, 2014). At three FACE sites, (Battipaglia *et al.*, 2013) found changes in intrinsic water-use efficiency between tree

species growing at ambient atmospheric (control) and elevated CO₂ treatments. An increase in $\delta^{13}\text{C}$ of tree-ring cellulose for all species at all CO₂ elevated study sites was observed, but elevated CO₂ resulted in only a slight increase in $\delta^{18}\text{O}$ for some species during some years of the experiment (scenario #1 or #2, Tab. 2).

Natural springs emitting geological CO₂ can be used as long-term CO₂ fertilization experiment with a source isotope signal similar to atmospheric air, thus facilitating the application of the DI-model. Oak trees in a Mediterranean ecosystem in Italy exposed to ca. 800 ppm CO₂ during their lifetime demonstrated increased $\Delta^{13}\text{C}$ values (decreased $\delta^{13}\text{C}$) and unchanged $\delta^{18}\text{O}$ values (scenario #1), indicating a down-regulation of A_{net} and constant g_s for these rather harsh conditions in a dry and nutrient-limited environment (Saurer *et al.*, 2003).

To summarize: The wide variety of isotopic patterns in response to CO₂ changes reflects the variability of species-specific responses to elevated CO₂. This includes a) stimulation of A_{net} under high CO₂ levels (Bader *et al.*, 2013; Streit *et al.*, 2014), b) down regulation of photosynthesis (Grams *et al.*, 2007, Sharma & Williams, 2009), c) reduction in stomatal conductance, or d) no stomatal response to changes in CO₂ (Keele *et al.*, 2007; Bader *et al.*, 2013; Streit *et al.*, 2014; Klein *et al.*, 2016) as summarized in Table 2. Therefore, each isotope data set should be analysed for its response to increasing CO₂ before drawing any further conclusion

e) Atmospheric Pollution

Ozone (O₃): Atmospheric pollution has been shown to affect plants on local and global scales (Omsa *et al.*, 2005; Mils *et al.*, 2018). At the regional scale, O₃ is the most relevant phytotoxic air pollutant (Ashmore, 2005; Ainsworth *et al.*, 2012). Species respond differently to elevated O₃ depending on uptake, mesophyll exposure, detoxification capacity, and plant age (Fuhrer & Booker, 2003, Matyssek & Sandermann, 2003, Paoletti *et al.*, 2020). A unique DI-model study observed that young, but not adult, beech trees were affected by elevated O₃ concentrations (Grams *et al.*, 2007), with an increase in $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ in leaf cellulose (scenario #2). Increasing $\delta^{18}\text{O}$ values in response to O₃ exposure have also been observed in other studies with various species; this was explained by reduced g_s values (Gessler *et al.*, 2009b, Grams & Matyssek, 2010). However, the response of $\delta^{13}\text{C}$ and thus in A_{net} was more variable (scenario #2 and #3). A_{net} responses to O₃ might potentially be biased by the increased enzymatic activity of phospho-enol pyruvate carboxylase (PEPC) (Saurer *et al.*, 1995; Doubnerova & Ryslava, 2011). In contrast to RuBisCo, the net PEPC isotope effect results in a net enrichment of ^{13}C , as PEPC uses bicarbonate as substrate rather than CO₂ (Vogel 1993). Since the equilibrium effect between bicarbonate and CO₂, is larger than the relatively small isotope effect associated with PEP carboxylation, the plant tissue is increased in $\delta^{13}\text{C}$ relative to the RuBisCo fractionation. This observation would intuitively be the result of a reduced c_i/c_a ratio, thus suggesting an increased A_{net} . However, this is in contrast to observed, reduced leaf gas-exchange fluxes with higher c_i/c_a ratios in plants exposed to O₃, which should result in more negative $\delta^{13}\text{C}$ values (Farquhar *et al.*, 1989). The impact of O₃ on C-isotope fractionation demonstrates that once the traditional fractionation model for C3 plants is no longer applicable, the model will yield physiologically implausible results.

Sulphur dioxide (SO₂): Since high ambient SO₂ concentrations have been a rather common problem in the industrialized countries, particularly before the mid-1980s, this pollutant may have had a stronger impact on the stable isotope ratios of plant organic matter than previously recognized. Wagner & Wagner (2006) report that the long-term trend in tree ring $\delta^{13}\text{C}$ showed an extraordinary increase between 1945 and 1990 and a rapid decrease after 1990, mirroring trends in atmospheric SO₂ concentrations. Savard *et al.*, (2004; 2020) showed that sulphur dioxide emissions from smelters induced an increase in $\delta^{13}\text{C}$ by up to 4‰ solely be explained by a reduction of stomatal conductance. SO₂ exposure impacts $\delta^{18}\text{O}$ of leaf water only to a minor degree by reducing stomatal opening (Farquhar & Lloyd, 1993, Sensula & Wilczynski, 2017). Thus, the combined physiological response to high pollution levels is far more pronounced in $\delta^{13}\text{C}$ (Martin *et al.*, 1988, Wagner & Wagner, 2006). Based on the reported isotope patterns, we expect scenarios #1 and #2, i.e. a strong increase in $\delta^{13}\text{C}$ and none-to-moderate change in $\delta^{18}\text{O}$. However, gas exchange responses

suggested by the model do not correspond with published CO_2 and H_2O gas exchange values. Atkinson & Winner, (1987), Kropff *et al.*, (1990), Wedler *et al.*, (1995); Douan *et al.*, (2019) unanimously report a reduction in A_{net} and g_s , which would match Scenarios #3 and #4 (Table 2). The impact of enzymatic detoxification mechanisms, (Randewig *et al.*, 2012), or a possible enhanced PEP-carboxylase activity as found for Ozone, lead to a considerable ^{13}C enrichment, which outweighs the standing model of C-isotope fractionation.

The isotope effects on $\delta^{13}\text{C}$ induced by SO_2 detoxification are not taken into account by the C-isotope fractionation model for C3 plants. Thus, the DI-model output does not correspond with the plant physiological response. It is critical to keep these SO_2 effects in mind when analysing isotopic chronologies from tree ring that originate from regions and periods with large SO_2 emissions (i.e. smelter industries, coal burning, etc.).

Gaseous nitrogen, (NO_2): We cannot exclude effects of gaseous NH_x or NO on plant metabolism and C and O isotope fractionation. Since we found no literature or data describing isotopic effects resulting from these compounds, we discuss only the effects of NO_2 on plant physiology (Sparks, 2009; Wellburn, 1990). Plants demonstrate a proportional increase in $\delta^{13}\text{C}$ with increasing NO_2 concentrations (Siegwolf *et al.*, 2001; Bukata & Kyser, 2005). An increase in $\delta^{13}\text{C}$ with a concomitant decrease of $\delta^{18}\text{O}$, irrespective of the soil nitrogen supply, has also been observed in a growth chamber NO_2 fumigation experiment on *Populus euramericana* (Siegwolf *et al.*, 2001). The interpretation of the C and O isotopic patterns (increase in A_{net} and g_s) corresponded with measured CO_2 and H_2O gas exchange (scenario #8, Tab 2). This was subsequently confirmed by field studies (Saurer *et al.*, 2004a; Guerrieri *et al.*, 2009), but the pattern might be changed when drought occurs. Obviously, protection against drought has the higher priority for plant survival, thus the drought response outweighs the influence of NO_2 (Guerrieri *et al.*, 2010).

To summarize: Air pollutants have remarkable effects on isotopic fractionation, particularly O_3 and SO_2 . Specific changes of enzyme activities mostly result in altered isotope fractionation, which do not agree and are not plausible in the context of well-accepted isotope fractionation and gas exchange principles. While NO_2 can have a fertilizing effect, it can also be toxic for plants at higher concentrations.

f) Soil-N fertilization ($\text{NH}_4^+\text{NO}_3^-$):

The significance of N-depositions and incorporation into the plants is described in Savard *et al.*, (2019) and Etzold *et al.*, 2020. An increase in soil N supply generally reduces $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ (scenario #6), even while exposure to atmospheric NO_2 increases $\delta^{13}\text{C}$ and reduces $\delta^{18}\text{O}$ (Siegwolf *et al.*, 2001). These two different nitrogen sources, NO_2 compared to soil N supply, can have opposite effects on carbon and water relations. One experiment has demonstrated that the addition of N to the soil increases the ratio of A_{net} to g_s in favour of A_{net} , indicating a N-fertilization effect (Guerrieri *et al.*, 2011). This is consistent with the response for resource improvement. However, it has also been observed that a change in the soil N supply to subalpine species resulted in isotopic patterns that were specific to plant functional groups (Bassin *et al.*, 2009). For instance, in the sedge *Carex sempervirens*, both leaf C and O isotope values increased, suggesting a decrease in g_s and a slight increase or constant A_{net} (scenario #2). In contrast, both isotopes declined in forb species, suggesting a constant A_{net} at an increasing g_s (scenario #6). Talhelm *et al.* (2011) and Marshall *et al.*, (2022) also observed a diversity of isotope responses as a result of soil N supply. In their study, *Acer saccharum* foliage and leaf litter material from four different sites was analysed, resulting in four different DI-scenarios (#1, #6, #7 and #8).

To summarize: While plants exposed to atmospheric NO_2 showed a consistent isotope pattern (scenario #8), we find highly diverse isotope responses for soil N supplied plants (Table 2), with a slight trend towards scenario #6. This high diversity of responses may be a result of variable soil conditions (pH, soil moisture and structure, acidic or loamy soil etc.) or it is coupled with other environmental effect. But it also reflects that increased soil N input is not always beneficial, but can shift the competitive balance between species resulting in “winners” and “losers” that respond very differently.

Reconciling Theory and Reality, General Conclusions

The theories of CO₂ and H₂O leaf gas exchange are explicitly linked with those of C and O isotope fractionation, as both isotopic patterns occur concurrently in leaves (Dongmann *et al.* , 1974, Farquhar *et al.* , 1982, Farquhar *et al.* , 1989, Farquhar & Lloyd, 1993, Cernusak *et al.* , 2016). However, when applying the DI-model to real measurements, our review reveals that we sometimes find deviations between measurements and theory.

What is the cause for such deviations between measured data and measured values?

While measured data are mostly analysed given one or maybe two driving variables, a whole spectrum of environmental factors of varying intensities impact gas exchange processes, plant metabolism, and isotopic fractionation in the field. Often these impacts are difficult to identify or to disentangle, e.g. temperature and air humidity (VPD) (see section 5a and 5b). With increasing air temperature, VPD also increases. Changes in isotopic ratios in plants may have been attributed solely to temperature, whereas the major impact was in fact VPD. The effect of varying CO₂ is often difficult to detect depending on site conditions, nutrients, species or climate. This is especially true given that studies of CO₂ in this context are generally carried out retrospectively on tree rings. While stomata respond readily to changes in CO₂ in lab experiments, hardly any stomatal response to CO₂ was observed in field and FACE studies (Körner *et al.* , 2005, Bader *et al.* , 2013, Streit *et al.* , 2014, Klein *et al.* , 2016). As for O₃ exposed plants, the C-fractionation model for C3 plants is no longer valid because PEPC activity is increased, resulting in a considerable increase in $\delta^{13}\text{C}$ of plant organic matter (Saurer *et al.* , 1995). A similar effect is found for SO₂. Thus, studies focusing on a specific plant-environment interaction may easily overlook the concurrent impact of other factors.

Does this invalidate the DI-model?

Whenever the DI-model “fails to fulfil our expectations” the first conclusion is that the DI-model does not work. However, the mechanistic basis of the conceptual model makes it a powerful diagnostic tool. Analysing isotope data in a physiologically mechanistic context facilitates the detection of anomalies or non-plausible responses, which otherwise would not be recognized if $\delta^{13}\text{C}$ or $\delta^{18}\text{O}$ data were studied and interpreted independently from each other. A good example is the tree-ring chronologies from a period and location with heavy SO₂ pollution (Wagner & Wagner, 2006). The anomaly of the carbon isotope values would hardly be detectable by evaluating the $\delta^{13}\text{C}$ time series separately. With the application of the DI-model, the physiologically unrealistic interpretation became apparent and demands further exploration (See section 5e, SO₂). Yet, for a plausible diagnosis of such data constellations, a thorough analysis of plant functional response to specific environmental impacts is essential. Expecting a “plug and play tool,” which can be blindly applied without the essential physiological background can lead us astray and result in wrong conclusions, as also found in the literature. To counteract such misinterpretations, Roden & Siegwolf (2012) published a list of points to consider when the DI-model is applied. Furthermore, the data analysis with the DI-model can be extended by including additional parameters, such as growth data (e.g. tree-ring width; Gessler *et al.* , 2018), or anatomical parameters (Churakova *et al.* , 2019), or adapting the C-isotope fractionation model by considering other enzymatic fractionation processes e.g. including the C4 plant fractionation approach as suggested by Farquhar *et al.* , 1989 and Ubierna *et al.* , 2018.

The use of the DI-model showed that certain isotope patterns can be indicative for an increase or decrease of resource limitation in a wide range of conditions. Based on various studies, we found that scenarios 2-4 (Figure 2) mostly reflect a decrease of resource availability or more stressful conditions that is consistent with gas-exchange results. In this interpretation, there is no need for a perfect match between isotope and gas-exchange scenarios, but rather a range of scenarios is considered. Regarding the application of the DI-model, this re-iterates the qualitative nature of the model, but also its wide-ranging application.

To conclude: Ultimately, whether conceptual (qualitative) or mechanistic (quantitative), models are a synthesis of our current knowledge and provide a powerful platform to test and interpret our measured data. Inconsistencies reveal the lack of consideration of a certain mechanism or reveal knowledge gaps in our understanding of metabolic plant processes. Mismatches between models (representing our current understanding) and reality open doors for further research and enhance our understanding of plant-environment

interactions. Given these considerations, the DI-model has proven to be a successful and powerful tool in plant ecophysiological research.

Authors contributions

R.T.W.S., and G.R.G. planned, and together with M.M.L. and M.S. designed the study. All authors collected the literature, and drafted a first manuscript during a workshop (Stans, CH), which was completed by R.T.W.S., M.S., M.M.L., and G.R.G. All authors commented and accepted the last version of the manuscript.

Acknowledgements

R.T.W.S. acknowledges financial support from the Swiss National Science Foundation under grant numbers CRSII3_136295/1 and 31003A_153428/1. G.R.G. was supported by funding from the European Community's Seventh Framework Program (FP7/2007-2013) under grant agreement 290605 (COFUND: PSI-FELLOW). M.M.L. was supported by the SNF Ambizione grant ("TreeCarbo", No. PZ00P2_179978). O.C-S. was supported by the Russian Science Foundation (RSF) Grant number 21-17-00006 (<https://rscf.ru/en/project/21-17-00006/>) granted to O.C-S. We thank Milena Scandella and Lola Schmid for assistance in assembling the data and in conducting the literature research.

Conflict of interests: None

References

- Andreu-Hayles, L., Lévesque, M., Guerrieri, R., Siegwolf, R.T.W., Körner, C. (2022) Chapt. 14 Limits and Strengths of Tree-Ring Stable Isotopes. In: Siegwolf, R.T.W., Brooks, J.R., Roden, J., Saurer, M. (eds) *Stable Isotopes in Tree Rings. Tree Physiology*, Vol 8. Springer, Cham. https://doi.org/10.1007/978-3-030-92698-4_14
- Allen C.D., Macalady A.K., Chenchouni H., Bachelet D., McDowell N., Vennetier M., Kitzberger T., Rigling A., Breshears D.D. & Hogg E.T. (2010) A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *Forest Ecology and Management* , **259** , 660-684.
- Allen S.T., Kirchner J.W., Braun S., Siegwolf R.T.W. & Goldsmith G.R. (2019) Seasonal origins of soil water used by trees. *Hydrology and Earth System Sciences* , **23** , 1199-1210.
- Ashmore M.R. (2005) Assessing the future global impacts of ozone on vegetation. *Plant Cell and Environment* , **28** , 949-964.
- Ainsworth EA, Yendrek CR, Sitch S, Collins WJ, & Emberson LD (2012) The effects of tropospheric ozone on net primary productivity and implications for climate change. *Annual Review of Plant Biology* , **63** , 637–661. <https://doi.org/10.1146/annurev-arplant-042110-103829>
- Atkinson C.J. & Winner W.E. (1987) GAS-EXCHANGE CHARACTERISTICS OF HETEROMELES-ARBUTIFOLIA DURING FUMIGATION WITH SULFUR-DIOXIDE. *New Phytologist* , **106** , 423-436.
- Badeck F.W., Tcherkez G., Nogues S., Piel C. & Ghashghaie J. (2005) Post-photo synthetic fractionation of stable carbon isotopes between plant organs - a widespread phenomenon. *Rapid Communications in Mass Spectrometry* , **19** , 1381-1391.
- Bader, M.K.F., Leuzinger, S., Keel, S.G., Siegwolf, R.T.W., Hagedorn, F., Schleppi, P. & Körner, C. (2013) Central European hardwood trees in a high-CO₂ future: synthesis of an 8-year forest canopy CO₂ enrichment project. *Journal of Ecology* , **101** , 1509-1519.
- Barbour MM, Farquhar GD. (2000) Relative humidity- and ABA-induced variation in carbon and oxygen isotope ratios of cotton leaves. *Plant, Cell & Environment* **23**: 473–485.
- Barbour M.M. (2007) Stable oxygen isotope composition of plant tissue: a review. *Functional Plant Biology* , **34** , 83-94.

- Barbour M.M., Farquhar G.D. & Buckley T.N. (2017b) Leaf water stable isotopes and water transport outside the xylem. *Plant Cell and Environment* , **40** , 914-920.
- Barbour M.M., Roden J.S., Farquhar G.D. & Ehleringer J.R. (2004) Expressing leaf water and cellulose oxygen isotope ratios as enrichment above source water reveals evidence of a Peclet effect. *Oecologia* , **138** , 426-435.
- Barbour M.M., Ryazanova S., Tcherkez G. (2017a) Respiratory Effects on the Carbon Isotope Discrimination Near the Compensation Point. In: Tcherkez G., Ghashghaie J. (eds) *Plant Respiration: Metabolic Fluxes and Carbon Balance. Advances in Photosynthesis and Respiration (Including Bioenergy and Related Processes)*, vol 43. pp. 144-160. Springer, Cham. https://doi.org/10.1007/978-3-319-68703-2_7
- Barbour MM, Loucos KE, Lockhart EL, et al. (2021) Can hydraulic design explain patterns of leaf water isotopic enrichment in C 3 plants? *Plant Cell Environ.* 2021;44: 432–444. <https://doi.org/10.1111/pce.13943>
- Bassin S., Werner R.A., Soergel K., Volk M., Buchmann N. & Fuhrer J. (2009) Effects of combined ozone and nitrogen deposition on the in situ properties of eleven key plant species of a subalpine pasture. *Oecologia* , **158** , 747-756.
- Battipaglia G., Saurer M., Cherubini P., Calfapietra C., McCarthy H.R., Norby R.J. & Cotrufo M.F. (2013) Elevated CO₂ increases tree-level intrinsic water use efficiency: insights from carbon and oxygen isotope analyses in tree rings across three forest FACE sites. *New Phytologist* , **197** , 544-554.
- Bigeleisen J. (1965) CHEMISTRY OF ISOTOPES. *Science* ,**147** , 463-+.
- Boegelein R., Hassdenteufel M., Thomas F.M. & Werner W. (2012) Comparison of leaf gas exchange and stable isotope signature of water-soluble compounds along canopy gradients of co-occurring Douglas-fir and European beech. *Plant Cell and Environment* ,**35** , 1245-1257.
- Boegelein R., Lehmann M.M. & Thomas F.M. (2019) Differences in carbon isotope leaf-to-phloem fractionation and mixing patterns along a vertical gradient in mature European beech and Douglas fir. *New Phytologist* , **222** , 1803-1815.
- Bowen G.J., Wassenaar L.I., Hobson K.A. (2005) Global application of stable hydrogen and oxygen isotopes to wildlife forensics. *Oecologia***143** , 337–348. DOI 10.1007/s00442-004-1813-y
- Bowen G.J., Liu Z., Vander Zanden H.B., Zhao L., Takahashi G. (2014) Geographic assignment with stable isotopes in IsoMAP. *Methods in Ecology and Evolution* 5: 201–206.
- Breshears D.D., Adams H.D., Eamus D., McDowell N.G., Law D.J., Will R.E., Williams A.P. & Zou C.B. (2013) The critical amplifying role of increasing atmospheric moisture demand on tree mortality and associated regional die-off. *Frontiers in Plant Science* , **4** .
- Brinkmann N., Seeger S., Weiler M., Buchmann N., Eugster W., Kahmen A. (2018) Employing stable isotopes to determine the residence times of soil water and the temporal origin of water taken up by *Fagus sylvatica* and *Picea abies* in a temperate forest. *New Phytologist*,**219**, 1300–1313. doi: 10.1111/nph.15255
- Brodribb T.J., Bowman D.M.J.S., Grierson P.F., Murphy B.P., Nichols S. & Prior L.D. (2013) Conservative water management in the widespread conifer genus *Callitris*. *Aob Plants* , **5** .
- Brüggemann N., Gessler A., Kayler Z., Keel S.G., Badeck F., Barthel M., Boeckx P., Buchmann N., Brugnoli E., Esperschütz J., Gavrichkova O., Ghashghaie J., Gomez-Casanovas N., Keitel C., Knohl A., Kuptz D., Palacio S., Salmon Y., Uchida Y. & Bahn M. (2011) Carbon allocation and carbon isotope fluxes in the plant-soil-atmosphere continuum: a review. *Biogeosciences* , **8** , 3457-3489.
- Bukata A.R. & Kyser T.K. (2005) Response of the nitrogen isotopic composition of tree-rings following tree-clearing and land-use change. *Environmental Science & Technology* , **39** , 7777-7783.

- Busch FA, Holloway-Phillips M, Stuart-Williams H, Farquhar GD. 2020. Revisiting carbon isotope discrimination in C3 plants shows respiration rules when photosynthesis is low. *Nature Plants* 6: 245–258
- Cernusak L.A., Barbour M.M., Arndt S.K., Cheesman A.W., English N.B., Feild T.S., Helliker B.R., Holloway-Phillips M.M., Holtum J.A.M., Kahmen A., McInerney F.A., Munksgaard N.C., Simonin K.A., Song X., Stuart-Williams H., West J.B. & Farquhar G.D. (2016) Stable isotopes in leaf water of terrestrial plants. *Plant, Cell & Environment* ,**39** , 1087-1102.
- Cernusak L.A., Pate J.S. & Farquhar G.D. (2002) Diurnal variation in the stable isotope composition of water and dry matter in fruiting *Lupinus angustifolius* under field conditions. *Plant Cell and Environment* , **25** , 893-907.
- Cernusak L.A., Ubierna N., Winter K., Holtum J.A.M., Marshall J.D. & Farquhar G.D. (2013) Environmental and physiological determinants of carbon isotope discrimination in terrestrial plants. *New Phytologist* , **200** , 950-965.
- Chevillat V.S., Siegwolf R.T.W., Pepin S. & Körner C. (2005) Tissue-specific variation of delta C-13 in mature canopy trees in a temperate forest in central Europe. *Basic and Applied Ecology* ,**6** , 519-534.
- Churakova O.V., Fonti M.V., Saurer M., Guillet S., Corona C., Fonti P., Myglan V.S., Kirdyanov A.V., Naumova O.V., Ovchinnikov D.V., Shashkin A.V., Panyushkina I.P., Buntgen U., Hughes M.K., Vaganov E.A., Siegwolf R.T.W. & Stoffel M. (2019) Siberian tree-ring and stable isotope proxies as indicators of temperature and moisture changes after major stratospheric volcanic eruptions. *Climate of the Past* ,**15** , 685-700.
- Churakova O.V., Lehmann M.M., Saurer M., Fonti M.V., Siegwolf R.T.W. & Bigler C. (2018) Compound-specific carbon isotopes and concentrations of carbohydrates and organic acids as indicators of tree decline in mountain pine. *Forests* , **9** .
- Craig H. & Gordon L.I. (1965) *Deuterium and Oxygen 18 Variations in the Ocean and the Marine Atmosphere* . Consiglio nazionale delle ricerche, Laboratorio de geologia nucleare.
- Dansgaard, W. (1964) Stable isotopes in precipitation. *Tellus* ,**16** , 436-468.
- Dawson T.E., Mambelli S., Plamboeck A.H., Templer P.H. & Tu K.P. (2002) Stable isotopes in plant ecology. *Annual Review of Ecology and Systematics* , **33** , 507-559.
- Dawson T.E. & Siegwolf R.T.W. (2007) *Stable Isotopes as Indicators of Ecological Change* (Terrestrial Ecology Series ed.). Academic Press.
- Dongmann G., Nürnberg H.W., Förstel H. & Wagener K. (1974) On the enrichment of H₂¹⁸O in the leaves of transpiring plants. *Radiation and Environmental Biophysics* ,**11** , 41-52.
- Duan J, Fu B, Kang H, Song Z, Jia M, Cao D, Wei A (2019) Response of gas-exchange characteristics and chlorophyll fluorescence to acute sulfur dioxide exposure in landscape plants, *Ecotoxicology and Environmental Safety*, 171, 122-129. doi.org/10.1016/j.ecoenv.2018.12.064,
- Doubnerová V, Ryslavá H (2011) What can enzymes of C4photosynthesis do for C3plants under stress? *Plant Science* 180 (2011) 575–583.
- Dubbert, M., Cuntz, M., Piayda A., & Werner, C. (2014). Oxygen isotope signatures of transpired water vapor: The role of isotopic non-steadystate transpiration under natural conditions. *New Phytologist*, 203, 1242–1252.
- Dubbert, M., Kübert, A., & Werner, C. (2017). Impact of leaf traits on temporal dynamics of transpired oxygen isotope signatures and its impact on atmospheric vapor. *Frontiers in Plant Science*, 8, 5.
- Ehleringer J.R., Buchmann N. & Flanagan L.B. (2000) Carbon isotope ratios in belowground carbon cycle processes. *Ecological Applications* , **10** , 412-422.

- Ehleringer J.R. & Cerling T.E. (1995) Atmospheric CO₂ and the ratio of intercellular to ambient CO₂ concentrations in plants. *Tree Physiology* , **15** , 105-111.
- Etzold S, Ferretti M, Reinds GJ, Solberg S, Gessler A, Waldner P, Schaub M, Simpson D, Benham S, Hansen K, Ingerslev M, Jonard M, Karlsson PE, Lindroos A-J, Marchetto A, Manninger M, Meessenburg H, Merilä P, Nöjd P, Rautio P, Sanders T.G.M., Seidling W, Skudnik M, Thimonier A, Verstraeten A, Vesterdal L, Vejpustkova M, de Vries W, (2020). Nitrogen deposition is the most important environmental driver of growth of pure, even-aged and managed European forests . *Forest Ecology and Management* 458, 117762. doi.org/10.1016/j.foreco.2019.117762 .
- Evans J.R. & von Caemmerer S. (2013) Temperature response of carbon isotope discrimination and mesophyll conductance in tobacco. *Plant, Cell & Environment* **36**, 745–756. doi: 10.1111/j.1365- 3040.2012.02591.x.
- Evans J.R. (2021) Mesophyll conductance: walls, membranes and spatial complexity. *New Phytologist*. **229** , 1864–1876 doi: 10.1111/nph.16968
- Farquhar G.D., O’Leary M.H. & Berry J.A. (1982) On the relationship between carbon isotope discrimination and the intercellular carbon dioxide concentration in leaves. *Australian Journal of Plant Physiology* , **9** , 121-137.
- Farquhar G.D., Ehleringer J.R. & Hubick K.T. (1989) Carbon isotope discrimination and photosynthesis. *Annual Review of Plant Physiology and Plant Molecular Biology* , **40** , 503-537.
- Farquhar G.D. & Lloyd J. (1993) Carbon and Oxygen Isotope Effects in the Exchange of Carbon Dioxide between Terrestrial Plants and the Atmosphere. In: *Stable Isotopes and Plant Carbon-water Relations* , pp. 47-70. Academic Press, San Diego.
- Farquhar GD, Cernusak LA (2005) On the isotopic composition of leaf water in the non-steady state. *Funct Plant Biol* 32:293–303.
- Farquhar G.D. & Gan K.S. (2003) On the progressive enrichment of the oxygen isotopic composition of water along a leaf. *Plant, Cell and Environment* 26, 1579–1597
- Farquhar, G. D., Griffani, D. S., & Barbour, M. M. (2021). The effects on isotopic composition of leaf water and transpiration of adding a gas-exchange cuvette. *Plant, Cell & Environment*, 44(9), 2844–2857. [https://doi.org/ 10.1111/pce.14076](https://doi.org/10.1111/pce.14076)
- Flexas J., Ribas-Carbo M., Diaz-Espejo A., Galmes J. & Medrano H. (2008) Mesophyll conductance to CO₂: current knowledge and future prospects. *Plant Cell and Environment* , **31** , 602-621.
- Frank D.C., Poulter B., Saurer M., Esper J., Huntingford C., Helle G., Treydte K., Zimmermann N.E., Schleser G.H., Ahlstrom A., Ciais P., Friedlingstein P., Levis S., Lomas M., Sitch S., Viovy N., Andreu-Hayles L., Bednarz Z., Berninger F., Boettger T., D’Alessandro C.M., Daux V., Filot M., Grabner M., Gutierrez E., Haupt M., Hilasvuori E., Jungner H., Kalela-Brundin M., Krapiec M., Leuenberger M., Loader N.J., Marah H., Masson-Delmotte V., Pazdur A., Pawelczyk S., Pierre M., Planells O., Pukiene R., Reynolds-Henne C.E., Rinne K.T., Saracino A., Sonninen E., Stievenard M., Switsur V.R., Szczepanek M., Szychowska-Krapiec E., Todaro L., Waterhouse J.S. & Weigl M. (2015) Water-use efficiency and transpiration across European forests during the Anthropocene. *Nature Climate Change* , **5** , 579-+.
- Fuhrer J. & Booker F. (2003) Ecological issues related to ozone: agricultural issues. *Environment International* , **29** , 141-154.
- Galiano Pérez L., Timofeeva G., Saurer M., Siegwolf R., Martínez-Vilalta J., Hommel R. & Gessler A. (2017) The fate of recently fixed carbon after drought release: towards unravelling C storage regulation in *Tilia platyphyllos* and *Pinus sylvestris* . *Plant, Cell & Environment* , **40** , 1711–1724.
- Gat J., Mook W., Meijer H. (2001) Environmental isotopes in the hydrological cycle, principles and applications, vol. II, Atmospheric water. Vienna, Austria: UNESCO/IAEA.

- Gessler A., Brandes E., Buchmann N., Helle G., Rennenberg H. & Barnard R.L. (2009a) Tracing carbon and oxygen isotope signals from newly assimilated sugars in the leaves to the tree-ring archive. *Plant Cell and Environment* , **32** , 780-795.
- Gessler A., Cailleret M., Joseph J., Schoenbeck L., Schaub M., Lehmann M., Treydte K., Rigling A., Timofeeva G. & Saurer M. (2018) Drought induced tree mortality - a tree-ring isotope based conceptual model to assess mechanisms and predispositions. *New Phytologist* , **219** , 485-490.
- Gessler A., Loew M., Heerdt C., Op de Beeck M., Schumacher J., Grams T.E.E., Bahnweg G., Ceulemans R., Werner H., Matyssek R., Rennenberg H. & Haberer K. (2009b) Within-canopy and ozone fumigation effects on delta C-13 and Delta O-18 in adult beech (*Fagus sylvatica*) trees: relation to meteorological and gas exchange parameters. *Tree Physiology* , **29** , 1349-1365.
- Gessler A., Pedro Ferrio J., Hommel R., Treydte K., Werner R.A. & Monson R.K. (2014) Stable isotopes in tree rings: towards a mechanistic understanding of isotope fractionation and mixing processes from the leaves to the wood. *Tree Physiology* , **34** , 796-818.
- Ghashghaie J., Badeck F.-W., Lanigan G., Nogues S., Tcherkez G., Deleens E., Cornic G. & Griffiths H. (2003) Carbon isotope fractionation during dark respiration and photorespiration in C3 plants. *Phytochemistry Reviews* , **2** , 145-161.yc
- Gimeno T.E., Company C.E., Drake J.E., Barton C.V.M, Tjoelker M.G., Ubierna N., and Marshall J.D., (2020) Whole-tree mesophyll conductance reconciles isotopic and gas- exchange estimates of water-use efficiency Whole-tree mesophyll conductance reconciles isotopic and gas- exchange estimates of water-use efficiency. *New Phytologist* **229**, 2535–2547. doi: 10.1111/nph.17088
- Giuggiola A., Ogee J., Rigling A., Gessler A., Bugmann H. & Treydte K. (2016) Improvement of water and light availability after thinning at a xeric site: which matters more? A dual isotope approach. *New Phytologist* , **210** , 108-121.
- Gleixner G. & Schmidt H.L. (1997) Carbon isotope effects on the fructose-1,6-bisphosphate aldolase reaction, origin for non-statistical C-13 distributions in carbohydrates. *Journal of Biological Chemistry* , **272** , 5382-5387.
- Gleixner G.R., Scrimgeour C., Schmidt H.-L. & Viola R. (1998) Stable isotope distribution in the major metabolites of source and sink organs of *Solanum tuberosum* L.: a powerful tool in the study of metabolic partitioning in intact plants. *Planta* , **207** , 241-245.
- Goldsmith G.R., Allen S.T., Braun S., Siegwolf R.T.W, Kirchner J.W. (2022). Climatic influences on summer use of winter precipitation by trees. *Geophysical Research Letters* 49: e2022GL098323.
- Goldsmith G.R., Allen S.T., Braun S., Engbersen N., Gonzalez-Quijano C.R., Kirchner J.W. & Siegwolf R.T.W. (2019) Spatial variation in throughfall, soil, and plant water isotopes in a temperate forest. *Ecohydrology* , **12** .
- Goldsmith G.R., Lehmann M.M., Cernusak L.A., Arend M. & Siegwolf R.T.W. (2017) Inferring foliar water uptake using stable isotopes of water. *Oecologia* , **184** , 763-766.
- Grams T.E.E., Kozovits A.R., Haeberle K.-H., Matyssek R. & Dawson T.E. (2007) Combining delta C-13 and delta O-18 analyses to unravel competition, CO2 and O-3 effects on the physiological performance of different-aged trees. *Plant Cell and Environment* , **30** , 1023-1034.
- Grams T.E.E. & Matyssek R. (2010) Stable isotope signatures reflect competitiveness between trees under changed CO2/O-3 regimes. *Environmental Pollution* , **158** , 1036-1042.
- Griffiths H., Robinson D. & Gardingen P.v. (1997) *The integration of biological, ecological and geological processes* . Bios Scientific Publishers Ltd., Oxford; UK.

- Grossiord C., Buckley T.N., Cernusak L.A., Novick K.A., Poulter B., Siegwolf R., Sperry J.S. & McDowell N.G. (2020) Plant responses to rising evaporative demand. *New Phytologist* , **226**,1550–1566 doi: 10.1111/nph.16485
- Guerrieri M.R., Siegwolf R.T.W., Saurer M., Jaeggi M., Cherubini P., Ripullone F. & Borghetti M. (2009) Impact of different nitrogen emission sources on tree physiology as assessed by a triple stable isotope approach. *Atmospheric Environment* , **43** , 410-418.
- Guerrieri R., Mencuccini M., Sheppard L.J., Saurer M., Perks M.P., Levy P., Sutton M.A., Borghetti M. & Grace J. (2011) The legacy of enhanced N and S deposition as revealed by the combined analysis of delta 13C, delta 18O and delta 15N in tree rings. *Global Change Biology* ,**17** , 1946-1962.
- Guerrieri R., Siegwolf R., Saurer M., Ripullone F., Mencuccini M. & Borghetti M. (2010) Anthropogenic NOx emissions alter the intrinsic water-use efficiency (WUEi) for Quercus cerris stands under Mediterranean climate conditions. *Environmental Pollution* ,**158** , 2841-2847.
- Helle G. & Schleser G.H. (2004) Beyond CO2-fixation by RuBisCo - an interpretation of C-13/C-12 variations in tree rings from novel intra-seasonal studies on broad-leaf trees. *Plant Cell and Environment* , **27** , 367-380.
- Hirl RT, Ogee J., Ostler U., Schaufele R., Baca Cabrera JC., Zhu J., Schleip J., Wingate L., Schnyder H. (2021) Temperature-sensitive biochemical18O-fractionation andhumidity-dependent attenuation factor are needed to predictδ18O of cellulose from leaf water in a grassland ecosystem. *New Phytologist* 229:3156–3171doi: 10.1111/nph.17111
- Jaeggi M., Saurer M., Fuhrer J. & Siegwolf R. (2002) The relationship between the stable carbon isotope composition of needle bulk material, starch, and tree rings in Picea abies. *Oecologia* , **131** , 325-332.
- Kagawa A, (2022) Foliar water uptake as a source of hydrogen and oxygen in plant biomassTree Physiology 00, 1–21 <https://doi.org/10.1093/treephys/tpac055>
- Kahmen A., Sachse D., Arndt S.K., Tu K.P., Farrington H., Vitousek P.M. & Dawson T.E. (2011) Cellulose δ18O is an index of leaf-to-air vapor pressure difference (VPD) in tropical plants.*Proceedings of the National Academy of Sciences of the United States of America* , **108** , 1981-1986.
- Kahmen A., Simonin K., Tu K.P., Merchant A., Callister A., Siegwolf R., Dawson T.E. & Arndt S.K. (2008) Effects of environmental parameters, leaf physiological properties and leaf water relations on leaf water delta(18)O enrichment in different Eucalyptus species. *Plant Cell and Environment* , **31** , 738-751.
- Keel S.G., Pepin S., Leuzinger S. & Koerner C. (2007) Stomatal conductance in mature deciduous forest trees exposed to elevated CO2.*Trees-Structure and Function* , **21** , 151-159.
- Keitel C., Matzarakis A., Rennenberg H. & Gessler A. (2006) Carbon isotopic composition and oxygen isotopic enrichment in phloem and total leaf organic matter of European beech (Fagus sylvatica L.) along a climate gradient. *Plant Cell and Environment* , **29** , 1492-1507.
- Klein T., Bader M.K.F., Leuzinger S., Mildner M., Schleppi P., Siegwolf R.T.W. & Korner C. (2016) Growth and carbon relations of mature Picea abies trees under 5years of free-air CO2 enrichment. *Journal of Ecology* , **104** , 1720-1733.
- Körner C. (2015) Paradigm shift in plant growth control. *Current Opinion in Plant Biology* , **25** , 107-114.
- Körner C., Asshoff R., Bignucolo O., Hattenschwiler S., Keel S.G., Pelaez-Riedl S., Pepin S., Siegwolf R.T.W. & Zotz G. (2005) Carbon flux and growth in mature deciduous forest trees exposed to elevated CO2.*Science* , **309** , 1360-1362.
- Kropff M.J., Smeets W.L.M., Meijer E.M.J., Vanderzalm A.J.A. & Bakx E.J. (1990) EFFECTS OF SULFUR-DIOXIDE ON LEAF PHOTOSYNTHESIS - THE ROLE OF TEMPERATURE AND HUMIDITY. *Physiologia Plantarum* , **80** , 655-661.

- Lanigan G.J., Betson N., Griffiths H., Seibt U. (2008) Carbon Isotope Fractionation during Photorespiration and Carboxylation in *Senecio*. *Plant Physiology*, **148** , pp. 2013–2020.
- Larcher W. (2003) *Physiological Plant Ecology - Ecophysiology and Stress Physiology of Functional Groups* . (4rd ed.). Springer, Berlin.
- Lehmann M.M., Gamarra B., Kahmen A., Siegwolf R.T.W. & Saurer M. (2017) Oxygen isotope fractionations across individual leaf carbohydrates in grass and tree species. *Plant, Cell & Environment* , **40** , 1658–1670.
- Lehmann M.M., Ghiasi S., George G.M., Cormier M.-A., Gessler A., Saurer M. & Werner R.A. (2019a) Influence of starch deficiency on photosynthetic and post-photosynthetic carbon isotope fractionations. *Journal of Experimental Botany* , **70** , 1829–1841.
- Lehmann M.M., Goldsmith G.R., Mirande-Ney C., Weigt R.B., Schoenbeck L., Kahmen A., Gessler A., Siegwolf R.T.W. & Saurer M. (2020) The O-18-signal transfer from water vapour to leaf water and assimilates varies among plant species and growth forms. *Plant Cell Environ.* 2020;43:510–523 .
- Lehmann M.M., Goldsmith G.R., Schmid L., Gessler A., Saurer M. & Siegwolf R.T.W. (2018) The effect of O-18-labelled water vapour on the oxygen isotope ratio of water and assimilates in plants at high humidity. *New Phytologist* , **217** , 105–116.
- Liu T., Barbour M.M., Yu D., Rao S., Song X. (2021) Mesophyll conductance exerts a significant limitation on photosynthesis during light induction. *New Phytologist* (2021) doi: 10.1111/nph.17757.
- Liu X., An W., Leavitt S.W., Wang W., Xu G., Zeng X. & Qin D. (2014) Recent strengthening of correlations between tree-ring delta C-13 and delta O-18 in mesic western China: Implications to climatic reconstruction and physiological responses. *Global and Planetary Change* , **113** , 23–33.
- Marshall, J.D., Brooks, J.R., Talhelm, A.F. (2022). Forest Management and Tree-Ring Isotopes. In: Siegwolf, R.T.W., Brooks, J.R., Roden, J., Saurer, M. (eds) *Stable Isotopes in Tree Rings*. Tree Physiology, vol 8. Springer, Cham. https://doi.org/10.1007/978-3-030-92698-4_23
- Martin-Benito D., Del Rio M., Heinrich I., Helle G. & Canellas I. (2010) Response of climate-growth relationships and water use efficiency to thinning in a *Pinus nigra* afforestation. *Forest Ecology and Management* , **259** , 967–975.
- Martin B., Bytnerowicz A. & Thorstenson Y.R. (1988) Effects of Air-Pollutants on the Composition of Stable Carbon Isotopes, $\delta^{13}\text{C}$, of Leaves and Wood, and on Leaf Injury. *Plant Physiology* , **88** , 218–223.
- Matyssek R. & Sandermann H. (2003) Impact of ozone on trees: an ecophysiological perspective. *Progress in Botany* **64** , **64** , 349–404.
- Maunoury-Danger F., Bathellier C., Laurette J., Fresneau C., Ghashghaie J., Damesin C. & Tcherkez G. (2009) Is there any $^{12}\text{C}/^{13}\text{C}$ fractionation during starch remobilisation and sucrose export in potato tubers? *Rapid Communications in Mass Spectrometry* , **23** , 2527–2533.
- Mauve C., Bleton J., Bathellier C., Lelarge-Trouverie C., Guerard F., Ghashghaie J., Tchaplal A. & Tcherkez G. (2009) Kinetic $^{12}\text{C}/^{13}\text{C}$ isotope fractionation by invertase: evidence for a small in vitro isotope effect and comparison of two techniques for the isotopic analysis of carbohydrates. *Rapid Communications in Mass Spectrometry* , **23** , 2499–2506.
- McCarroll D. & Loader N.J. (2004) Stable isotopes in tree rings. *Quaternary Science Reviews* , **23** , 771–801.
- Mills G, Pleijel H, Malley CS, Sinha B, Cooper OR, Schultz MG, Neufeld HS, Simpson D, Sharps K, Feng Z, Gerosa G, Harmens H, Kobayashi K, Saxena P, Paoletti E, Sinha V, Xu X (2018) Tropospheric Ozone Assessment Report: Present-day tropospheric ozone distribution and trends relevant to vegetation. *Elem Sci Anth* , **6**(1), p.47. doi.org/10.1525/elementa.302

- Moreno-Gutierrez C., Dawson T.E., Nicolas E. & Ignacio Querejeta J. (2012) Isotopes reveal contrasting water use strategies among coexisting plant species in a Mediterranean ecosystem. *New Phytologist* ,**196** , 489-496.
- Mueller M.H., Alaoui A., Kuells C., Leistert H., Meusburger K., Stumpp C., Weiler M. & Alewell C. (2014) Tracking water pathways in steep hillslopes by delta O-18 depth profiles of soil water. *Journal of Hydrology* , **519** , 340-352.
- Omsa K., Nouchi I. & De Kok L. (2005) *Plant Responses to Air Pollution and Global Change* Springer, Tokyo Berlin Heidelberg New York
- Paoletti E, Grulke NE, Matyssek R (2020) Ozone Amplifies Water Loss from Mature Trees in the Short Term but decreases It in the Long Term.*Forests* **11** (1), 46; <https://doi.org/10.3390/f11010046>
- Pflug E.E., Siegwolf R., Buchmann N., Dobberty M., Kuster T.M., Guenthardt-Goerg M.S. & Arend M. (2015) Growth cessation uncouples isotopic signals in leaves and tree rings of drought-exposed oak trees. *Tree Physiology* , **35** , 1095-1105.
- Poage M.A. and Chamberlain C.P., (2001) Empirical Relationships Between Elevation and the Stable Isotope Composition of Precipitation and Surface Waters: Considerations for Studies of Paleoelevation Change. *American Journal of Science*, Vol. **301** , 1–15.
- Randewig D., Hamisch D., Herschbach C., Eiblmeier M., Gehl C., Jurgeleit J., Skerra J., Mendel R.R., Rennenberg H. & Haensch R. (2012) Sulfite oxidase controls sulfur metabolism under SO₂ exposure in *Arabidopsis thaliana*. *Plant Cell and Environment* , **35** , 100-115.
- Rasheed F., Dreyer E., Richard B., Brignolas F., Brendel O. & Le Thiec D. (2015) Vapour pressure deficit during growth has little impact on genotypic differences of transpiration efficiency at leaf and whole-plant level: an example from *Populus nigra* L. *Plant Cell and Environment* , **38** , 670-684.
- Rinne K.T., Saurer M., Kirdeyanov A.V., Bryukhanova M.V., Prokushkin A.S., Churakova O.V. & Siegwolf R.T.W. (2015) Examining the response of needle carbohydrates from Siberian larch trees to climate using compound-specific $\delta^{13}\text{C}$ and concentration analyses.*Plant, Cell & Environment* , **38** , 2340-2352.
- Roden J., Kahmen A., Buchmann N., Siegwolf R.T.W. (2015) The enigma of effective path length for $\delta^{18}\text{O}$ enrichment in leaf water of conifers. *Plant, Cell & Environment* **38**: 2551–2565.
- Roden J. & Siegwolf R. (2012) Is the dual-isotope conceptual model fully operational? *Tree Physiology* , **32** , 1179-1182.
- Roden J.S., Bowling D.R., McDowell N.G., Bond B.J. & Ehleringer J.R. (2005) Carbon and oxygen isotope ratios of tree ring cellulose along a precipitation transect in Oregon, United States. *Journal of Geophysical Research-Biogeosciences* , **110** .
- Roden J.S. & Ehleringer J.R. (1999) Observations of hydrogen and oxygen isotopes in leaf water confirm the Craig-Gordon model under wide-ranging environmental conditions. *Plant Physiology* , **120** , 1165-1173.
- Roden J.S. & Farquhar G.D. (2012) A controlled test of the dual-isotope approach for the interpretation of stable carbon and oxygen isotope ratio variation in tree rings. *Tree Physiology* , **32** , 490-503.
- Rozanski K, Araguas-Araguas L, Gonfiantini R (1993) Isotopic patterns in modern global precipitation. In: Swart PK, Lohmann KL, McKenzie J, Savin S (eds) Climate change in continental isotopic records. Geophysical Monograph 78 American Geophysical Union, Washington DC, pp 1–37
- Sarris D., Siegwolf R. & Körner C. (2013) Inter- and intra-annual stable carbon and oxygen isotope signals in response to drought in Mediterranean pines. *Agricultural and Forest Meteorology* ,**168** , 59-68.
- Saurer M., Aellen K. & Siegwolf R. (1997) Correlating $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ in cellulose of trees.*Plant Cell and Environment* , **20** , 1543-1550.

- Saurer M., Cherubini P., Ammann M., De Cinti B. & Siegwolf R. (2004a) First detection of nitrogen from NO_x in tree rings: a $^{15}\text{N}/^{14}\text{N}$ study near a motorway. *Atmospheric Environment* , **38** , 2779-2787.
- Saurer M., Cherubini P., Bonani G. & Siegwolf R. (2003) Tracing carbon uptake from a natural CO_2 spring into tree rings: an isotope approach. *Tree Physiology* , **23** , 997-1004.
- Saurer M., Kirdyanov A.V., Prokushkin A.S., Rinne K.T. & Siegwolf R.T.W. (2016) The impact of an inverse climate-isotope relationship in soil water on the oxygen-isotope composition of *Larix gmelinii* in Siberia. *New Phytologist* , **209** , 955-964.
- Saurer M., Maurer S., Matyssek R., Landolt W., Günthardt-Goerg M.S. & Siegenthaler U. (1995) The influence of ozone and nutrition on $\delta^{13}\text{C}$ in *Betula pendula* . *Oecologia* , **103** , 397-406.
- Saurer M., Siegwolf R.T.W. & Schweingruber F.H. (2004b) Carbon isotope discrimination indicates improving water-use efficiency of trees in northern Eurasia over the last 100 years. *Global Change Biology* , **10** , 2109-2120.
- Savard M.M., Begin C., Parent M., Smirnoff A. & Marion J. (2004) Effects of smelter sulfur dioxide emissions: A spatiotemporal perspective using carbon isotopes in tree rings. *Journal of Environmental Quality* , **33** , 13-26.
- Savard MM, Bégin C, Marion J (2020) Response strategies of boreal spruce trees to anthropogenic changes in air quality and rising pCO_2 . *Environm. Poll.*, v. 261. doi.org/10.1016/j.envpol.2020.114209
- Savard MM, Bégin C, Laganière J, Martineau C, Marion J, Stefani, FOP, Séguin A, Smirnoff A, Bergeron J, Morency MJ, Paré D (2019) Anthropogenic N – a global issue examined at regional scale from soils, to fungi, roots and tree rings. *E3S Web Conf.* 98
- Scheidegger Y., Saurer M., Bahn M. & Siegwolf R. (2000) Linking stable oxygen and carbon isotopes with stomatal conductance and photosynthetic capacity: a conceptual model. *Oecologia* , **125** , 350-357.
- Schimel D.S., House J.I., Hibbard K.A., Bousquet P., Ciais P., Peylin P., Braswell B.H., Apps M.J., Baker D., Bondeau A., Canadell J., Churkina G., Cramer W., Denning A.S., Field C.B., Friedlingstein P., Goodale C., Heimann M., Houghton R.A., Melillo J.M., Moore B., Murdiyarso D., Noble I., Pacala S.W., Prentice I.C., Raupach M.R., Rayner P.J., Scholes R.J., Steffen W.L. & Wirth C. (2001) Recent patterns and mechanisms of carbon exchange by terrestrial ecosystems. *Nature* , **414** , 169-172.
- Seibt, U., Wingate, L., Berry, J. A., & Lloyd, J. (2006) Non-steady state effects in diurnal $\delta^{18}\text{O}$ discrimination by *Picea sitchensis* branches in the field. *Plant, Cell and Environment*, **29**, 928–939.
- Sensula B. & Wilczynski S. (2017) Climatic signals in tree-ring width and stable isotopes composition of *Pinus sylvestris* L. growing in the industrialized area nearby Kedzierzyn-Kozle. *Geochronometria* , **44** , 240-255.
- Sharkey T.D. (2012) Editorial: mesophyll conductance: constraint on carbon acquisition by C_3 plants. *Plant, Cell & Environment* **35**, 1881–1883.
- Sharma S. & Williams D.G. (2009) Carbon and oxygen isotope analysis of leaf biomass reveals contrasting photosynthetic responses to elevated CO_2 near geologic vents in Yellowstone National Park. *Biogeosciences* , **6** , 25-31.
- Siegwolf R.T.W., Matyssek R., Saurer M., Maurer S., Günthardt-Goerg M.S., Schmutz P. & Bucher J.B. (2001) Stable isotope analysis reveals differential effects of soil nitrogen and nitrogen dioxide on the water use efficiency in hybrid poplar leaves. *New Phytologist* , **149** , 233-246.
- Song X, Loucos KE, Simonin KA, Farquhar GD, Barbour MM (2015a) Measurements of transpiration isotopologues and leaf water to assess enrichment models in cotton. *New Phytologist*, **206** , 637–646.

- Song, X., Simonin Kevin, A., Loucos Karen, E., & Barbour Margaret, M. (2015b). Modelling non-steady-state isotope enrichment of leaf water in a gas-exchange cuvette environment. *Plant, Cell and Environment*, **38**, 2618–2628.
- Song X., Barbour M.M., Farquhar G.D., Vann D.R. & Helliker B.R. (2013) Transpiration rate relates to within- and across-species variations in effective path length in a leaf water model of oxygen isotope enrichment. *Plant Cell and Environment* , **36** , 1338-1351.
- Song X., Barbour M.M., Saurer M. & Helliker B.R. (2011) Examining the large-scale convergence of photosynthesis-weighted tree leaf temperatures through stable oxygen isotope analysis of multiple data sets. *New Phytologist* , **192** , 912-924.
- Song, X., Lorrey, A., Barbour, M.M. (2022). Environmental, Physiological and Biochemical Processes Determining the Oxygen Isotope Ratio of Tree-Ring Cellulose. In: Siegwolf, R.T.W., Brooks, J.R., Roden, J., Saurer, M. (eds) *Stable Isotopes in Tree Rings. Tree Physiology*, vol 8. Springer, Cham. https://doi.org/10.1007/978-3-030-92698-4_10
- Sparks JP (2009) Ecological ramifications of the direct foliar uptake of nitrogen. *Oecologia* 159:1–13.
- Sprenger M., Leistert H., Gimbel K. & Weiler M. (2016) Illuminating hydrological processes at the soil-vegetation-atmosphere interface with water stable isotopes. *Reviews of Geophysics* , **54** , 674-704.
- Sternberg L. & DeNiro M.J.D. (1983) BIOGEOCHEMICAL IMPLICATIONS OF THE ISOTOPIC EQUILIBRIUM FRACTIONATION FACTOR BETWEEN THE OXYGEN-ATOMS OF ACETONE AND WATER. *Geochimica Et Cosmochimica Acta* , **47** , 2271-2274.
- Streit K., Rinne K.T., Hagedorn F., Dawes M.A., Saurer M., Hoch G., Werner R.A., Buchmann N. & Siegwolf R.T.W. (2013) Tracing fresh assimilates through *Larix decidua* exposed to elevated CO₂ and soil warming at the alpine treeline using compound-specific stable isotope analysis. *The New phytologist* , **197** , 838-849.
- Streit K., Siegwolf R.T.W., Hagedorn F., Schaub M. & Buchmann N. (2014) Lack of photosynthetic or stomatal regulation after 9 years of elevated CO₂ and 4 years of soil warming in two conifer species at the alpine treeline. *Plant Cell and Environment* , **37** , 315-326.
- Talhelm A.F., Pregitzer K.S. & Burton A.J. (2011) No evidence that chronic nitrogen additions increase photosynthesis in mature sugar maple forests. *Ecological Applications* , **21** , 2413-2424.
- Tcherkez G., Farquhar G., Badeck F. & Ghashghaie J. (2004) Theoretical considerations about carbon isotope distribution in glucose of C-3 plants. *Functional Plant Biology* , **31** , 857-877.
- Tcherkez G., Mahe A. & Hodges M. (2011)¹²C/¹³C fractionations in plant primary metabolism. *Trends in Plant Science* , **16** , 499-506.
- Tcherkez G, Gauthier P, Buckley TN, Busch FA, Barbour MM, Bruhn D, Heskell MA, Gong XY, Crous KY, Griffin K et al. (2017). Leaf day respiration: low CO₂ flux but high significance for metabolism and carbon balance. *New Phytologist* **216** , 986–1001.
- Ubierna N., Holloway-Phillips MM., Farquhar G.D. (2018) Using Stable Carbon Isotopes to Study C₃ and C₄Photosynthesis: Models and Calculations. In: Covshoff S. (eds) *Photosynthesis. Methods in Molecular Biology*, vol 1770. Humana Press, New York, NY. https://doi.org/10.1007/978-1-4939-7786-4_10.
- Ubierna N, Farquhar GD (2014) Advances in measurements and models of photosynthetic carbon isotope discrimination in C3 plants. *Plant Cell Environ* **37** , 1494–1498
- Voelker S.L., Brooks J.R., Meinzer F.C., Anderson R., Bader M.K.F., Battipaglia G., Becklin K.M., Beerling D., Bert D., Betancourt J.L., Dawson T.E., Domec J.C., Guyette R.P., Korner C., Leavitt S.W., Linder S., Marshall J.D., Mildner M., Ogee J., Panyushkina I., Plummer H.J., Pregitzer K.S., Saurer M., Smith A.R., Siegwolf R.T.W., Stambaugh M.C., Talhelm A.F., Tardif J.C., Van de Water P.K., Ward J.K. & Wingate L. (2016) A dynamic leaf gas-exchange strategy is conserved in woody plants under changing ambient CO₂:

evidence from carbon isotope discrimination in paleo and CO₂ enrichment studies. *Global Change Biology* , **22** , 889-902.

Vogel, J.C. (1980). Fractionation of the Carbon Isotopes During Photosynthesis. In: Fractionation of the Carbon Isotopes During Photosynthesis. Sitzungsberichte der Heidelberger Akademie der Wissenschaften, vol 1980 / 3. Springer, Berlin, Heidelberg. https://doi.org/10.1007/978-3-642-46428-7_1

Vogel J. C. (1993) Variability of Carbon Isotope Fractionation during Photosynthesis. In: *Stable Isotopes and Plant Carbon-water Relations* , pp. 29-46. Academic Press, San Diego

von Caemmerer S, Evans JR. (2015) Temperature responses of mesophyll conductance differ greatly between species. *Plant, Cell & Environment* **38** , 629–637. doi: 10.1111/pce.12449

Wagner R. & Wagner E. (2006) Influence of air pollution and site conditions on trends of carbon and oxygen isotope ratios in tree ring cellulose. *Isotopes in Environmental and Health Studies* , **42** , 351-365.

Waring R.H. & Silvester W.B. (1994) Variation in foliar $\delta^{13}\text{C}$ values within the crowns of *Pinus radiata* trees. *Tree Physiology* , **14** , 1203-1213.

Wedler M., Weikert R.M. & Lippert M. (1995) PHOTOSYNTHETIC PERFORMANCE, CHLOROPLAST PIGMENTS AND MINERAL-CONTENT OF NORWAY SPRUCE (PICEA ABIES(L) KARST) EXPOSED TO SO₂ AND O₃ IN AN OPEN-AIR FUMIGATION EXPERIMENT. *Plant Cell and Environment* , **18** , 263-276.

Wei T.M., Tcherkez G. , Xu M.W., Schäufele R., Schnyder H., Yusheng Y., Xiao Y.G. (2021) Accounting for mesophyll conductance substantially improves ^{13}C -based estimates of intrinsic water-use efficiency. *New Phytologist* , 229: 1326–1338 doi: 10.1111/nph.16958

Wellburn A.R. (1990) Why are atmospheric oxides of nitrogen usually phytotoxic and not alternative fertilizers. *New Phytologist* , **115** , 395-429.

Wieser G., Oberhuber W., Gruber A., Leo M., Matyssek R. & Grams T.E.E. (2016) Stable Water Use Efficiency under Climate Change of Three Sympatric Conifer Species at the Alpine Treeline. *Frontiers in Plant Science* , **7** .

Yakir D. & Deniro M.J. (1990) Oxygen and hydrogen isotope fractionation during cellulose metabolism in *lemna-gibba* L. *Plant Physiology* , **93** , 325-332.

Yakir D, DeNiro MJ, Gat JR (1990) Natural deuterium and oxygen-18 enrichment in leaf water of cotton plants grown under wet and dry conditions: evidence for water compartmentation and its dynamics. *Plant Cell Environ* 13:49–56.

Zweifel R., Sterck F., Braun S., Buchmann N., Eugster W., Gessler A., Häni M., Peters R.L., Walthert L., Wilhelm M., Zieminska K., Sophia Etzold S. *New Phytologist* (2021) **231** , 2174–2185 doi:10.1111/nph.17552

Zwieniecki, M. A., Brodribb, T. J., & Holbrook, N. M. (2007). Hydraulic design of leaves: Insights from rehydration kinetics. *Plant, Cell & Environment*, 30, 910–921.

Table 1: Summary of research studies containing plant ^{13}C and ^{18}O measurements that cite and apply the DI-model of Scheidegger et al., (2000). Information listed contains the type of material analysed, the availability of gas-exchange data and the region where the study was carried out. Controlled conditions indicate greenhouse or growth chamber studies.

Plant functional type	No. of studies	Wood isotopes only	Leaf isotopes only	Leaves & wood or phloem isotopes	Pl
Gymno-sperms	78	59	15	4	7
Angio-sperms	79	30	43	6	26
Both	27	13	11	3	5

Table 2: Impact of the dominant environmental drivers on tree physiological parameters and stable C and O isotope fractionation: the theoretical responses might deviate from field observations, since other environmental factors impact gas exchange and isotopic fractionation besides the dominant drivers, which does not always correspond with all responses from field observations. g_s ... stomatal conductance; A_{net} ... net photosynthesis; c_i/c_a ... ratio between leaf intercellular versus ambient CO_2 mole fractions; $\Delta^{13}C$... net carbon isotope fractionation; e_a/e_i ... ratio between ambient and substomatal partial water vapour molfractions; $\delta^{18}O$... oxygen isotope fractionation; VPD ... Vapour Pressure Deficit; LAVPD ... Leaf to Air Vapour Pressure Deficit; FACE Free air CO_2 exposure. DI-Model scenario corresponds with the scenario numbers in Figure 1. The red scenario numbers indicate non-plausible gas exchange responses, as derived from the C and O isotope values.

Dominant driving variables	g_s	A_{net}	c_i/c_a	$\delta^{13}C / \Delta^{13}C$	e_a/e_i	$\delta^{18}O / \Delta^{18}O$	DI-Model Scenario
VPD	Decreasing with increasing VPD. The degree of decrease is species dependent	Initially constant, then decreasing with increasing VPD	Progressively decreasing with increasing VPD.	$\delta^{13}C$: increases $\Delta^{13}C$: decreases with increasing VPD	Decreasing with increasing VPD	Both increase with increasing VPD	2, 3 7, 8 (isotopic decoupling Sect. 5a)
Temperature	High at low temperatures, \pm constant over the temperature range between 15°C and 40°C, given that VPD stays constantly low (VPD [?] 0.5 KPa). Decreasing with increasing temperature, mostly due to increasing VPD	Follows an optimum curve increasing from low to higher temperatures decreasing at higher temperatures with an average optimal range between 15°C –30°C. Temperature compensation points on average at 3°C and 40°C	Follows an inverse curve of A_{net} ; with increasing temperature the ratio decreases to remain constant for the optimal range of A_{net} , then increases with decreasing A_{net} .	$\delta^{13}C$: Follows an optimum curve, and is decreasing in the low and high temperature ranges. $\Delta^{13}C$: Follows an inverse curve of A_N , with increases in the low and high temperature ranges.	Decreasing trend with increasing temperature, as a result of a higher leaf than ambient temperature (increasing trend of LAVPD)	$\delta^{18}O$ and $\Delta^{18}O$: Increasing trend due to decreasing e_a/e_i caused by increasing leaf temperature, resulting in increasing LAVPD	1 & 5

Dominant driving variables	g_s	A_{net}	c_i/c_a	$\delta^{13}C / \Delta^{13}C$	e_a/e_i	$\delta^{18}O / \Delta^{18}O$	DI-Model Scenario
PAR	Increasing, following a saturation curve	Increasing, following a saturation curve; light saturation is species and light exposition (sunlit shade) dependent	Declines logarithmically (x-axis mirrors the saturation curve) depending on the A_{net}/g_s ratio	$\delta^{13}C$: increases, following the A_{net} curve $\Delta^{13}C$: decreases, following an inverse curve of A_{net}	Constant or decrease with increasing PAR	Both constant or decrease with increasing PAR	7 or 8
CO ₂	Lab conditions: decreasing with increasing CO ₂ Field studies (FACE) minimal decrease to no response (see Sect. 5d)	Increasing, following a saturation curve, degree of CO ₂ saturation is species- and nutrient-specific.	Lab conditions: decreasing with increasing CO ₂ Field studies: a) decreasing if c_i constant b) c_i/c_a constant c) increasing if $c_a - c_i$ constant	Lab conditions: $\delta^{13}C$ increasing $\Delta^{13}C$ decreasing Field studies a) $\delta^{13}C$ increasing $\Delta^{13}C$ decreasing b) $\delta^{13}C$ decreasing $\Delta^{13}C$ constant c) $\delta^{13}C$ decreasing $\Delta^{13}C$ increasing	Lab conditions: decreasing with increasing CO ₂ Field studies (FACE) constant or slight decrease	$\delta^{18}O$ and $\Delta^{18}O$: Lab conditions: increasing with increasing CO ₂ $\delta^{18}O$ and $\Delta^{18}O$: Field studies (FACE) constant or slight increase	a) 1 or 2 b, c) 4, 5 1, (2) (see Sect. 5d)
Ozone	Initially decreases, but with increasing duration stomata become non-functional	Decreases with progressive duration of exposure (depending on the concentration)	Increases with progressive duration of exposure (depending on the O ₃ concentration)	$\delta^{13}C$: increases while $\Delta^{13}C$ decreases as a consequence of increased PEP carboxylase activity	Increases with progressive duration of exposure (depending on the O ₃ concentration)	Decreases with progressive duration of exposure (depending on the O ₃ concentration)	1 & 2 (PEP-carboxylase driven isotope patterns) 3 (supported by gas exchange measurements)

Dominant driving variables	g_s	A_{net}	c_i/c_a	$\delta^{13}C / \Delta^{13}C$	e_a/e_i	$\delta^{18}O / \Delta^{18}O$	DI-Model Scenario
SO ₂	Initially decreases, but with increasing duration stomata become non-functional	Decreases with progressive duration of exposure (depending on the concentration)	Increases with progressive duration of exposure (depending on the SO ₂ concentration)	$\delta^{13}C$: increases $\Delta^{13}C$: decreases; possibly as a consequence of increased PEP carboxylase activity and other detox. mechanisms	Constant or increasing with progressive duration of exposure (depending on the SO ₂ concentration)	Constant or decreasing with progressive duration of exposure (depending on the SO ₂ concentration)	1 & 2 (biochemically driven isotope patterns) 3 & 4 (based on gas exchange measurements)
NO ₂	Increases for non-toxic concentrations	Increases for non-toxic concentrations	Decreases for non-toxic concentrations	$\delta^{13}C$: increases $\Delta^{13}C$: decreases	decreases	decreases	8
Soil N supply NO ₃ ⁻ and NH ₄ ⁺ Highly variable responses, strongly plant species and Habitat dependent (see Sect. 5f)	a) Increasing b) Decreasing c) Constant d) Increasing e) Constant to decreasing	a) Increasing b) Constant c) Increasing d) Increasing e) Increasing	a) Increasing when A_{net} / g_s ratio is in favour of g_{st} b) Decreasing c) Decreasing d) constant e) decreasing	a) Constant $\delta^{13}C$ and $\Delta^{13}C$ b) $\delta^{13}C$ constant to slightly decreasing and $\Delta^{13}C$ constant c) $\delta^{13}C$ increasing, $\Delta^{13}C$ decreasing d) $\delta^{13}C$ and $\Delta^{13}C$ constant e) $\delta^{13}C$ increasing, $\Delta^{13}C$ decreasing	a) Increasing b) Decreasing c) Constant d) Increasing e) Increasing	$\delta^{18}O$ and $\Delta^{18}O$ decreasing b) $\delta^{18}O$ and $\Delta^{18}O$ increasing c) $\delta^{18}O$ and $\Delta^{18}O$ constant d) $\delta^{18}O$ and $\Delta^{18}O$ decreasing e) $\delta^{18}O$ and $\Delta^{18}O$ decreasing	a) 6 b) 2 c) 1 d) 7 e) 8




Figure 1 : The updated dual C-O isotope model according to Scheidegger et al., (2000). Arrows compare “two varying situations: A and B” as explained in the text. The symbol “ [?] ” indicates no significant change. Arrows pointing up (—) or down (—) stand for an increase/decrease of the respective values. $\Delta^{18}\text{O}$ is the isotopic fractionation for ^{18}O vs. ^{16}O ; $\delta^{13}\text{C}$ is the value for the stable carbon isotope; c_i/c_a is the ratio between the intercellular vs. ambient CO_2 mole fractions; A_{net} is the net photosynthesis (sensu Larcher, 2003); and g_s the stomatal conductance. Red dotted arrows show intermediate A_{net} - g_s -responses not covered in the original model.




Figure 2: Stable C/O isotope and A_{net}/g_s scenarios and their environmental drivers as observed from experimental studies, including a) Isotope patterns reflecting DI-model scenarios #1-8 and b) corresponding observed gas exchange scenarios #1-8. The yellow and grey areas reflect scenarios with generally increasing

and decreasing resource availability, respectively.


Hosted file

image4.emf available at <https://authorea.com/users/449155/articles/595709-updating-the-dual-c-and-o-isotope-gas-exchange-model-a-concept-to-understand-plant-responses-to-the-environment-and-its-implications-for-tree-rings>

Figure 3: Pattern of carbon and oxygen isotopes in a) leaves and b) tree-ring cellulose of *Quercus robur*. Black arrows indicate the directional trend in isotopic patterns due to drought treatments. Open circles represent irrigated plants (control) and open triangles irrigated and warmed plants. Filled circles represent drought treatment and filled triangles stand for drought treatments and warmed plants. Data are means \pm SE, $n = 8$, experiment from 2009 (Pflug *et al.*, 2015). The red arrow and symbols represent the expected values, according to the isotopic pattern in the leaves, whereas the black arrow and symbols are measured data.

Breakout Box 1

Figure 4: The mechanistic linkages between plant physiology and isotopes of carbon at leaf level. c_a and c_i stand for ambient and inter cellular CO_2 mol fractions respectively, g_s and g_m for stomatal and mesophyll conductance; a is the fractionation of CO_2 in the air and through stomata, b the fractionation during carboxylation and A_N is the net photosynthesis.



Carbon Isotopes: Figure 4 above shows a situation with increasing environmental stress from left to right, e.g. increasing drought. The variability of the $^{13}\text{C}/^{12}\text{C}$ isotope ratio in organic matter is predominantly determined by the extent of CO_2 and H_2O gas exchange during photosynthesis (Farquhar *et al.*, 1989, Farquhar *et al.*, 1982, Vogel, 1980). The degree of the stomatal opening regulates the water loss (i.e. transpiration). This leads to a concomitant decrease in the diffusion of CO_2 along the gradient between the ambient air, (c_a) and the leaf intercellular spaces (c_i). Thus, depending on the stomatal conductance (g_s) and the drawdown of $[\text{CO}_2]$ by photosynthesis (A_{net}), c_i will decrease along with the c_i/c_a ratio (see Fig 4). Farquhar *et al.*, (1989; 1982) showed that a reduction in c_i/c_a , decreases the fractionation between ^{13}C and ^{12}C ($[\delta^{13}\text{C}]$). This results in a higher $^{13}\text{C}/^{12}\text{C}$ ratio because the partial pressure of $^{12}\text{CO}_2$ decreases faster, as a consequence of the isotopic effect, (Bigeisen, 1965) than that of the $^{13}\text{CO}_2$, resulting in an increase of the $^{13}\text{CO}_2$ relative to the $^{12}\text{CO}_2$ partial pressure. This higher $^{13}\text{CO}_2$ partial pressure ratio leads to a higher incorporation rate of ^{13}C , expressed in decreased $[\delta^{13}\text{C}]$ (or less negative $\delta^{13}\text{C}$) values of organic matter synthesized under the respective conditions. Consequently, $\delta^{13}\text{C}$ values reflect the interplay between g_s and A_{net} , which is controlled by environmental variations in water supply and demand, as well as photosynthetically active radiation,


temperature or CO₂ concentration. This isotope ratio is transferred to sugars, the primary photosynthetic product, and used for metabolic processes and biomass synthesis (see also Cernusak *et al.* , 2013).

However, the observed (net) fractionation ($\Delta^{13}C_{observed}$ [?] $\Delta^{13}C_{net}$) is a composite of numerous, complex biochemical and physical processes, and can be summarized as $\Delta^{13}C_{observed} = \Delta_b - \Delta_{gs} - \Delta_{gm} - \Delta_e - \Delta_f$, where each fractionation term is flux-rate weighted, (Ubierna *et al.*, 2018). A range of fractionation values are provided by Ubierna & Farquhar (2014). Δ_b is the fractionation associated with the carboxylation of CO₂ via RuBisCo, in the absence of any respiratory fractionation. Δ_{gs} is the fractionation associated with diffusion of CO₂ through the boundary layer and the stomata. Δ_{gm} is the fractionation associated with mesophyll conductance g_m . Δ_e reflects fractionations associated with the day-respiration rate R_d (cytoplasmatic decarboxylation, mitochondrial metabolism, C-remobilization and respiration of the heterotrophic tissues, Tcherkez *et al.*, 2017). Since Δ_b and Δ_{gs} are linked to the largest C-flux and they reflect the largest part of the observed fractionation in the biomass, while Δ_e and Δ_f are associated with the small respiratory C-fluxes and are often neglected. Thus, the $\Delta^{13}C_{observed}$ can be described by the simplified equation 11 The simplified net isotope fractionation for C3-plant is described as $\Delta^{13}C_{observed} = a + (b - a) * c_i / c_a$, with a , the fractionation of CO₂ in the air and through stomata, b the fractionation during carboxylation and c_i and c_a are the mol fractions for the inter cellular and ambient CO₂ concentration. (Ubierna & Farquhar, 2014; Cernusak *et al.*, 2016). However, for plants operating at low light conditions, (e.g. understory plants) the respiratory fractionations might become more relevant (Barbour *et al.*, 2017; Busch *et al.*, 2020; Liu *et al.* ; 2021). A comprehensive overview on day respiration and its impact of e is given in Tcherkez *et al.*, (2017). Moreover, after entering the intercellular spaces the CO₂ molecules traverse the mesophyll cell walls and chloroplast membranes with their diffusional resistance r_{mb} and r_c respectively, which is summarized as $g_m = 1/(r_{mb} + r_c)$. As g_m can contribute significantly to the limitation of photosynthesis by regulating the chloroplast CO₂ concentration (c_c), it impacts the ¹³C-isotope fractionation (Δ_{gm}) accordingly (Evans 2021, Evans & Von Caemmerer, 2013; Sharkey *et al.*; 2012). Yet its experimental determination is challenging and often not possible therefore mostly a model based estimation.

What are the implications of Δ_{gm} , Δ_e and Δ_f with regard to the use of the Dual C and O isotope approach? Mostly the observed fractionation, $\Delta^{13}C_{observed}$ or $\delta^{13}C_{observed}$ is used, which is driven by the most dominating C-flux that is controlled by A_{net} and g_s with its associated fractionation Δ_b and Δ_{gs} . Therefore Δ_{gm} , Δ_e and Δ_f can be neglected. However, as soon as any of these components are separately studied, or derivatives of the fractionation model, e.g. intrinsic water use efficiency (iWUE, Wei *et al.*, 2021, Gimeno *et al.*, 2020) it is essential that the respective fractionation components are taken into account accordingly.

Breakout Box 2

Figure 5: The mechanistic linkages between plant physiology and isotopes of oxygen at the leaf level. e_a and e_i stand for ambient and inter cellular water vapor mol fractions respectively, g_s and g_m for stomatal and mesophyll conductance. ϵ_e and ϵ_k are the equilibrium and kinetic fractionation factors respectively.



Oxygen Isotopes: We must keep in mind that major ^{18}O fractionation processes occur on two different levels: 1st processes, which alter the isotope ratio before water is taken up by plants and 2nd changes, which are a result of ^{18}O fractionation processes during H_2O and CO_2 gas exchange and assimilate incorporation into plant organic matter.

1. The isotope ratio of precipitation water varies depending on i) its **origin** . i.e. water vapour from warm regions like the Mediterranean is more enriched in H_2^{18}O than northern vapour masses (Rozanski et al., 1993). Furthermore continental (Bowen *et al.*, 2005, 2014) or altitudinal (Poage MA & Chamberlain CP, 2001) effects change $\delta^{18}\text{O}$ of precipitation significantly. ii) **Seasonality** of the precipitation is another source of variation in $\delta^{18}\text{O}$ of precipitation water. That, however, is closely linked to the temperature dependence of $\delta^{18}\text{O}$ resulting in an annual sinusoid pattern (Daansgard, 1964, Gat et al., 2001). Further fractionations occur during atmospheric transport, rainfall, canopy throughfall and infiltration in the soil. Depending on the soil structure, season and climate, $\delta^{18}\text{O}$ of soil water shows a great vertical variability (Sprenger *et al.* , 2016), with typically higher $\delta^{18}\text{O}$ values in upper soil layers due to evaporative effects. iii) As plant roots seem to take up water from any **soil depth** , where its accessibility enables its uptake under a minimum of energy, roots potentially have a large range in soil depths from where they absorb water. Brinkmann et al., (2018) Allen et al., (2019) and Goldsmith et al., (2022) found that considerable amounts of absorbed soil water originate from winter water, resulting in temporal shifts (described as temporal origin) of soil $\delta^{18}\text{O}$. Accordingly this leads to a strongly dampened variability of the source water isotope signal (xylem water), the isotopic basis for leaf water H_2^{18}O enrichment in leaves.

2. The variability of the $\delta^{18}\text{O}$ values in leaf organic matter is predominantly determined by the extent of leaf H_2O gas exchange during photosynthesis. Cernusak et al, (2016) and Song et al., (2022) with literature therein provide a summarizing overview of the various aspects of leaf water enrichment. Therefore, we cover only briefly the most important aspects on leaf water enrichment. While leaf water transpires out of the stomata, the lighter H_2^{16}O evaporates more readily than the heavier H_2^{18}O molecules, leaving the remaining leaf water enriched in H_2^{18}O (Craig & Gordon, 1965; Dongmann *et al.* , 1974) (See Figure 5). These authors (CG & D) described leaf water enrichment ($\delta^{18}\text{O}_{\text{LW}}$) at the location of evaporation as a function of the ratio of the ambient versus the leaf the intercellular water vapor mole fraction, e_a/e_i ([?]RH), the $\delta^{18}\text{O}_{\text{source}}$ of source (xylem) water and $\delta^{18}\text{O}_{\text{vap}}$ of ambient water vapor. The extent to which leaf water is enriched in H_2^{18}O is inverse proportional to the ratio e_a/e_i as indicated in the above Figure 5. However, the calculated values according to CG & D tend to overestimate $\delta^{18}\text{O}_{\text{LW}}$ relative to observed values. As a consequence, Farquhar and Lloyd (1993) took the transpiration rate into account by i) introducing the **Péclet effect** . It describes the diffusional flux of enriched water from the location of enrichment into the leaf water body as opposed to a convective water flux of unenriched water molecules from the xylem into the leaf, replenishing the transpired water loss (E). Thus, with increasing E a decrease in $\delta^{18}\text{O}_{\text{LW}}$ is observed (see Figure 5), which results for numerous species in a negative relationship between stomatal conductance (g_s) versus $\delta^{18}\text{O}_{\text{LW}}$ (see

Barbour et al., 2021). A so called pathlength (L) stands for a transport distance of water molecules from the intracellular (through the cell wall) to the leaf inter cellular spaces. This path length (L), however, scarcely represents a real leaf trait and often unrealistic values were used to match predicted versus observed data. The applicability of the Péclet approach is likely tied to the hydraulic leaf design (Zwieniecki et al., 2007) for which three designs “with different pathways for water movements and levels of mesophyll connectedness were defined” (Barbour et al., 2021). Another approach was the application of **ii) a two-pool model** (Song et al., 2015a; Yakiret al., 1990) yielding better estimates for some species, in particular for plants where the leaf was strongly compartmented into a pool with non-enriched and a pool linked to the location of enrichment. Roden et al., (2015) found improved estimates for coniferous plants when the two-pool model was combined with the Péclet correction. Based on a study on 27 species Barbour et al (2021) made recommendations how to consider the hydraulic design for the Péclet correction or the two-pool model or the combination of both. In this context it is also useful to consider the spatial patterns of isotopic composition within leaves, as the isotopic composition tends to be more enriched towards the tip of the leaf or at greater distance from the mid-vein (Cernusak et al., 2016). Farquhar & Gan (2003) considered such effects by expanding the one-dimensional Péclet model, separating convection-diffusion effects in the leaf xylem and lamina.

Another process, which significantly impacts $\delta^{18}\text{O}_{\text{LW}}$ is **iii) the back diffusion** of ambient water vapor into the substomatal cavities (bidirectional fluxes, Seibt et al., 2006, Farquhar et al., 2021), depending on g_s , the ambient water vapor mole fraction (e_a) and its isotopic value ($\delta^{18}\text{O}_{a\tau\mu-a\tau} + \epsilon_\epsilon$). Particularly at high ambient humidity conditions and high g_s (but low E), facilitate a high bidirectional diffusion rate of water molecules (Lehmann et al., 2018; 2020; Kagawa, 2022), impacting $\delta^{18}\text{O}_{\text{LW}}$ considerably and thus $\delta^{18}\text{O}_{\text{organic}}$. Similar to the Péclet effect, the influx of ^{18}O depleted water vapor into the leaf intercellular spaces diminishes the $\delta^{18}\text{O}_{\text{LW}}$ enrichment, with an increasing water vapor influx resulting in a negative relationship between g_s and $\delta^{18}\text{O}_{\text{LW}}$ (Farquhar et al., 2021). This process is often overseen, probably due to the assumption that the isotopic values of source water and ambient water vapor are in equilibrium, which is rarely the case in the field (Boegelein et al., 2019).

iv) Mostly, for the prediction of $\delta^{18}\text{O}_{\text{LW}}$ Isotopic Steady State (ISS) conditions are assumed, yielding often good agreements between observed and predicted values. Changes of ISS / NSS (non-steady state) occur when the transpired water has an isotopic composition differing from that of source water (Farquhar & Cernusak 2005). As the diurnal dynamic of environmental drivers (RH, T_{leaf} , PaR) is considerable the impact on $\delta^{18}\text{O}_{\text{LW}}$ is accordingly, as the prevalent conditions are isotopic NSS. For temporal short-term analyses various studies found remarkable improvements when comparing observed with predicted $\delta^{18}\text{O}_{\text{LW}}$ values applying the NSS approach (Farquhar & Cernusak, 2005; Seibt et al., 2006, Dubbert et al., 2014). In recent years the combined use of gas exchange systems coupled to isotope laser spectrometers allowed for an online monitoring of the isotopic fractionation of $\delta^{18}\text{O}_{\text{LW}}$ under NSS. Song et al., (2015) extended the NSS approach of Farquhar & Cernusak, (2015) considering the mixing of the transpired water with the prevalent cuvette vapor with their respective isotope values. In a next step Dubbert et al., (2017) included leaf traits such as g_s , stomatal density and leaf water content. Farquhar et al., (2021) provides a good discussion on the latest developments on ISS and NSS leaf water enrichment.

v) In contrast to leaf water enrichment, the ISS assumption yields suitable results for organic matter, since the transport of sugars in the phloem and the synthesis of organic material, i.e. cellulose are much slower processes than leaf water enrichment. Note that the organic assimilates ($\delta^{18}\text{O}_{\text{organic}}$) are enriched in ^{18}O relative to the leaf water by 27‰ biochemical fractionation $\epsilon_{\beta_{10}}$ (Sternberg et al., 1983, Lehmann et al., 2017). The assimilates are a mixture of carbohydrates formed under various environmental conditions, representing an integration of the short-term isotope signal variation within a time frame of hours to days (Andreu-Hayles et al., 2022). Furthermore, reserve carbohydrates mix with freshly formed assimilates (Gessler et al., 2014). The mechanistic coherence between leaf water enrichment and sucrose synthesis is summarized in Barbour and Farquhar (2000) and the mathematical description of the oxygen exchange between the carbonyl group and the source water during cellulose synthesis and its application is given in Barbour et al., (2007). Hirl et al., (2021) described the climate sensitivity of these exchange processes and $\epsilon_{\beta_{10}}$ and its significance for the interpretation in climate reconstructions and physiological processes.

Supporting Information

Table S1: Reviewed studies containing both isotope and gas exchange measurements. DI-model scenarios are given for isotope patterns (ISO) and for the gas exchange (GAS). RH = relative humidity, WSOM = Water soluble organic matter, A_{net} = net photosynthetic assimilation rate, A_{sat} = light-saturated photosynthetic assimilation rate. RH = relative humidity. VPD = Vapour pressure deficit. Leaf material: 1= lower, inner canopy, 2=lower, outer canopy, 3=upper, inner and outer canopy.

ISO	GAS	TREATMENT	SPECIES TREATED	SPECIES TYPE	GASEX TYPE	TISSUE	BIOME	CLASSIFICATION	Author
0	0	control to weed removal	<i>Corymbia citri-odora</i>	tree	A _{sat} and g _s	leaf bulk material ¹	Temperate Broadleaf and Mixed Forests	angiosperm	Hu et al 2020
0	0	control to increased temperature (under low RH)	<i>Eucalyptus globulus Labill.</i>	tree	A _{net} and g _s	stem cellulose	Controlled conditions	angiosperm	Ro & qu 2020
0	0	control to defoliation (across growing periods)	<i>Abies balsamea</i>	tree	A _{sat} and g _s	stem cellulose	Controlled conditions	gymnosperm	Sin et al 2020
0	5	control to low light (across both RH)	<i>Eucalyptus globulus Labill.</i>	tree	A _{net} and g _s	stem cellulose	Controlled conditions	angiosperm	Ro & qu 2020
2	2	grass/herb to shrub/tree	<i>various</i>	various	A _{net} and g _s	leaf cellulose	Mediterranean Forests, Woodlands and Shrub	both	Mc Gu et al 2020
2	2	low to high VPD (across genotypes)	<i>Populus nigra</i>	tree	A _{sat} and g _s	leaf bulk material and sugar	Controlled conditions	angiosperm	Ra et al 2020

ISO	GAS	TREATMENT	SPECIES TREATED	SPECIES TYPE	GASEX TYPE	TISSUE	BIOME	CLASSIFICATION	ANALYSIS
2	3	low to high in-traspecific competition intensity	<i>Stipa tenacissima</i>	grass	A_{net} and g_s	leaf bulk material	Mediterranean Forests, Woodlands and Shrub	angiosperm	Ra et al 2006
3	3	open to dense forest	<i>Pinus halepensis</i>	tree	A_{net} and g_s	stem cellulose	Mediterranean Forests, Woodlands and Shrub	gymnosperm	McGuire et al 2006
3	3	control to increased temperature (under high RH)	<i>Eucalyptus globulus Labill.</i>	tree	A_{net} and g_s	stem cellulose	Controlled conditions	angiosperm	Rodriguez & Quirion 2006
3	0	control to drought (across both RH)	<i>Eucalyptus globulus Labill.</i>	tree	A_{net} and g_s	stem cellulose	Controlled conditions	angiosperm	Rodriguez & Quirion 2006
5	5	control to nitrogen deficiency (across both RH)	<i>Eucalyptus globulus Labill.</i>	tree	A_{net} and g_s	stem cellulose	Controlled conditions	angiosperm	Rodriguez & Quirion 2006
6	6	control to weed removal	<i>Corymbia citriodora</i>	tree	A_{sat} and g_s	leaf bulk material ²	Temperate Broadleaf and Mixed Forests	angiosperm	Hu et al 2006
6	6	ambient to high NO ₃	<i>Acer saccharum</i>	tree	A_{net} and g_s	leaf bulk material	Temperate Broadleaf and Mixed Forests	angiosperm	Talbot et al 2006

ISO	GAS	TREATMENT	SPECIES TREATED	SPECIES TYPE	GASEX TYPE	TISSUE	BIOME	CLASSIFICATION	ANALYSIS
6	7	control to irrigation	<i>Triticum durum</i>	grass	A _{net} and g _s	leaf bulk material	Mediterranean Forests, Woodlands and Scrub	angiosperm	Canopy Bo et 20
6	7	ambient to high soil N (across NO ₂ exposures)	<i>Populus x euramerica</i>	tree	A _{net} and g _s	leaf bulk material	Controlled conditions	angiosperm	Sie et 20
7	7	moderate to heavy thinning	<i>Pinus halepensis</i>	tree	A _{sat} and g _s	leaf bulk material	Mediterranean Forests, Woodlands and Scrub	gymnosperm	M Gu et 20
7	7	shift in time (over years)	<i>Larix de-cidua</i>	tree	A _{sat} and g _s	stem cellulose	Temperate Coniferous Forests	gymnosperm	W et 20
7	0	low to high RH (across treatments: drought, nitrogen, light)	<i>Eucalyptus globulus Labill.</i>	tree	A _{net} and g _s	stem cellulose	Controlled conditions	angiosperm	R & qu 20
8	8	base to top canopy gradient	<i>Pseudotsuga men-ziesii</i>	tree	A _{net} and g _s	leaf and phloem WSOM	Temperate Broadleaf and Mixed Forests	gymnosperm	Bo et 20
8	8	control to weed removal	<i>Corymbia citri-odora</i>	tree	A _{sat} and g _s	leaf bulk material ³	Temperate Broadleaf and Mixed Forests	angiosperm	Hu et 20
8	7	base to top canopy gradient	<i>Fagus sylvatica</i>	tree	A _{net} and g _s	leaf and phloem WSOM	Temperate Broadleaf and Mixed Forests	angiosperm	Bo et 20

ISO	GAS	TREATMENT	SPECIES TREATED	SPECIES TYPE	GASEX TYPE	TISSUE	BIOME	CLASSIFICATION	Author
8	7	low to high NO ₂ at low soil N	<i>Populus x eu-ramerica-cana</i>	tree	A _{net} and g _s	leaf bulk material	Controlled conditions	angiosperm	Sie et al 2001
8	0	low to high NO ₂ at high soil N	<i>Populus x eu-ramerica-cana</i>	tree	A _{net} and g _s	leaf bulk material	Controlled conditions	angiosperm	Sie et al 2001
5	5	temperature	<i>H. squamatum</i>	shrub	A _{net} and g _s	leaf bulk material	Arid	angiosperm	Leon-Sanchez et al 2016
3	3	Drought 2014	<i>Triticale</i>	crop	A _{net} and g _s	leaf bulk material	Arid	angiosperm	Mu et al 2013
7	3	Drought 2013	<i>Triticale</i>	crop	A _{net} and g _s	leaf bulk material	Arid	angiosperm	Mu et al 2013
8	3	Drought	<i>Capsicum annum</i>	herb	A _{net} and g _s	leaf bulk material	Controlled conditions	angiosperm	Ser et al 2013
2	2	drought	<i>Lycopersicon esculentum</i> L	herb	A _{net} and g _s	leaf bulk material	Controlled conditions	angiosperm	Wang et al 2013
0	0	nitrogen	<i>Lycopersicon esculentum</i> L	herb	A _{net} and g _s	leaf bulk material	Controlled conditions	angiosperm	Wang et al 2013

References for Table S1

Boegelein R, Hassdenteufel M, Thomas FM, Werner W (2012) Comparison of leaf gas exchange and stable isotope signature of water-soluble compounds along canopy gradients of co-occurring Douglas-fir and European beech. *Plant Cell and Environment* 35:1245-1257. doi: 10.1111/j.1365-3040.2012.02486.x

Cabrera-Bosquet, Llorenc, Albrizio Rossella, Nogues Salvador, Luis Araus Jose (2011). Dual Δ¹³C/δ¹⁸O response to water and nitrogen availability and its relationship with yield in field-grown durum wheat. *Plant, Cell & Environm.*, 34, 418–433 doi: 10.1111/j.1365-3040.2010.02252.x

Huang Zhiquan, Zhihong Xu, Timothy J. Blumfield, Ken Bubb (2008). Foliar δ¹³C and δ¹⁸O reveal differential physiological responses of canopy foliage to pre-planting weed control in a young spotted gum (*Corymbia citriodora* subsp. *Variegata*) plantation Volume 28, Issue 10, Pages 1535–1543, <https://doi.org/10.1093/treephys/28.10.1535>

Leon-Sanchez L, Nicolas E, Nortes PA, Maestre FT, Querejeta JI (2016) Photosynthesis and growth reduction with warming are driven by nonstomatal limitations in a Mediterranean semi-arid shrub. *Ecology and Evolution* 6:2725-2738. doi: 10.1002/ece3.2074

- Moreno-Gutierrez C et al., (2011) Leaf delta ^{18}O of remaining trees is affected by thinning intensity in a semiarid pine forest. *Plant Cell and Environment* 34:1009-1019. doi: 10.1111/j.1365-3040.2011.02300.x
- Moreno-Gutierrez C, Dawson TE, Nicolas E, Ignacio Querejeta J (2012) Isotopes reveal contrasting water use strategies among coexisting plant species in a Mediterranean ecosystem. *New Phytologist* 196:489-496. doi: 10.1111/j.1469-8137.2012.04276.x
- Munjonji L, Ayisi KK, Vandewalle B, Haesaert G, Boeckx P (2016) Combining carbon-13 and oxygen-18 to unravel triticale grain yield and physiological response to water stress. *Field Crops Research* 195:36-49. doi: 10.1016/j.fcr.2016.06.001
- Ramirez, David A, Querejeta, Jose I, Bellot Juan (2009). Bulk leaf delta O-18 and delta C-13 reflect the intensity of intraspecific competition for water in a semi-arid tussock grassland. *Plant, Cell and Environment*, 32, 1346–1356 doi: 10.1111/j.1365-3040.2009.02002.x
- Rasheed F, Dreyer E, Richard B, Brignolas F, Brendel O, Le Thiec D (2015) Vapour pressure deficit during growth has little impact on genotypic differences of transpiration efficiency at leaf and whole-plant level: an example from *Populus nigra* L. *Plant Cell and Environment* 38:670-684. doi: 10.1111/pce.12423
- Roden JS, Farquhar GD (2012) A controlled test of the dual-isotope approach for the interpretation of stable carbon and oxygen isotope ratio variation in tree rings. *Tree Physiology* 32:490-503. doi: 10.1093/treephys/tps019
- Serret MD et al., (2018) Interactive Effects of CO₂ Concentration and Water Regime on Stable Isotope Signatures, Nitrogen Assimilation and Growth in Sweet Pepper. *Frontiers in Plant Science* 8. doi: 10.3389/fpls.2017.02180
- Siegwolf RTW et al., (2001) Stable isotope analysis reveals differential effects of soil nitrogen and nitrogen dioxide on the water use efficiency in hybrid poplar leaves. *New Phytologist* 149:233-246. doi: 10.1046/j.1469-8137.2001.00032.x
- Simard Sonia, Hubert Morin, Cornelia Krause, William M. Buhay, Kerstin Treydte (2012). Tree-ring widths and isotopes of artificially defoliated balsam firs: A simulation of spruce budworm outbreaks in Eastern Canada. *Environmental and Experimental Botany* 81, 44–54.
- Talhelm AF, Pregitzer KS, Burton AJ (2011) No evidence that chronic nitrogen additions increase photosynthesis in mature sugar maple forests. *Ecol. Appl.* 21:2413-2424. doi: 10.1890/10-2076.1
- Wang Chao, Shuxia Wu, Moussa Tankari, Ximei Zhang, Li Li, Daozhi Gong, Weiping Hao, Yanqing Zhang, Xurong Mei, Yufei Wang, Fulai Liu, Yaosheng Wang (2018). Stomatal aperture rather than nitrogen nutrition determined water use efficiency of tomato plants under nitrogen fertigation. *Agricultural and water management* (209), 94-101.
- Wieser G, Oberhuber W, Gruber A, Leo M, Matyssek R, Grams TEE (2016) Stable Water Use Efficiency under Climate Change of Three Sympatric Conifer Species at the Alpine Treeline. *Frontiers in Plant Science* 7. doi: 10.3389/fpls.2016.00799

Hosted file

Revised_Main-Text-Siegwolf-et-al_221113.docx available at <https://authorea.com/users/449155/articles/595709-updating-the-dual-c-and-o-isotope-gas-exchange-model-a-concept-to-understand-plant-responses-to-the-environment-and-its-implications-for-tree-rings>

Hosted file

Figure 1.docx available at <https://authorea.com/users/449155/articles/595709-updating-the-dual-c-and-o-isotope-gas-exchange-model-a-concept-to-understand-plant-responses-to-the-environment-and-its-implications-for-tree-rings>

Hosted file

Figure 2.docx available at <https://authorea.com/users/449155/articles/595709-updating-the-dual-c-and-o-isotope-gas-exchange-model-a-concept-to-understand-plant-responses-to-the-environment-and-its-implications-for-tree-rings>

Hosted file

Figure 3.docx available at <https://authorea.com/users/449155/articles/595709-updating-the-dual-c-and-o-isotope-gas-exchange-model-a-concept-to-understand-plant-responses-to-the-environment-and-its-implications-for-tree-rings>

Hosted file

Figure 4.docx available at <https://authorea.com/users/449155/articles/595709-updating-the-dual-c-and-o-isotope-gas-exchange-model-a-concept-to-understand-plant-responses-to-the-environment-and-its-implications-for-tree-rings>

Hosted file

Figure 5.docx available at <https://authorea.com/users/449155/articles/595709-updating-the-dual-c-and-o-isotope-gas-exchange-model-a-concept-to-understand-plant-responses-to-the-environment-and-its-implications-for-tree-rings>

Hosted file

Table 1.docx available at <https://authorea.com/users/449155/articles/595709-updating-the-dual-c-and-o-isotope-gas-exchange-model-a-concept-to-understand-plant-responses-to-the-environment-and-its-implications-for-tree-rings>

Hosted file

Table 2.docx available at <https://authorea.com/users/449155/articles/595709-updating-the-dual-c-and-o-isotope-gas-exchange-model-a-concept-to-understand-plant-responses-to-the-environment-and-its-implications-for-tree-rings>