

Dynamics of a diffusive predator-prey model with prey-stage structure and prey-taxis

Yan Li¹, Xiuzhen Fan², and Hui Hao²

¹China University of Petroleum Huadong - Qingdao Campus

²China University of Petroleum East China - Qingdao Campus

March 07, 2024

Abstract

This paper is concerned with a diffusive predator-prey model with prey-taxis and prey-structure under the homogeneous Neumann boundary condition. The stability of the unique positive constant equilibrium of the predator-prey model is derived. Hopf bifurcation and steady state bifurcation are also concluded.

Dynamics of a diffusive predator-prey model with prey-stage structure and prey-taxis*

Yan Li[†], Xiuzhen Fan, Hui Hao

College of Science, China University of Petroleum(East China),
Qingdao 266580, PR China

Abstract

This paper is concerned with a diffusive predator-prey model with prey-taxis and prey-structure under the homogeneous Neumann boundary condition. The stability of the unique positive constant equilibrium of the predator-prey model is derived. Hopf bifurcation and steady state bifurcation are also concluded.

Keywords: Predator-prey model; Hopf bifurcation; Steady state bifurcation; Prey-taxis.

1 Introduction

Predator-prey models have always been considered to be classical, and the source of all the research work in population models during the past century is the Lotka-Volterra model [1, 2]. In the predator-prey models, the functional response is one of the crucial factors, which affect population dynamics. Typically, the Lotka-Volterra interaction term can be classified into many different types, for instance, Holling type I-IV [3, 4], Holling-Tanner type [5, 6], Beddington-DeAngelis type [7, 8, 9], ratio-dependent type [10], and Ivlev type [11]. Dynamic structure of the system is not only related to the response function but also may depend on many other factors such as location, age and mature delay and so on. The life histories of plants, insects, and animal life histories exhibit enormous diversity. Metamorphosis may carry the same individual through several totally different niches during a lifetime. Specialized stages may exist for dispersal or dormancy. The vital rates (rates of survival, development, and reproduction) almost always depend on age, size, or development stage. Population growth models that include age, stage or body size structure often predict complex population dynamics. Due to the above realistic evidences, the stage-structured models have received much attention in recent years, see [13, 14, 15, 16, 17, 18, 19, 20, 21, 22]. Generally speaking, population growth models that include stage structure predict more complex population dynamics than those without stage structure.

In the spatial-temporal predator-prey interaction, with the exception of the random diffusion of predators and the prey, the variations of the predator's

*This work was supported by Shandong Provincial Natural Science Foundation (Nos.ZR2019MA006,ZR2021MA025,ZR2021MA028).

[†]Corresponding author

Email address: liyan@upc.edu.cn

velocity are often directed by prey gradient i.e. prey-taxis. Kareiva and Odell [23] first derived a prey-taxis model to describe the predator aggregation in high prey density areas. Since then, more and more scholars have studied the predator-prey model with prey-taxis. For example, Lee et al. showed that the prey-taxis tends to reduce the likelihood of pattern formation and effective bio-control in [24]. Wang et al. [25] investigated the global existence, boundedness and global stabilities of the equilibria for a two predators and one prey model with prey-taxis. We refer to [26, 27, 28, 29, 30] for other interesting works on models with prey-taxis. It has been recognized that the systems with prey-taxis may undergo more rich dynamics and generate different spatial patterns than without prey-taxis.

In [12], the authors established the following predator-prey model with general functional response and stage-structure for the prey:

$$\begin{cases} \frac{du}{dt} = av - bu - \gamma u^2 - g(u)w, & t > 0, \\ \frac{dv}{dt} = u - v, & t > 0, \\ \frac{dw}{dt} = w(-r + \delta g(u)), & t > 0 \end{cases} \quad (1)$$

where u, v are the population densities of immature and mature prey species, respectively. w denotes the density of predator population. $a, b, r, \delta > 0$, and the functional response $g(u)$ also satisfies

$g(0) = 0$, $g'(u) > 0$ ($u \geq 0$), and $0 < g(u) < L$, L is a positive constant.

For the background of (1), the readers can refer to [12]. The authors studied the stability of equilibrium points for this ODE system via linearization and the Lyapunov method, and showed that Hopf bifurcation occurs in paper [12].

It is known that the distributions of populations, in general, being heterogeneous, depend not only on time, but also on the spatial positions in habitat. So it is natural and more precise to study the corresponding PDE problem. In paper [12], the authors also considered the following corresponding reaction-diffusion system:

$$\begin{cases} \frac{\partial u}{\partial t} - d_1 \Delta u = av - bu - \gamma u^2 - g(u)w, & x \in \Omega, t > 0, \\ \frac{\partial v}{\partial t} - d_2 \Delta v = u - v, & x \in \Omega, t > 0, \\ \frac{\partial w}{\partial t} - d_3 \Delta w = w(-r + \delta g(u)), & x \in \Omega, t > 0, \\ \frac{\partial u}{\partial n} = \frac{\partial v}{\partial n} = \frac{\partial w}{\partial n} = 0, & x \in \partial\Omega, t > 0, \\ u(x, 0) = u_0(x), v(x, 0) = v_0(x), w(x, 0) = w_0(x) \geq 0 & x \in \Omega \end{cases} \quad (2)$$

where $\Omega \subseteq \mathbb{R}^N$ ($N \geq 1$) is a bounded domain with smooth boundary $\partial\Omega$, n is the outward unit normal vector of the boundary $\partial\Omega$. d_1, d_2 and d_3 are

1 positive constants which stand for the random diffusion rates of the three species,
2 respectively. The homogeneous Neumann boundary condition indicates that
3 the predator-prey system is self-contained with zero population flux across the
4 boundary. $u_0(x)$, $v_0(x)$ and $w_0(x)$ are nonnegative smooth functions on $\bar{\Omega}$. In
5 [12], the existence and uniform boundedness of global solutions and stability
6 of equilibrium points for the corresponding reaction-diffusion problem (2) were
7 discussed. In addition, by using the topological degree theory, the existence
8 of nontrivial steady states of system (2) under certain situations was showed,
9 and some nonexistence of nontrivial steady state results were also obtained. We
10 refer to [31, 32, 33, 34, 35] for the studies on the reaction-diffusion in other
11 three-species predator-prey models.

12 In this paper, we introduce prey-taxis term into problem (2) and investigate
13 the effect of the prey-taxis on the dynamics of predator-prey model. Thus, we
14 shall consider the following model:

$$\begin{cases} \frac{\partial u}{\partial t} - d_1 \Delta u + \chi \nabla(u \nabla v) = av - bu - \gamma u^2 - g(u)w, & x \in \Omega, t > 0, \\ \frac{\partial v}{\partial t} - d_2 \Delta v = u - v, & x \in \Omega, t > 0, \\ \frac{\partial w}{\partial t} - d_3 \Delta w + \rho \nabla(w \nabla u) = w(-r + \delta g(u)), & x \in \Omega, t > 0, \\ \frac{\partial u}{\partial n} = \frac{\partial v}{\partial n} = \frac{\partial w}{\partial n} = 0, & x \in \partial\Omega, t > 0, \\ u(x, 0) = u_0(x), v(x, 0) = v_0(x), w(x, 0) = w_0(x) \geq 0 & x \in \Omega \end{cases} \quad (3)$$

15 where the terms $\chi \nabla(u \nabla v)$ and $\rho \nabla(w \nabla u)$ are actually taxis mechanisms and
16 $\chi, \rho > 0$ are their taxis rates, respectively. The term $\chi \nabla(u \nabla v)$ models the
17 movement of the immature prey which is directed toward the increasing mature
18 prey densities. The term $\rho \nabla(w \nabla u)$ accounts for prey-taxis which describes the
19 phenomenon that the predator has the tendency to move increasingly toward
20 the immature prey gradient direction. In the following paper, we will study the
21 Hopf bifurcation and steady-state bifurcation of problem (3) by choosing the
22 prey-taxis rate.

23 The outline of this paper is as follows. In Section 2, after analyzing the
24 characteristic equation, we conclude the stability of constant equilibria of prob-
25 lem (3). In Section 3, we research the existence of periodic solutions bifurcating
26 from the unique positive constant equilibrium of problem (3). In Section 4,
27 we consider steady state bifurcations to show the existence of nontrivial steady
28 state solutions of (3).

Throughout the paper, μ_k denotes the eigenvalues of $-\Delta$ in Ω under the
homogeneous Neumann boundary condition satisfying

$$0 = \mu_0 < \mu_1 \leq \mu_2 < \cdots \mu_k < \cdots < \infty.$$

29 **2 Stability of constant equilibria of problem (3)**

30 In this section we will study the stability of constant equilibria of problem (3).

It is easy to see that the trivial equilibrium point $(0, 0, 0)$ always exists. If $a > b$, semi-trivial equilibrium point $(\bar{u}, \bar{v}, 0) = (\frac{a-b}{\gamma}, \frac{a-b}{\gamma}, 0)$ exists. Especially if

$$a - b - \gamma u^* > 0, \quad \text{and} \quad \delta L > r \quad (4)$$

a unique positive constant equilibrium (u^*, v^*, w^*) also exists, where

$$u^* = v^* = g^{-1}\left(\frac{r}{\delta}\right), \quad w^* = \frac{\delta}{r}(a - b - \gamma u^*)u^*.$$

When $\chi = \rho = 0$, in [12], the stability of constant equilibria of (1) and (2) has been studied. We shall perform linearized stability analysis to see the effects of prey-taxis coefficients χ and ρ .

Theorem 2.1. *For problem (3),*

(1) *If $a < b$, then $(0, 0, 0)$ is locally asymptotically stable; if $a > b$, then $(0, 0, 0)$ is unstable;*

(2) *Let $a > b$, $r > \delta L$ hold. If*

$$\chi \leq \frac{\gamma}{a-b}[d_1 + d_2(2a-b)],$$

$(\bar{u}, \bar{v}, 0)$ is locally asymptotically stable;

(3) *Assume that (4) holds. If*

$$\chi \leq \min\{\chi_1, \chi_2\}, \quad \text{and} \quad a \leq \eta \quad (5)$$

where

$$\begin{aligned} \chi_1 &= \frac{d_1 + d_2\eta + d_3(\eta+1)}{u^*}, \\ \chi_2 &= \frac{d_1 + d_2\eta + \frac{d_2}{d_3}\rho w^* g(u^*)}{u^*}, \\ \eta &= b + 2\gamma u^* + g'(u^*)w^*, \end{aligned}$$

then (u^*, v^*, w^*) is locally asymptotically stable.

Proof We will prove the above points in turn.

(1) Linearizing problem (3) at the trivial equilibrium $(0, 0, 0)$, the Jacobi matrix at $(0, 0, 0)$ is as follows

$$J_{(0,0,0)} = \begin{pmatrix} -d_1\mu_k - b & a & 0 \\ 1 & -d_2\mu_k - 1 & 0 \\ 0 & 0 & -d_3\mu_k - r \end{pmatrix}, \quad k = 0, 1, 2, \dots,$$

so the characteristic equation is

$$[\lambda^2 + (d_1\mu_k + b + d_2\mu_k + 1)\lambda + d_1d_2\mu_k^2 + (d_1 + d_2b)\mu_k + b - a](\lambda + d_3\mu_k + r) = 0.$$

Hence, according to Routh-Hurwitz criterion, we obtain that if $a < b$, then $(0, 0, 0)$ is locally asymptotically stable; if $a > b$, then $(0, 0, 0)$ is unstable; if $a = b$, when $k = 0$, we have $\mu_k = 0$, so the characteristic equation is

$$\lambda(\lambda + b + 1)(\lambda + r) = 0,$$

- 1 then the roots of the equation are $\lambda_1 = 0, \lambda_2 = -(b + 1) < 0, \lambda_3 = -r < 0$.
- 2 There is a zero eigenvalue, so we need to use the manifold theorem to judge the
- 3 stability, which will not be discussed here.

(2) Similar to the above, the Jacobi matrix at $(\bar{u}, \bar{v}, 0)$ is

$$J_{(\bar{u}, \bar{v}, 0)} = \begin{pmatrix} -d_1\mu_k - b - 2\gamma\bar{u} & \mu_k\chi\bar{u} + a & -g(\bar{u}) \\ 1 & -d_2\mu_k - 1 & 0 \\ 0 & 0 & -d_3\mu_k - r + \delta g(\bar{u}) \end{pmatrix}, \quad k = 0, 1, 2, \dots,$$

the characteristic equation at $(\bar{u}, \bar{v}, 0)$ is

$$[\lambda^2 + (d_1\mu_k + 2a - b + d_2\mu_k + 1)\lambda + (d_1\mu_k + 2a - b)(d_2\mu_k + 1) - (\chi\frac{a-b}{\gamma}\mu_k + a)](\lambda + d_3\mu_k + r - \delta g(\frac{a-b}{\gamma})) = 0.$$

It is observed that $r - \delta g(\frac{a-b}{\gamma}) > 0$ since $r > \delta L$ holds. Moreover according to Routh-Hurwitz criterion, it is easy to get that if

$$\chi \leq \frac{\gamma}{a-b}[d_1 + d_2(2a - b)],$$

- 4 then $(\bar{u}, \bar{v}, 0)$ is locally asymptotically stable.
- 5 (3) Due to the standard linearized stability principle, the linearized stability
- 6 of (u^*, v^*, w^*) is determined by eigenvalues of the following matrixes

$$N_k = \begin{pmatrix} -d_1\mu_k - \eta & \mu_k\chi u^* + a & -g(u^*) \\ 1 & -d_2\mu_k - 1 & 0 \\ \mu_k\rho w^* + \delta w^*g'(u^*) & 0 & -d_3\mu_k \end{pmatrix}, \quad k = 0, 1, 2, \dots \quad (6)$$

- 7 Hence, the characteristic equation for N_k is as follows:

$$\lambda^3 + A_1\lambda^2 + A_2\lambda + A_3 = 0, \quad (7)$$

- 8 where

$$\begin{aligned} A_1 &= (d_1 + d_2 + d_3)\mu_k + \eta + 1, \\ A_2 &= (d_1d_2 + d_1d_3 + d_2d_3)\mu_k^2 + [d_1 + d_2\eta + d_3(\eta + 1) + \rho w^*g(u^*) - \chi u^*]\mu_k \\ &\quad + [\eta + \delta w^*g(u^*)g'(u^*) - a], \\ A_3 &= d_1d_2d_3\mu_k^3 + [(d_1 + d_2\eta)d_3 + d_2\rho w^*g(u^*) - d_3\chi u^*]\mu_k^2 \\ &\quad + [d_3\eta + d_2\delta w^*g(u^*)g'(u^*) + \rho w^*g(u^*) - ad_3]\mu_k + \delta w^*g(u^*)g'(u^*). \end{aligned}$$

1 Firstly it is noted that $A_1 > 0$ for any $k \geq 0$. In addition, if

$$\chi < \chi_3, \quad a < \eta + \delta w^* g(u^*) g'(u^*), \quad (8)$$

2 then $A_2 > 0$ for any $k \geq 0$, where $\chi_3 = \frac{d_1 + d_2 \eta + d_3(\eta + 1) + \rho w^* g(u^*)}{u^*}$. And if

$$\chi \leq \chi_2, \quad a \leq \eta + \frac{d_2}{d_3} \delta w^* g(u^*) g'(u^*) + \frac{1}{d_3} \rho w^* g(u^*), \quad (9)$$

3 then $A_3 > 0$ for any $k \geq 0$, where χ_2 has been mentioned above.

Next, direct computations show that

$$A_1 A_2 - A_3 = B_1 \mu_k^3 + B_2 \mu_k^2 + B_3 \mu_k + B_4$$

4 where

$$\begin{aligned} B_1 &= (d_1 + d_2 + d_3)(d_1 d_2 + d_1 d_3 + d_2 d_3) - d_1 d_2 d_3, \\ B_2 &= (d_1 d_2 + d_1 d_3 + d_2 d_3)(\eta + 1) + d_1[d_1 + d_2 \eta + d_3(\eta + 1) + \rho w^* g(u^*) - \chi u^*] \\ &\quad + d_2[d_1 + d_2 \eta + d_3(\eta + 1) - \chi u^*] + d_3[d_3(\eta + 1) + \rho w^* g(u^*)], \\ B_3 &= \eta[d_1 + d_2 \eta + d_3(\eta + 1) + \rho w^* g(u^*) - \chi u^*] + d_1[\eta + \delta w^* g(u^*) g'(u^*) - a] \\ &\quad + d_2(\eta - a) + d_3 \delta w^* g(u^*) g'(u^*) + [d_1 + d_2 \eta + d_3(\eta + 1) - \chi u^*], \\ B_4 &= \eta[\eta + \delta w^* g(u^*) g'(u^*) - a] + \eta - a. \end{aligned}$$

5 According to the above formula, we can easily get $B_1 > 0$ for any $k \geq 0$; if
6 $\chi \leq \chi_1$, then $B_2 > 0$ holds; if $\chi \leq \chi_1$, and $a \leq \eta$, then $B_3 > 0$; if $a \leq \eta$, then
7 $B_4 > 0$. Therefore, $A_1 A_2 - A_3 > 0$ holds if $\chi \leq \chi_1$, and $a \leq \eta$, where χ_1, η have
8 been mentioned above.

9 In summary, if (8), (9) and $\chi \leq \chi_1, a \leq \eta$ hold, that is $\chi \leq \min\{\chi_1, \chi_2, \chi_3\} =$
10 $\min\{\chi_1, \chi_2\}$ and $a \leq \eta$ hold, then we have $A_1, A_2, A_3 > 0$ and $A_1 A_2 - A_3 > 0$.
11 According to Routh-Hurwitz criterion, if (4) and (5) hold, then (u^*, v^*, w^*) is
12 locally asymptotically stable.

13 Remark

14 When $\rho = 0$, we get $\chi_2 < \chi_1 = \chi_3$. Therefore, condition (5) can be changed
15 to $\chi \leq \chi_2$ and $a \leq \eta$, which makes the stability conditions more concise. When
16 $\rho \neq 0$, we have $\chi_1 < \chi_3$, but the relationship between χ_2 and χ_1, χ_3 cannot be
17 determined. It can be seen that ρ has a direct influence on the size relationship
18 of χ_1, χ_2 and χ_3 , also indirectly affects the stability conditions.

19 Compared with the case of $\chi = \rho = 0$, the stability of constant equilibria of
20 the model after introducing the prey-taxis is controlled by the prey-taxis rates.
21 The fluctuation of the value of χ has a direct impact on the stability.

22 3 Hopf bifurcation of problem (3)

23 In this section we are going to analyze the conditions about the parameters
24 under which Hopf bifurcation occurs near the unique positive constant solution

- 1 (u^*, v^*, w^*) of problem (3). We shall apply Theorem 6.1 of paper [31] to derive
 2 the emergence of Hopf bifurcation.

Denote

$$H = \{\chi_j^H : A_1 A_2(\chi_j^H) - A_3(\chi_j^H) = 0\},$$

- 3 where H is the Hopf bifurcation curve.

Theorem 3.1. Assume that (4) and $a \leq \eta$ hold. There exists $j \geq 1$ such that

$$\min\{\chi_1, \chi_2\} < \chi_j^H \leq \min\{\chi_2, \chi_3\}$$

holds, where χ_1, χ_2 and χ_3 have been mentioned in Theorem 2.1. Suppose that
 (H1) for some $j \in N$, μ_j is a simple eigenvalue of $-\Delta$ in Ω with Neumann
 boundary condition and the corresponding eigenfunction is $y_j(x)$;

(H2) for any $k \neq j$, $\chi_j^H \neq \chi_k^H$.

Then

1. (3) has a unique one-parameter family $\{\beta(s) : 0 < s < \varepsilon\}$ of nontrivial
 periodic orbits near $(\chi, u, v, w) = (\chi_j^H, u^*, v^*, w^*)$. More precisely, let $X = \{u \in$
 $H^2(\Omega) \mid \frac{\partial u}{\partial n}|_{\partial\Omega} = 0\}$, there exist $\varepsilon > 0$ and C^∞ function $s \mapsto (U_j(s), T_j(s), \chi_j(s))$
 from $s \in (-\varepsilon, \varepsilon)$ to $C^1(\mathbb{R}, X^3) \times (0, \infty) \times \mathbb{R}$ satisfying

$$(U_j(0), T_j(0), \chi_j(0)) = \left((u^*, v^*, w^*), \frac{2\pi}{\nu_0}, \chi_j^H \right),$$

and

$$U_j(s, x, t) = (u^*, v^*, w^*) + sy_j(x)[V_j^+ \exp(i\nu_0 t) + V_j^- \exp(i\nu_0 t)] + o(s),$$

where $\nu_0 = \sqrt{A_2}$, A_2 has been mentioned in (7), and V_j^\pm is an eigenvector
 satisfying $N_j V_j^\pm = i\nu_0 V_j^\pm$;

2. for $0 < |s| < \varepsilon$, $\beta(s) = \beta(U_j(s)) = \{U_j(s, \cdot, t) : t \in \mathbb{R}\}$ is a nontrivial periodic
 orbit of (3) of period $T_j(s)$;

3. if $0 < s_1 < s_2 < \varepsilon$, then $\beta(s_1) \neq \beta(s_2)$;

4. there exists $\tau > 0$ such that if (3) has a nontrivial periodic solution $\tilde{U}(x, t)$
 of period T for some $\chi \in \mathbb{R}$ with

$$|\chi - \chi_j^H| < \tau, \quad |T - \frac{2\pi}{\nu_0}| < \tau, \quad \max|\tilde{U}(x, t) - (u^*, v^*, w^*)| < \tau,$$

- 4 then $\chi = \chi_j(s)$ and $\tilde{U}(x, t) = U_j(s, x, t + \theta)$ for some $s \in (0, \varepsilon)$ and some $\theta \in \mathbb{R}$.

- 5 **Proof** We illustrate all the conditions listed in Theorem 6.1 of paper [31]
 6 one by one.

Step 1: we first show that, at $\chi = \chi_j^H$, there is $\omega_0 > 0$ such that $\pm i\omega_0$
 are simple eigenvalues of (7). By $\chi_j^H \leq \chi_2$ and $a \leq \eta$, it is easy to see that
 $A_3(\chi_j^H) > 0$. And due to $\chi_j^H < \chi_3$ and $a \leq \eta$, we have $A_2(\chi_j^H) > 0$. The roots
 of the characteristic equation (7) with $\chi = \chi_j^H$ are

$$\lambda_j = -A_1 < 0, \quad \lambda_j^\pm = \pm \sqrt{A_2(\chi_j^H)}.$$

Therefore, the matrix $N_j(\chi_j^H)$ defined in (6) admits a pair of purely imaginary eigenvalues $\pm i\sqrt{A_2(\chi_j^H)}$.

Step 2: we next show that, at $\chi = \chi_j^H$, (7) has not eigenvalues of the form $\pm iq\sqrt{A_2(\chi_j^H)}$ for $q \in N \setminus \{\pm 1\}$. Due to μ_j is a simple eigenvalue of $-\Delta$ and $\chi_j^H \neq \chi_k^H$ for $j \neq k$, the characteristic equation (7) has no root of the form $iq\sqrt{A_2(\chi_j^H)}$ with $q \in N^+ \setminus \{\pm 1\}$.

Step 3: finally, we prove that, for χ near χ_j^H , N_j has a unique eigenvalue $\sigma(\chi) + i\nu(\chi)$ such that $\sigma(\chi_j^H) = 0$, $\nu(\chi_j^H) > 0$ and $\sigma'(\chi_j^H) \neq 0$.

Let $\alpha(\chi)$ and $\sigma(\chi) \pm i\nu(\chi)$ be the three roots of (7) in a neighbourhood of χ_j^H . Clearly, $\alpha(\chi)$, $\sigma(\chi)$ and $\nu(\chi)$ are real analytic function of χ , and $\alpha(\chi_j^H) = -A_1 < 0$, $\sigma(\chi_j^H) = 0$, $\nu(\chi_j^H) = \sqrt{A_2(\chi_j^H)} > 0$.

Next, we show the transversality condition $\sigma'(\chi_j^H) \neq 0$.

Plugging $\alpha(\chi)$, $\sigma(\chi) \pm i\nu(\chi)$ into (7), we obtain

$$A_1 = -\alpha - 2\sigma, \quad A_2(\chi) = \sigma^2 + \nu^2 + 2\alpha\sigma, \quad A_3(\chi) = -\alpha(\sigma^2 + \nu^2). \quad (10)$$

By differentiating the three equations in (10) with respect to χ and using the definitions of A_1 , A_2 and A_3 , we have

$$\alpha' + 2\sigma' = 0, \quad 2\sigma\sigma' + 2\nu\nu' + 2\alpha'\sigma + 2\alpha\sigma' = -u^*\mu_j, \quad (11)$$

$$\alpha'\nu^2 + \alpha'\sigma^2 + 2\alpha\nu\nu' + 2\alpha\sigma\sigma' = d_3u^*\mu_j^2. \quad (12)$$

Note that $\sigma(\chi_j^H) = 0$. It follows from (11) and (12) that, at $\chi = \chi_j^H$,

$$2\nu\nu' + 2\alpha\sigma' = -u^*\mu_j, \quad 2\alpha\nu\nu' + \alpha'\nu^2 = d_3u^*\mu_j^2.$$

Observe that $\alpha(\chi_j^H) = -A_1$ and $\alpha'(\chi_j^H) = -2\sigma'(\chi_j^H)$. Thus, at $\chi = \chi_j^H$,

$$2\alpha\nu\nu' + 2\alpha^2\sigma' = 2\alpha\nu\nu' - \alpha'\alpha^2 = -\alpha u^*\mu_j,$$

$$2\alpha\nu\nu' + \alpha'\nu^2 = d_3u^*\mu_j^2,$$

which implies

$$\begin{aligned} \alpha'(\chi_j^H) &= \frac{(d_3\mu_j + \alpha)u^*\mu_j}{\alpha^2 + \nu^2} \Big|_{\chi=\chi_j^H} \\ &= \frac{(d_3\mu_j - A_1)u^*\mu_j}{A_1^2 + A_2} \Big|_{\chi=\chi_j^H}. \end{aligned}$$

Due to the definition of A_1 such that $d_3\mu_j - A_1 < 0$, so we obtain that $\alpha'(\chi_j^H) < 0$, and hence $\sigma'(\chi_j^H) = -\frac{1}{2}\alpha'(\chi_j^H) > 0$. This gives the transversality condition mentioned above.

Noticing that (3) is normally parabolic and steps 1 – 3 ensure the conditions $(H_1) - (H_3)$ given in Theorem 6.1 of [31], respectively. Our desired conclusions are deduced by Theorem 6.1 of [31].

Remark

1 In paper [12], the authors derived the emergence of Hopf bifurcation at
 2 $\gamma = \gamma^*$ by choosing γ as bifurcation parameter, where $\gamma^* = \frac{\zeta_0 - b - \frac{\delta}{r} u^* g'(u^*)(a-b)}{(2 - \frac{\delta}{r} u^* g'(u^*)) u^*}$.
 3 Compared with [12], we introduced the prey-taxis rates χ and ρ . By choosing
 4 χ as a bifurcation parameter, Hopf bifurcation can also be generated at χ_j^H .
 5 When $\rho = 0$, it can be accurately shown that periodic solutions will arise near
 6 χ_2 .

7 4 Steady state bifurcation of problem (3)

8 Note that the steady state solutions of (3) satisfy

$$\begin{cases} -d_1 \Delta u + \chi \nabla(u \nabla v) = av - bu - \gamma u^2 - g(u)w, & x \in \Omega, \\ -d_2 \Delta v = u - v, & x \in \Omega, \\ -d_3 \Delta w + \rho \nabla(w \nabla u) = w(-r + \delta g(u)), & x \in \Omega, \\ \frac{\partial u}{\partial n} = \frac{\partial v}{\partial n} = \frac{\partial w}{\partial n} = 0, & x \in \partial \Omega. \end{cases} \quad (13)$$

9 In this section, we prove the existence of nonconstant solutions of (13). In
 10 order to achieve this goal, we shall use χ as bifurcation parameter and apply
 11 the bifurcation theory in Theorem 4.3 of [37].

Denote

$$S = \{\chi_j^S : A_3(\chi_j^S) = 0\},$$

12 where S is the steady state bifurcation curve.

Theorem 4.1. *Assume that the parameters that condition (H1) in Theorem 3.1 is satisfied, and also*

(S1) for any $k \neq j$, $\chi_j^S \neq \chi_k^S$.

Then (13) has a unique one-parameter family $\Gamma_j = \{(\hat{U}_j(s), \hat{\chi}_j(s)) : -\varepsilon < s < \varepsilon\}$ of nontrivial solutions near $(u, v, w, \chi) = (u^, v^*, w^*, \chi_j^S)$. More precisely, there exist $\varepsilon > 0$ and C^∞ function $s \mapsto (\hat{U}_j(s), \hat{\chi}_j(s))$ from $s \in (-\varepsilon, \varepsilon)$ to $X^3 \times \mathbb{R}$ satisfying*

$$(\hat{U}_j(0), \hat{\chi}_j(0)) = ((u^*, v^*, w^*), \chi_j^S),$$

13 and

$$\begin{aligned} \hat{U}_j(s, x) &= (u^*, v^*, w^*) + sy_j(x) \left(d_2 \mu_j + 1, 1, \frac{(\rho w^* \mu_j + \delta w^* g'(u^*))(d_2 \mu_j + 1)}{d_3 \mu_j} \right) \\ &\quad + s(h_{1,j}(s), h_{2,j}(s), h_{3,j}(s)), \end{aligned}$$

14 such that $h_{1,j}(0) = h_{2,j}(0) = h_{3,j}(0) = 0$.

Proof Let $X = H^2(\Omega) = \{u \in H^2(\Omega) \mid \frac{\partial u}{\partial n}|_{\partial \Omega} = 0\}$, $Y = L^2(\Omega)$, $Y_0 = \{u \in L^2(\Omega) \mid \int_\Omega u(x) dx = 0\}$. Define a mapping $F : X^3 \times \mathbb{R} \rightarrow Y_0 \times Y^2 \times \mathbb{R}$ by

$$F(u, v, w, \chi) = \begin{pmatrix} d_1 \Delta u - \chi \nabla(u \nabla v) + av - bu - \gamma u^2 - g(u)w \\ d_2 \Delta v + u - v \\ d_3 \Delta w - \rho \nabla(w \nabla u) + w(-r + \delta g(u)) \end{pmatrix}.$$

1 We apply Theorem 4.3 of [37] to the equation $F(u, v, w, \chi) = 0$ at $(u^*, v^*, w^*, \chi_j^S)$.
2 Clearly, $F(u^*, v^*, w^*, \chi_j^S) = 0$, and F is continuously differentiable. We verify
3 the conditions in Theorem 4.3 of [37] in the following steps.
4 **Step 1** $F_U(u^*, v^*, w^*, \chi_j^S)$ is a Fredholm operator with index zero, and the kernel
5 space $N(F_U(u^*, v^*, w^*, \chi_j^S))$ is a one-dimensional, where $U = (u, v, w)$.

According to the Lemma 2.3 in [36], one can show that the linear operator
 $F_U(u^*, v^*, w^*, \chi_j^S) : X^3 \rightarrow Y_0 \times Y^2 \times \mathbb{R}$ is a Fredholm operator with index zero.
To prove that $N(F_U(u^*, v^*, w^*, \chi_j^S)) \neq \{0\}$, we calculate that

$$F_U(u^*, v^*, w^*, \chi_j^S)[\phi, \psi, \varphi] = \begin{pmatrix} d_1 \Delta \phi - \chi_j^S \nabla(u^* \nabla \psi) + a\psi - b\phi - 2\gamma u^* \phi - g(u^*)\varphi \\ d_2 \Delta \psi + \phi - \psi \\ d_3 \Delta \varphi - \rho \nabla(w^* \nabla \phi) + \varphi(-r + \delta g(u^*)) \end{pmatrix}.$$

Let $(\phi, \psi, \varphi) (\neq 0) \in NF_U(u^*, v^*, w^*, \chi_j^S)$, so

$$F_U(u^*, v^*, w^*, \chi_j^S)[\phi, \psi, \varphi] = 0.$$

6 The above equation has a non-zero solution, which is equivalent to that 0 is the
7 eigenvalue of N_j . It is easy to verify that when $\chi = \chi_j^S$, 0 is the eigenvalue of
8 N_j and the corresponding eigenfunction is

$$(\bar{a}_j, \bar{b}_j, \bar{c}_j)y_j = \left(d_2\mu_j + 1, 1, \frac{(\rho w^* \mu_j + \delta w^* g'(u^*))(d_2\mu_j + 1)}{d_3\mu_j} \right) y_j. \quad (14)$$

9 From the condition **(H1)**, the eigenvector is unique up to a constant multi-
10 ple. Thus one has $N(F_U(u^*, v^*, w^*, \chi_j^S)) = \text{span}\{(\bar{a}_j, \bar{b}_j, \bar{c}_j)y_j\}$, which is one-
11 dimensional.

12 **Step 2** $F_{\chi_U}(u^*, v^*, w^*, \chi_j^S)[(\bar{a}_j, \bar{b}_j, \bar{c}_j)y_j] \notin R(F_U(u^*, v^*, w^*, \chi_j^S))$.

13 We claim that the range space $R(F_U(u^*, v^*, w^*, \chi_j^S))$ can be characterized
14 as follows:

$$\begin{aligned} & R(F_U(u^*, v^*, w^*, \chi_j^S)) \\ &= \left\{ (h_1, h_2, h_3, \tau) \in Y_0 \times Y^2 \times \mathbb{R} : \int_{\Omega} (a_j^* h_1 + b_j^* h_2 + c_j^* h_3) y_j \, dx = 0 \right\}, \end{aligned} \quad (15)$$

where (a_j^*, b_j^*, c_j^*) is a non-zero eigenvector for the eigenvalue $\lambda = 0$ of N_j^T (the
transpose of N_j defined in (6)):

$$(a_j^*, b_j^*, c_j^*)y_j = \left(d_2\mu_j + 1, \chi_j^S \mu_j u^* + a, -\frac{g(u^*)(d_2\mu_j + 1)}{d_3\mu_j} \right) y_j.$$

Indeed if $(h_1, h_2, h_3, \tau) \in R(F_U(u^*, v^*, w^*, \chi_j^S))$, then there exists $(\phi_1, \psi_1, \varphi_1) \in$
 X^3 such that

$$F_U(u^*, v^*, w^*, \chi_j^S)[(\phi_1, \psi_1, \varphi_1)] = (h_1, h_2, h_3, \tau).$$

Define

$$L[\phi, \psi, \varphi] = \begin{pmatrix} d_1 \Delta \phi - \chi_j^S u^* \Delta \psi + a\psi - b\phi - 2\gamma u^* \phi - g(u^*)\varphi \\ d_2 \Delta \psi + \phi - \psi \\ d_3 \Delta \varphi - \rho w^* \Delta \phi + \varphi(-r + \delta g(u^*)) \end{pmatrix},$$

and its adjoint operator

$$L^*[\phi, \psi, \varphi] = \begin{pmatrix} d_1 \Delta \phi - b\phi - 2\gamma u^* \phi + \psi - \rho w^* \Delta \varphi \\ d_2 \Delta \psi - \chi_j^S u^* \Delta \phi + a\phi - \psi \\ d_3 \Delta \varphi - g(u^*)\phi + \varphi(-r + \delta g(u^*)) \end{pmatrix}.$$

Thus, we have

$$\begin{aligned} \langle (h_1, h_2, h_3), (a_j^*, b_j^*, c_j^*)y_j \rangle &= \langle L[(\phi_1, \psi_1, \varphi_1)], (a_j^*, b_j^*, c_j^*)y_j \rangle \\ &= \langle (\phi_1, \psi_1, \varphi_1), L^*[(a_j^*, b_j^*, c_j^*)y_j] \rangle \\ &= \langle (\phi_1, \psi_1, \varphi_1), N_j^*(a_j^*, b_j^*, c_j^*)y_j \rangle \\ &= 0, \end{aligned}$$

where $\langle \cdot, \cdot \rangle$ is the inner product in $[L^2(\Omega)]^3$. This illustrates that if

$$(h_1, h_2, h_3, \tau) \in R(F_U(u^*, v^*, w^*, \chi_j^S)),$$

then

$$\int_{\Omega} (a_j^* h_1 + b_j^* h_2 + c_j^* h_3) y_j \, dx = 0. \quad (16)$$

Due to (16) defines a codimension-1 set in $Y_0 \times Y^2 \times \mathbb{R}$, and we obtain that

$$\text{codim } R(F_U(u^*, v^*, w^*, \chi_j^S)) = \dim N(F_U(u^*, v^*, w^*, \chi_j^S)) = 1,$$

therefore, $R(F_U(u^*, v^*, w^*, \chi_j^S))$ must be in form of (15).

Now it's worth noting that

$$F_{\chi U}(u^*, v^*, w^*, \chi_j^S)[(\bar{a}_j, \bar{b}_j, \bar{c}_j)y_j] = (-u^* \bar{b}_j \Delta y_j, 0, 0) = (u^* \mu_j y_j, 0, 0),$$

then according to (15), we get

$$\begin{aligned} \int_{\Omega} (a_j^* h_1 + b_j^* h_2 + c_j^* h_3) y_j \, dx &= \int_{\Omega} u^* \mu_j y_j a_j^* y_j \, dx \\ &= \int_{\Omega} u^* \mu_j (d_2 \mu_j + 1) y_j^2 \, dx > 0. \end{aligned}$$

Hence, $F_{\chi U}(u^*, v^*, w^*, \chi_j^S)[(\bar{a}_j, \bar{b}_j, \bar{c}_j)y_j] \notin R(F_U(u^*, v^*, w^*, \chi_j^S))$. This conclusion has been proved.

5 Conclusions

In this paper, we study the dynamics of a three-component predator-prey model with prey-taxis and stage structure for the prey under the homogeneous Neumann boundary condition. The main contributions of the present paper consist of three parts: stability analysis of constant equilibria, Hopf bifurcation analysis and steady state bifurcation analysis. For the first problem, we mainly

1 use the eigenvalue method to analyze and obtain the stability of the constant
2 equilibria. We conclude that the sufficiently strong taxis effect χ destabilizes
3 the stability of the positive equilibrium regardless of the influence of another
4 taxis mechanism ρ ; for the second problem, choosing the prey-taxis sensitivity
5 coefficient as a bifurcation parameter, we get the existence of Hopf bifurcation;
6 for the third problem, the emergence of non-constant steady state is concluded
7 at a chemotactic parameter by the bifurcation theorem. Our conclusions show
8 that taxis rate χ plays an important role in determining the stability of the
9 interior equilibrium and influencing the existence of time-periodic patterns and
10 non-constant steady state. However, we shall not perform related numerical
11 simulations in the present framework and we leave it for the interested readers.

12 **Acknowledgment.** We would like to thank the anonymous referees for their
13 helpful comments and suggestions.

14 References

- 15 [1] A.J. Lotka. Elements of physical biology[M]. Williams and Wilkins. (1925).
- 16 [2] H. Malchow, S.V. Petrovskii, E. Venturino. Spatiotemporal patterns in ecol-
17 ogy and epidemiology: theory, models, and simulation[M]. (2007).
- 18 [3] C.S. Holling. The functional response of predators to prey density and its
19 role in mimicry and population regulation[J]. Memoirs of the Entomological
20 Society of Canada. 97(45): 5-60(1965).
- 21 [4] J.F. Andrews. A mathematical model for the continuous culture of microor-
22 ganisms utilizing inhibitory substrates[J]. Biotechnology and Bioengineering.
23 10(6): 707-723(1968).
- 24 [5] J.T. Tanner. The stability and the intrinsic growth rates of prey and predator
25 populations[J]. Ecology. 56(4): 855-867(1975).
- 26 [6] P.A. Braza. The bifurcation structure of the Holling-Tanner model for
27 predator-prey interactions using two-timing[J]. SIAM Journal on Applied
28 Mathematics. 63(3): 889-904(2002).
- 29 [7] J. Beddington. Mutual interference between parasites or predators and its
30 effect on searching efficiency[J]. Journal of Animal Ecology. 44(1): 331-
31 340(1975).
- 32 [8] D.L. DeAngelis, R.A. Goldstein, R.V. O'Neill. A model for trophic interac-
33 tion[J]. Ecology. 56(2): 881-892(1975).
- 34 [9] X.C. Zhang, G.Q. Sun, Z. Jin. Spatial dynamics in a predator-prey mod-
35 el with Beddington-DeAngelis functional response[J]. Physical Review E.
36 85(2): 021924(2012).

- 1 [10] R. Arditi, L.R. Ginzburg. Coupling in predator-prey dynamics: ratio-
2 dependence[J]. *Journal of Theoretical Biology*. 139(3): 311-326(1989).
- 3 [11] R.E. Kooij, A. Zegeling. A predator-prey model with Ivlev's functional
4 response[J]. *Journal of Mathematical Analysis and Applications*. 198: 473-
5 489(1996).
- 6 [12] S.H. Xu. Dynamics of a general prey-predator model with prey-stage struc-
7 ture and diffusive effects[J]. *Computers and Mathematics with Applications*.
8 68(3): 405-423(2014).
- 9 [13] S.M. Baer, B.W. Kooi, Y.A. Kuznetsov, H.R. Thieme. Multiparametric
10 bifurcation analysis of a basic two-stage population model[J]. *SIAM Journal*
11 *on Applied Mathematics*. 66(4): 1339-1365(2006).
- 12 [14] K.L. Cooke, R.H. Elderkin, W. Huang. Predator-prey interactions with de-
13 lays due to juvenile maturation[J]. *SIAM Journal on Applied Mathematics*.
14 66(3): 1050-1079(2006).
- 15 [15] S.A. Gourley, Y. Kuang. A stage structured predator-prey model and its
16 dependence on maturation delay and death rate[J]. *Journal of Mathematical*
17 *Biology*. 49(2): 188-200(2004).
- 18 [16] A. Hastings, Age-dependent predation is not a simple process. I. Continuous
19 time models[J]. *Theoretical Population Biology*. 23(3): 347-362(1983).
- 20 [17] S.Q. Liu, E. Berett. A stage-structured predator-prey model of Beddington-
21 Deangelis type[J]. *SIAM Journal on Applied Mathematics*. 66(4): 1101-
22 1129(2006).
- 23 [18] S.Q. Liu, L.S. Chen, R. Agarwal. Recent progress on stage-structured
24 population dynamics[J]. *Mathematical and Computer Modelling*. 36(11-13):
25 1319-1360(2002).
- 26 [19] C.H. Ou, J.H. Wu. Spatial spread of rabies revisited: Influence of age-
27 dependent diffusion on nonlinear dynamics[J]. *SIAM Journal on Applied*
28 *Mathematics*. 67(1): 138-163(2006).
- 29 [20] W.D. Wang, G. Mulone, F. Salemi, V. Salone. Permanence and stability of
30 a stage-structured predator-prey model[J]. *Journal of Mathematical Analysis*
31 *and Applications*. 262(2): 499-528(2001).
- 32 [21] N.X. Yan, L.S. Chen. Global stability of a predator-prey system with
33 stage structure for the predator[J]. *Acta Mathematica Sinica*. 20(001): 63-
34 70(2004).
- 35 [22] X. Rui, M. Chaplain, F.A. Davidson. Persistence and global stability of a
36 ratio-dependent predator-prey model with stage structure[J]. *Applied Math-*
37 *ematics and Computation*. 158(3): 729-744(2004).

- 1 [23] P. Kareiva, G. Odell. Swarms of predators exhibit preytaxis if individual
2 predators use area-restricted search[J]. The American Naturalist. 130(2):
3 233-270(1987).
- 4 [24] J.M. Lee, T. Hillen, M.A. Lewis. Pattern formation in prey-taxis systems[J].
5 Journal of Biological Dynamics. 3(6): 551-573(2009).
- 6 [25] J.P. Wang, M.X. Wang. Boundedness and global stability of the two-
7 predator and one-prey models with nonlinear prey-taxis[J]. Zeitschrift für
8 Angewandte Mathematik und Physik. 69(3): 63(2018).
- 9 [26] B.E. Ainseba, M. Bendahmane, A. Noussair. A reaction-diffusion system
10 modeling predator-prey with prey-taxis[J]. Nonlinear Analysis: Real World
11 Applications. 9(5): 2086-2105(2008).
- 12 [27] C. Hauzy, F.D. Hulot, A. Gins, M. Loreau. Intra- and interspecific density-
13 dependent dispersal in an aquatic prey-predator system[J]. The Journal of
14 Animal Ecology. 76: 552-558(2007).
- 15 [28] J.I. Tello, D. Wrzosek. Predator-prey model with diffusion and indirect prey-
16 taxis[J]. Mathematical Models and Methods in Applied Sciences. 26(11):
17 2129-2162(2016).
- 18 [29] J.P. Wang, M.X. Wang. Global solution of a diffusive predator-prey model
19 with prey-taxis[J]. Computers and Mathematics with Applications. 77(10):
20 2676-2694(2019).
- 21 [30] M. Winkler. Asymptotic homogenization in a three-dimensional nutrient
22 taxis system involving food-supported proliferation[J]. Journal of Differential
23 Equations. 263(8): 4826-4869(2017).
- 24 [31] P. Liu, J.P. Shi, Z. Wang. Pattern formation of the attraction-repulsion
25 Keller-Segel system[J]. Discrete and Continuous Dynamical Systems Series
26 B. 18(10): 2597-2625(2013).
- 27 [32] J.P. Wang, M.X. Wang. A predator-prey model with taxis mechanisms and
28 stage structure for the predator[J]. Nonlinearity. 33(7): 1-39(2020).
- 29 [33] P. Panja. Stability and dynamics of a fractional-order three-species
30 predator-prey model[J]. Theory in Biosciences. 138: 251-259(2019).
- 31 [34] L. Jia. Analysis for a delayed three-species predator-prey model with
32 feedback controls and prey diffusion[J]. Journal of Mathematics. 2020: 1-
33 26(2020).
- 34 [35] M. Sambath, C. Gokila. Stability and bifurcation analysis of three species
35 predator-prey model with Holling Type II[J]. Discontinuity, Nonlinearity,
36 and Complexity. 8(2): 127-144(2019).

- 1 [36] X. Wang, Q. Xu. Spiky and transition layer steady states of chemotaxis
2 systems via global bifurcation and Helly's compactness theorem[J]. Journal
3 of Mathematical Biology. 66(6): 1241-1266(2013).
- 4 [37] J.P. Shi, X.F. Wang. On global bifurcation for quasilinear elliptic systems
5 on bounded domains[J]. Journal of Differential Equations. 246(7): 2788-
6 2812(2009).