

Exploring the Critical Coronal Transition Region: The Key to Uncovering the Genesis of the Solar Wind and Solar Eruptions

Authors: A. Vourlidas, N. Viall, J. M. Laming, S. Cranmer, N. Arge, C. DeForest, G. de Toma, A. Caspi, N-E Raouafi

Endorsers: B. Alterman, N. Alzate, J. Burkepile, B. Chandran, G. Chintzoglou, C. Downs, S. Gibson, Y.-K. Ko, C. Korendyke, M. Linton, J. Mason, E. Mason, S. di Matteo, M.-P. Miralles, J. Newmark, T. Nieves-Chinchilla, J. Raymond, Y. Rivera, O.C. St Cyr, B. Thompson, S. Tun-Beltran, J. Qiu, S. Savage, D. Seaton, L. Strachan, M. West

The Current State

Long the source of fascination for humanity, the solar corona¹ has increasingly become a central target of Heliophysics research as space-based research became to dominate the field. The corona is the source of both the *ambient* and *eruptive* solar wind. Thus, the effects of physical processes that operate in the corona permeate every aspect of Heliophysics, from solar to space physics to the study of planetary magnetospheres, ionospheres, and atmospheres to the interaction of our heliosphere with interstellar space.

Table 1 Top-level Synopsis of the State of Knowledge of two Major Questions in Heliophysics as of 2020.

How and where does the solar wind form?	
Current Status (top-level)	Issues
<ul style="list-style-type: none"> - Large-scale magnetic field is responsible for the structure of the corona - General picture is that the SW is bimodal SW: Fast SW (FSW) from open field coronal holes . Slow SW (SSW) from everywhere else, including closed fields. - There are a plethora of observations of corona and solar wind that do not fit the bimodal SW picture - SW energization can come from waves, turbulence, reconnection, instabilities, or combinations of multiple processes. 	<ul style="list-style-type: none"> - No routine measurements of the physical properties (temperature, magnetic field, speed, composition, waves, turbulence, etc.) in the critical range $\sim 1.2 - 5 R_s$ - Reliance on mathematical extrapolations of the phot. field and ad-hoc prescriptions of coronal heating to model the corona - Small-scale structure and variability in the low atmosphere has wide impacts in the corona (S-web, .) - Difficulty in inward extrapolation through the cusp region and S-web obscures the source of the slow solar wind - Bimodal SW picture doesn't capture time-dynamics, 3D complexity, or multi-scale feedback that are critical to the formation of the solar wind
How do eruptions form?	
Current Status (top-level)	Issues
<ul style="list-style-type: none"> - CMEs (and flares) are coronal phenomena powered by the release of magnetic energy via reconnection - CMEs are ejections of magnetic flux ropes and develop within 1-15 R_s - CME acceleration/speed peak mostly $< 3R_s/20R_s$, resp. - Similar processes drive eruptions over a wide spatio-temporal range, from jets (arcsecs/mins) to streamer-blowouts ($> 90^\circ/\text{days}$) - High-energy SEPs accelerated below 10 R_s 	<ul style="list-style-type: none"> - Impossible to predict eruption due to incomplete understanding of energy accumulation, role of instabilities, triggers, etc. - CME magnetic content unknown due to: lack of B_{cor} measurements; incomplete understanding of CME inner corona evolution and force balance; no thermal information - Incomplete understanding of CME early evolution (rotation, deflection, compression) - No routine measurements of physical properties of shocks and plasma prevent closure on particle acceleration processes

It is no surprise, then, that much of the current research effort and hardware investment in Heliophysics revolves around variations of a single (two-sided) overarching question: *How does the solar wind (quiescent and explosive) form?* The question has remained open ever since the discoveries of the solar wind in 1962 and of Coronal Mass Ejections (CMEs) in 1971.

We have made progress in the last half-century but despite the energy and treasure expended on these problems, the fundamental questions on how the solar wind and CMEs form remain open. Table 1 attempts to summarize, at a high level, the current status. Evidently, both are complex problems but we know enough to envision a complete solution within the next 30 years. This White Paper (WP) describes the broad outlines of a 30-yr plan to resolve these issues. Linked WPs² with more detailed descriptions of some key science investigations are referenced where necessary.


Research ‘chokepoints’ and Path Forward

¹ We define the corona as the region between two physical interfaces of the solar atmosphere: from $\beta > 1$ to $\beta < 1$ interface (\sim upper chromosphere) to the sub- to super-Alfvénic solar wind interface ($\sim 20 R_s$, nominally).

² See WPs ‘CME Acceleration in the Middle Corona’ by Mason+ & ‘Observing Solar Analogs to Understand the Sun’ by Youngblood+

The issues in Table 1 readily identify several ‘research chokepoints’ that hinder progress:

- **Incomplete measurements.** This is the key chokepoint. The physical properties of the corona above ~ 1.2 Rs are rarely measured, particularly temperature, magnetic field, flow speed, and composition. The various regions are covered by different instruments in different wavelengths (low corona in the EUV, middle corona in the visible). The photospheric magnetic field is usable only within 60° from central meridian.
- **Multi-scale processes and feedback³:** coronal physics encompasses a large range of temporal and spatial scales crossing several physical interfaces. The system is interconnected with feedback from small to large scales and vice versa, including long-range interactions.
- **Fragmented community:** The corona is observed across a range of wavelengths (radio to X-rays) and with different instrumentation (both remote and in-situ) not always operating concurrently. This has fragmented the community into distinct plasma regimes, largely clustering around regions where existing instrumentation has made observations widely available and where models can be sufficiently self-contained to be tractable. This disunity precipitates stovepiping of research topics, stifling cross-disciplinary innovation required to understand the system as a whole.



Achieving Closure by 2050

To accelerate scientific discovery, a systems approach is required with coordinated development along three axes:

- Better tools (e.g. data assimilation, big data techniques, model dissemination, virtual repositories)⁴
- Better measurements (e.g. the right measurements from the right location for the right duration)⁵
- United community (e.g. unify the research language, expand premise of DRIVE Science Centers, enhance interdisciplinary & international collaborations, train system-minded researchers).

These developments feed and strengthen the two pillars of the research enterprise—observations and theory/modeling---through a chain of interlinked science investigations as outlined in our indicative 30-year plan in Table 2.

³ Details in submitted WP ‘Mesoscale dynamics: the key to unlocking the universal physics of multiscale feedback’ by Kepko+

⁴ See submitted WP ‘The Sun-Earth Connection as a Single System: Data Analysis and Processing Needs of Current and Future Missions’ by Alzate+ for details

⁵ See submitted WP “Solving the Space Weather Problem” by Vourlidis+ for an example

Table 2 Indicative 30-year plan for Achieving Closure on the Outstanding Scientific Problems of Solar Wind and CME Formation

Goals	Questions	Investigation	0-10 years	Goals	Investigations	10-20 years	Goals	20-30 years	Goals
How and where does the Solar Wind Form? ⁶	- Where does the SW originate? - How is the SW released and accelerated? - What determines the SW composition and ionic state? - What is the origin & evolution of the mesoscale plasma & magn. field structure? - What is the origin of Alfvénic fluctuations? - How is SW turbulence driven and dissipated? - How do the SW kinetic distribution functions evolve?	- Trace magn. & plasma connectivity from phot through the corona - Measure physical properties of CCTR - Weed-out theories of SW heating/accel. (test predictions of SW heating/accel theories) - Advance coronal modeling - Investigate the wave/turb. properties of coronal plasma - Develop coronal magnetometry methods	- Connect abundance variation in the Sun and solar wind - Link small-scale structures from chrom. Through corona to heliosphere - Improve physical description of the corona - Map the wave properties of coronal plasma - Data-driven time-dependent (i.e., continuously updated boundary cond.) coronal modeling	- Trace energy & plasma flow throughout the corona - Trace magnetic connectivity throughout the corona	- First polar coverage of B-field & corona ⁷ - First sustained multi-height B-field ⁷ through the $\beta=1$ layer. - 2π corona/B-field coverage ⁷	- Achieve full coverage of the Sun in a few key regimes - Closure on the two overarching objectives. - Move towards prediction of solar eruptive activity - Move towards integration of solar knowledge to stellar systems			
	Tools	- Ingest/analyze multi-platform/view data - Ingest multi-view B-field data - Deploy Big Data methods - Begin 'fusing' remote/in-situ methodologies - Facilitate access/use of advanced models by the community		- Long booms: EUV/UV coatings; imag. Spectropolarimetry - High resolution X-ray imaging - AI approaches to long-term solar dbases - Practical multi-scale kinetic/MHD sims - Efficient time-dep data-driven sims - Fuse remote/in-situ methodology		Routine data assimilation			
	Measurements	- Physical properties of coronal and CME plasma with multiple instruments (e.g., PSP-SO-DKIST) - Measure coronal magnetic field - Off-limb spectroscopy, high-res imaging (VIS/EUV), - First polar, off Sun Earth line magnetic field - Expand remit of Drive Science Centers; FSTs	- Identify reliably SEP origins in indiv. events - Remote measurements of 'seed' populations - Improve coronal shock modeling - Advance description of forces acting on CMEs - quantitative B_{CME} estimates	- 2π B_{phot} coverage - High-sens. Spectroscopy for 'seed' detection - Multi-height B measurements - Time/space-resolved 3D imaging (poln or tomogr.)	- Establish 'Corona-as-a-system research' program		4π B_{phot} + corona coverage		
	Community	- What are the roles of ideal/non-ideal processes in CME eruption? - Where does the CME magn. field originate? - What is the energy budget and distribution in eruptions? - How do coronal shocks form and accelerate particles? - How do CMEs escape?	- particle acceleration & shock development <10 Rs - 3D CME evolution <10Rs - Eruption Energy budgets - Estimate CME magn. content - Feedback to surface (X-rays, γ -rays, CME effects on global B-field)	- Measure coronal currents in ARs - Semi-operational coronal MHD modeling - Detailed description of CME dynamics <10Rs	- Deploy CME prediction methods - Reliable mapping of B_{cor} to Alfvénic zone - Operational coronal modeling				
How do CMEs form?									

⁶ See submitted white paper by Viall+ for details

⁷ See submitted white papers on 4π /polar perspectives by Gibson+, Hassler+, Vourlidas+, Newmark+ for details