

<div> <div></div> <div>Mosquitoes</div> </div> <div>Microsporidia</div>	<i>A. cinereus</i>	<i>A. vexans</i>	<i>C. richiardii</i>	<i>O. annulipes</i>	<i>O. cantans</i>	<i>O. punctor</i>	<i>O. sticticus</i>	All	Prevalence (%) of microsporidian species	
	92.3% (n = 13)	63.2% (n = 19)	56.3% (n = 16)	54.0% (n = 63)	57.1% (n = 77)	72.7% (n = 11)	61.5% (n = 13)	59.9% (n = 212)	in all	in M-P
<i>Amblyospora salinaria</i>							1	1	0.47	0.67
<i>Amblyospora</i> sp.				1	1		1	3	1.42	2.00
<i>Encephalitozoon hellem</i>					2			2	0.94	1.33
<i>Enterocytozpora artemiae</i>				1				1	0.47	0.67
Microsporidium sp. PL01	12	10	9	31	27	8	7	104	49.1	69.33
<i>Nosema adaliae</i>	1			1		1		3	1.42	2.00
<i>N. ceranae</i>					2			2	0.94	1.33
<i>N. chrysorrhoeae/portugal</i>		2	1	4	16			23	10.8	15.33
<i>N. pieriae</i>			1	2				3	1.42	2.00
<i>N. thomsoni</i>		1						1	0.47	0.67
<i>Nosema</i> sp. CHW-2007a		2	1	2	1		1	7	3.30	4.67