

Table 1. Effect of risk groups on demographic and operative outcomes

| | 1 Risk factor (n=1073) | 2 or 3 Risk factors (n=446) | p value |
|--|---------------------------------------|--|--------------------|
| Age (yr) | 62.6±6.6 | 64.2±6.9 | 0.001 |
| 40-59 | 314 (29.8%) | 104 (23.7%) | 0.001 |
| 50-69 | 585 (55.5%) | 233 (53.1%) | |
| >=70 | 156 (14.8%) | 102 (23.2%) | |
| BMI | 27±3.5 | 27.9±3.4 | 0.002 |
| <25 | 107 (30.7%) | 56 (19.9%) | 0.009 |
| 25-29.9 | 173 (49.6%) | 160 (56.7%) | |
| >=30 | 69 (19.8%) | 66 (23.4%) | |
| Operation time (min) | 178.8±76.3 | 190.9±69.8 | 0.008 |
| Blood loss (ml) | 409.2±424 | 295.9±401 | 0.001 |
| Operation type | | | 0.001 |
| Open | 693(64.6%) | 151(33.9%) | 0.001 |
| Laparoscopic | 171(15.9%) | 59(13.2%) | |
| Robotic | 209(19.5%) | 236(52.9%) | |
| PLND | | | 0.001 |
| Yes | 567 (53.6%) | 403 (94.2%) | 0.001 |
| No | 490 (46.4%) | 25 (5.8%) | |
| PLND type | | | 0.001 |
| Obturator fossa | 205 (36.1%) | 64 (15.9%) | 0.001 |
| Extended | 355 (62.6%) | 338 (83.9%) | |
| Lymph node positivity | 35 (10.1%) | 97 (28%) | 0.001 |
| Catheter removal time (day) | 10.8±2.5 | 9.8±6.7 | 0.005 |
| Clavien complication | | | |
| 1 | 15 | 6 | |
| 2 | 3 | 0 | |
| 3a | 5 | 1 | |
| 3b | 4 | 8 | |
| 4a | 1 | 0 | |
| 5 | 1 | 0 | |