

Table 1: Demographic characteristics of 1093 pregnant women randomly assigned to undergo an additional ultrasound examination at 35-37 weeks' gestation according to follow up status

Abbreviations: y, years; IQR, interquartile range; SD, standard deviation.

| Variables | Completed (n = 1061) | Lost (n = 32) | <i>p</i> value |
|--|-------------------------|------------------|----------------|
| Maternal age (y) [median(IQR)] | 30 (26-34) | 27.5 (25-31.5) | 0.02 |
| Maternal height (m) mean \pm SD | 1.63 \pm 0.06 | 1.64 \pm 0.06 | 0.50 |
| Maternal weight at beginning of pregnancy (Kg) [median(IQR)] | 61 (55-70) | 60 (56-70) | 0.86 |
| Body mass index at beginning of pregnancy (Kg/m ²) [median(IQR)] | 23 (20.7-26.2) | 22.8 (20.9-25.3) | 0.90 |
| Increase in weight during pregnancy at randomization (Kg) [median(IQR)] | 11 (9-15) | 10 (8-14) | 0.65 |
| Parity (n [%]) | | | 0.61 |
| Nulliparous | 573 (54) | 13 (41) | |
| Multiparous | 488 (46) | 19 (59) | |
| Marital status (n [%]) | | | 0.08 |
| Single | 295 (28.1) | 14 (43.7) | |
| Married | 443 (42.2) | 6 (18.8) | |
| Co-habitant | 299 (28.4) | 12 (37.5) | |
| Divorced | 13 (1.2) | 0 (0) | |
| Widowed | 1 (0.1) | 0 (0) | |
| Ethnicity (n [%]) | | | 0.30 |
| White | 939 (88.5) | 26 (81.2) | |
| Black | 116 (10.9) | 6 (18.8) | |
| Mixt | 2 (0.19) | 0 (0) | |
| Asian | 4 (0.38) | 0 (0) | |
| Education (n [%]) | | | 0.06 |
| Doctoral level | 7 (0.7) | 0 (0) | |
| Master level | 44 (4.2) | 0 (0) | |
| Degree level | 337 (32) | 8 (25) | |
| High school (12th grade) | 337 (32) | 11 (34.4) | |
| Middle school (9th grade) | 256 (24.3) | 6 (18.7) | |
| Elementary school (4th grade) | 62 (5.9) | 7 (21.9) | |
| Less than elementary school | 10 (0.9) | 0 (0) | |
| Smoker at randomization (n [%]) | 164 (15.5) | 6 (18.8) | 0.61 |