

Table 3. The evaluation of the relationship between comorbidity, major surgery history, smoking and alcohol use and STAI-I, STAI-II and Beck Depression Inventory Scores of 24 Patients whose operation were postponed

Parameters	STAI-I	STAI-II	Beck Depression Inventory
Yes/No (n)	Median (min-max)	Median (min-max)	Median (min-max)
HT			
Yes (13)	47 (44-49)	40 (39-46)	3 (0-11)
No (11)	47 (45-49)	41 (40-47)	5 (0-13)
p^m	0.699	0.098	0.178
DM			
Yes (13)	47 (45-48)	41 (39-47)	5 (0-13)
No (11)	47 (44-49)	41 (40-46)	2 (0-11)
p^m	1	0.976	0.482
CVD			
Yes (9)	47 (46-49)	41 (39-47)	5 (1-13)
No (15)	46 (44-49)	41 (40-46)	3 (0-11)
p^m	0.118	0.465	0.148
Major Surgery History			
Yes (11)	46 (44-48)	42 (39-47)	5 (1-13)
No (13)	47 (45-49)	41 (40-46)	4 (0-11)
p^m	0.102	0.723	0.725
Smoking			
Yes (5)	47 (45-49)	41 (40-46)	7 (1-11)
No (19)	47 (44-49)	41 (39-47)	3 (0-13)
p^m	0.342	0.885	0.131
Alcohol			
Yes (5)	47 (44-49)	41 (40-46)	5 (0-7)
No (19)	47 (45-49)	41 (39-47)	4 (0-13)
p^m	0.201	0.885	0.971

p^m : Mann-Whitney U test, STAI: State-Trait Anxiety Inventory, HT: Hypertension, DM: Diabetes Mellitus, CVD: Cardiovascular Disease