

Table 2. Distribution of attacks according to the sources of infection and antimicrobial treatment preferences

Infections	Number of patients	Ratio (%)	Number of infection attacks	Ratio (%)	Monotherapy	Ratio (%)	Two combined therapy	Ratio (%)	Three combined therapy	Ratio (%)
Cholangitis	37	19.47	42	16.34	41	15.95	1	0.4	0	0
Intraabdominal abscess	32	16.85	74	28.79	45	17.5	20	7.81	9	3.5
Infected bilioma	28	14.74	28	10.89	23	8.94	5	1.95	0	0
Cystitis	22	11.58	22	8.56	21	8.17	1	0.4	0	0
Sepsis	18	9.47	26	10.11	13	5.05	6	2.34	6	2.34
Surgical site infection	17	8.95	17	6.61	16	6.22	1	0.4	0	0
Pneumonia	15	7.89	16	6.22	9	3.5	7	2.73	0	0
Acute pyelonephritis	12	6.32	22	8.56	18	7.0	3	1.17	1	0.4
Pancreatitis	5	2.63	6	2.34	6	2.34	0	0	0	0
COVID-19	1	0.5	1	0.38	1	0.38	0	0	0	0
Fasciola hepatica	1	0.5	1	0.38	1	0.38	0	0	0	0
Influenza and cholangitis	1	0.5	1	0.38	0	0	1	0.4	0	0
Catheter infection	1	0.5	1	0.38	0	0	1	0.4	0	0
TOTAL	190	100.00	257	100.00	195	75.89	46	17.89	16	6.22