

**Table 1** Demographic and clinical characteristics of all tumor thrombus patients treated with and without anticoagulation

Characteristic	Total patients Number (%) (N=50)	Patients w/ anticoagulation Number (%) (N=10)	Patients without anticoagulation Number (%) (N=40)
<b>Age</b>			
Less than 1 year	5 (10%)	0	5 (12.5%)
1-12 year	30 (60%)	5 (50%)	25 (62.5 %)
13-21 year	15 (30%)	5 (50%)	10 (25%)
<b>Sex</b>			
Male	34 (68%)	5 (50%)	29 (72.5%)
Female	16 (32%)	5 (50%)	11 (27.5%)
<b>Race/Ethnicity</b>			
Caucasian	15 (30%)	2 (20 %)	13 (32.5%)
African-American	5 (10%)	2 (20%)	3 (7.5%)
Hispanic	23 (46%)	3 (30%)	20 (50%)
Asian	7 (14%)	3 (30%)	4 (10%)
<b>Diagnosis</b>			
Wilms Tumor	6 (12%)	2 (20%)	4 (10%)
Hepatoblastoma	13 (26%)	0	13 (32.5%)
Neuroblastoma	3 (6%)	1 (10%)	2 (5%)
Ewing sarcoma	3 (6%)	1 (10%)	2 (5%)
Osteosarcoma	7 (14%)	2 (20%)	5 (12.5%)
Other sarcomas	7 (14%)	3 (30%)	4 (10%)
Other tumors	11 (22%)	1 (10%)	10 (25%)
<b>Imaging modality</b>			
CT	25 (50%)	6 (60%)	19 (47.5%)
MRI	20 (40%)	4 (40%)	16 (40%)
US	6 (12%)	0	6 (15%)
Echocardiography	10 (20%)	4 (40%)	6 (15%)
CT Angiogram	2 (4%)	2 (20%)	0
<b>Tumor thrombus</b>			
Involvement of IVC	18 (36%)	8 (80%)	10 (40%)
Involvement of heart	12 (24%)	6 (60%)	6 (15%)