

Table 1. Demographic and clinical parameters of two groups

Parameters	Group 1 (n=73)	Group 2 (n=61)	p value
Age	46,25±12,87	45,51±14,36	0,75
BMI	28,38 (21,51-37,89)	27,34 (20,20-47,56)	0,36
Preoperative Hb (g/dL)	14,50 (9,1-17,9)	14,30 (10,6-17)	0,72
Postoperative Hb (g/dL)	13,05 (7,9-17,0)	13,3 (9,6-15,9)	0,98
Preoperative Cre (mg/dL)	0,94 (0,6-2,63)	0,89 (0,47-2,24)	0,22
Postoperative Cre (mg/dL)	0,87 (0,51-1,75)	0,83 (0,49-1,48)	0,33
Preop vs Postop Hb**	-6,6 (<0,001)	-5,7 (<0,001)	
Preop vs Postop Cre**	-3,2 (0,001)	-3,7 (<0,001)	
Hydronephrosis (n, %)		0,67	
Absent	14 (19,2)	15 (24,6)	
Grade 1	21 (28,8)	21 (34,4)	
Grade 2	33 (45,2)	22 (36,1)	
Grade 3	4 (5,5)	3 (4,9)	
Grade 4	1 (1,4)	0 (0)	
Stone size (mm)	9 (5-16)	10 (5-24)	0,54
Stone density (HU)	986 (480-1428)	900 (254-1632)	0,06
Operation time (dk)	46,97±20,51	48,54±27,98	0,70
Stone free (n, %)			0,03
Residual +	3 (4,2)	9 (15,0)	
Residual -	69 (95,8)	51 (85,0)	
Complication (n, %)			0,87
Absent	56 (77,8)	46 (76,7)	
Present	16 (22,2)	14 (23,3)	

Data were given as mean ± SD in cases with normal distribution, and as median (min-max) in data that did not show normal distribution. For categorical data, it was shown as n (%). BMI: Body mass index; HU: Hounsfield unit; mm: millimeters. * p <0.05 ** It is shown as “Z value (p value)” for Wilcoxon test.