

Strategies	Strategy A (National)		Strategy B (Rural)		Strategy C (Urban)		
Access to CEmOC (%)	50.2		47.2		55.7		
CS rate (%)	17.2		12.8		28.2		
Differences between strategies	Strategy A (Baseline)		Change from Strategy A to B (B-A)		Change from Strategy A to C (C-A)		
	Per 10,000 women	Per total population	Per 10,000 women	Per total population	Per 10,000 women	Per total population	
	Pregnancies to full term	646	20,998,079	+2.1	+68,735	-0.2	-7,743
	Vaginal Deliveries	522	16,957,079	+47	+1,539,119	-84	-2,732,176
	Cesarean Sections	173	5,628,779	-58	-1,888,050	+116	+3,764,628
	Neonates that survive	621	20,180,625	-0.1	-4,780	+6.68	+217,006
	Neonates that die	10	334,498	+2.2	+71,934	-6.9	-224,572
	Maternal mortality	0.5	15,296	-0.12	+875,673	-18.7	-6,692
Maternal mortality per 100,000 live birth	75		-19		33		
Neonatal mortality per 1,000 live births	16		3		-11		
Complications associated with increased CS rates							
Previa	0.5	16,252	-0.09	-2,868	+0.38	+12,428	
Accreta	0.4	13,384	-0.06	-1,912	+0.38	+12,428	
Hysterectomy	0.3	10,516	+8.97	+291,572	+0.24	+7,648	
ICU admissions	0.1	2,868	-0.09	-2,772	+0.03	+956	
Uterine Rupture	0.4	117,585	+3.73	+15,296	+3.4	+4,780	
PPH requiring transfusion	2.2	701,685	+19.93	+16,252	+20	+34,415	
Complications associated with lack of access to CEmOC care							
Fistula	0.003	956	+0.03	0.0	0.0	-860	
Incontinence	0.009	2,868	+0.08	0.0	+0.1	0.0	
Stroke	0.021	6,692	+0.16	-956	+0.2	0.0	
Uterine Rupture	0.4	117,585	+3.73	+15,296	+3.4	+4,780	
PPH requiring transfusion	2.2	701,685	+19.93	+16,252	+20	+34,415	