

RESEARCH ARTICLE

Fuzzy Laplace-Adomian Decomposition method for solving Fuzzy Klein-Gordan equations.

Kishor A. Kshirsagar^{*1} | Vasant R. Nikam² | Shivaji A. Tarate¹ | Shrikisan B. Gaikwad¹

¹Department of Mathematics, New Arts, Commerce and Science College, Ahmednagar, Maharashtra, India

²Department of Mathematics, Loknete Vyankatrao Hiray Arts, Science and Commerce College, Nashik, Maharashtra, India

Correspondence

*Kishor A. Kshirsagar, New Arts, Commerce and Science College, Ahmednagar. Email: kishorkshirsagar@newartscs.ac.in

Present Address

New Arts, Commerce and Science College, Ahmednagar.

Summary

In this article, we have use Fuzzy Laplace-Adomian Decomposition method to evaluate the approximate solution of non linear and linear Time Fractional Klein-Gordan equations with suitable initial conditions in Fuzzy environment. We apply Fuzzy Laplace Transform and iterative method under caputo fractional derivative.

KEYWORDS:

Fuzzy partial differential equations, Fractional derivatives and integrals, Fuzzy Laplace Transform, Fuzzy Fractional Klein- Gordan equation

1 | INTRODUCTION

For describing uncertainty and interpreting ambiguous or subjective data in mathematical models, fuzzy set theory is a useful tool^{1,2,3,4,5}. Fuzzy differential equations (FDEs) are an interesting issue from a theoretical standpoint; Fei⁶ studied fuzzy random differential equations (FRDEs) with non-Lipschitz coefficients. 1986's Kaleva⁷ presented fuzzy differential equations. Kaleva⁸ introduced the FDE Cauchy issue. Lopez⁹ compared FDEs. Friedman and Kandel¹⁰ solved FDEs numerically. Mizukoshi et al.¹¹ explored FDEs. Chang and Zadeh¹² introduced fuzzy derivatives in 1972. Dubois and Prade¹³ introduced extension.

This work develops a method for approximating nonlinear PDE solutions. LADM combines ADM and Laplace transformations.^{14,15,16,17,18} successfully employed this strategy. Adomian's^{19,20} ADM has been used in biology, physics, and chemical processes. The method provides a convergent series that may be computed quickly. To solve the differential equation apply the Laplace transformation and decomposing nonlinear terms into Adomian polynomials generates a recursive, iterative technique.

In recent decades there has appeared a big number of nonlinear evolution equations^{21,22}. To solve the fuzzy heat equation, Allahviranloo used the Adomian decomposition technique (ADM)²³. Alikhani and Bahrami²⁴ presented Fuzzy number cross products for FPDEs. Jameel et al.²⁵ proposed a new approximation method for fuzzy heat equations. To solve partial differential equations, Stepnicka and Valasek were the first to employ a fuzzy transform (PDEs)²⁶. Finally, Jameel studied the heat equation's semianalytical solution in a fuzzy environment²⁷.

The Fuzzy Fractional Klein-Gordan equation²⁸

$$\frac{\partial^\alpha \tilde{\mathfrak{N}}(b, \wp; \delta)}{\partial \wp^\alpha} + \frac{\partial^2 \tilde{\mathfrak{N}}(b, \wp; \delta)}{\partial b^2} + a \tilde{\mathfrak{N}}(b, \wp; \delta) + b \tilde{\mathfrak{N}}^2(b, \wp; \delta) + c \tilde{\mathfrak{N}}^3(b, \wp; \delta) = \tilde{k}(\delta) \mathfrak{N}(b, \wp; \delta) \quad (1)$$

⁰ **Abbreviations:** FDE, Fuzzy Differential equations; FRDE, Fuzzy Random Differential equations ; PDE, Partial Differential Equations; LADM, Laplace Adomian Decomposition Method; FPDE, Fuzzy Partial Differential equations

subject to the fuzzy initial condition as

$$\tilde{\mathfrak{N}}(b, 0; \delta) = \tilde{\mathfrak{N}}_0 \quad \text{and} \quad \frac{\partial \tilde{\mathfrak{N}}(b, 0; \delta)}{\partial \wp} = \tilde{\mathfrak{N}}_1 \quad (2)$$

where $\mathfrak{N}(b, \wp; \delta)$ be function of fuzzy number value and α is a parameter that describes the order of the fractional derivatives in Caputo sense. a, b, c are constants, $\tilde{k}(\delta)$ is a fuzzy number.

2 | IMPORTANT TERMINOLOGY

Here, we present important definitions, theorems, and Lemmas of fractional calculus and popular operators in fuzzy environments.

Definition 1. ²⁹ The order $\alpha > 0$ Riemann-Liouville integral operator \mathcal{J}^α is defined as

$$\mathcal{J}^\alpha g(\wp) = \frac{1}{\Gamma(\alpha)} \int_0^\wp (\wp - \mu)^{\alpha-1} g(\mu) d\mu, \quad \wp > 0. \quad (3)$$

Definition 2. ^{30,31} In the case of the fuzzy number-valued function g , the R-L fractional integral of order α can be represented as follows:

$$\left[\mathcal{J}^\alpha \tilde{\mathfrak{N}}(\wp; \delta) \right] = \left[\mathcal{J}^\alpha \underline{\mathfrak{N}}(\wp; \delta), \mathcal{J}^\alpha \bar{\mathfrak{N}}(\wp; \delta) \right], \quad \wp > 0, \quad (4)$$

where

$$\mathcal{J}^\alpha \underline{\mathfrak{N}}(\wp) = \frac{1}{\Gamma(\alpha)} \int_0^\wp (\wp - \mu)^{\alpha-1} \underline{\mathfrak{N}}(\mu) d\mu, \quad \wp > 0,$$

$$\mathcal{J}^\alpha \bar{\mathfrak{N}}(\wp) = \frac{1}{\Gamma(\alpha)} \int_0^\wp (\wp - \mu)^{\alpha-1} \bar{\mathfrak{N}}(\mu) d\mu, \quad \wp > 0.$$

Definition 3. ²⁹ The fractional derivative of $g(\wp)$ in the Caputo sense is defined as follows:

$${}^c D_\wp^\alpha g(\wp) = \mathcal{J}^{\zeta-\alpha} D_\wp^\zeta g(\wp) = \begin{cases} \frac{1}{\Gamma(\zeta-\alpha)} \int_0^\wp \frac{g^\zeta(\mu) d\mu}{(\wp-\mu)^{\alpha+1-\zeta}} & \zeta-1 < \alpha < \zeta, \zeta \in \mathbb{N} \\ \frac{d^\zeta}{d\wp^\zeta} g(\wp) & \alpha = \zeta, \zeta \in \mathbb{N} \end{cases} \quad (5)$$

Definition 4. ^{30,31} Let $[\tilde{\mathfrak{N}}(\wp; \delta)] = [\underline{\mathfrak{N}}(\wp; \delta), \bar{\mathfrak{N}}(\wp; \delta)]$ be a parametric representation of fuzzy valued function $\tilde{\mathfrak{N}}(\wp; \delta)$, where $\delta \in [0, 1], 0 < \alpha < 1$ and $\wp \in (a, b)$.

(i) If $\tilde{\mathfrak{N}}(\wp; \delta)$ is Caputo-type fuzzy fractional differential function in the first form can be expressed in the following way:

$$\left[{}^c D_\wp^\alpha \tilde{\mathfrak{N}}(\wp; \delta) \right] = \left[{}^c D_\wp^\alpha \underline{\mathfrak{N}}(\wp; \delta), {}^c D_\wp^\alpha \bar{\mathfrak{N}}(\wp; \delta) \right].$$

(ii) If $\tilde{\mathfrak{N}}(\wp; \delta)$ is Caputo-type fuzzy fractional differential function in the second form can be expressed in the following way:

$$\left[{}^c D_\wp^\alpha \tilde{\mathfrak{N}}(\wp; \delta) \right] = \left[{}^c D_\wp^\alpha \bar{\mathfrak{N}}(\wp; \delta), {}^c D_\wp^\alpha \underline{\mathfrak{N}}(\wp; \delta) \right].$$

where,

$${}^c D_\wp^\alpha \underline{\mathfrak{N}}(\wp; \delta) = \frac{1}{\Gamma(\zeta-\alpha)} \int_0^\wp \frac{\underline{\mathfrak{N}}^\zeta(\mu) d\mu}{(\wp-\mu)^{\alpha+1-\zeta}}, \quad \zeta-1 < \alpha < \zeta, \quad \zeta \in \mathbb{N}$$

$${}^c D_\wp^\alpha \bar{\mathfrak{N}}(\wp; \delta) = \frac{1}{\Gamma(\zeta-\alpha)} \int_0^\wp \frac{\bar{\mathfrak{N}}^\zeta(\mu) d\mu}{(\wp-\mu)^{\alpha+1-\zeta}}, \quad \zeta-1 < \alpha < \zeta, \quad \zeta \in \mathbb{N}$$

Definition 5. ^{32,31} The Laplace transform of fuzzy function g is expressed as

$$\mathfrak{N}(p) = \mathfrak{L}\left[\tilde{\mathfrak{N}}(\wp; \delta)\right](p) = \int_0^{\infty} \text{Exp}(-p\wp) \odot \tilde{\mathfrak{N}}(\wp) d\wp, \quad p > 0 \text{ and integer.} \quad (6)$$

Definition 6. ³¹ The transform Fuzzy Laplace for Caputo Type Fuzzy Fractional Derivative is

$$\mathfrak{L}\left[{}^c D_{\wp}^{\alpha} \tilde{\mathfrak{N}}(\wp; \delta)\right](p) = p^{\alpha} \mathfrak{N}(p) - \sum_{q=0}^{\zeta-1} p^{\alpha-q-1} \tilde{\mathfrak{N}}^q(b, 0; \delta), \quad \alpha \in (\zeta - 1, \zeta] \quad (7)$$

Definition 7. The “Mittag-Leffler” operator $E_{\alpha}(\wp)$ is

$$E_{\alpha}(\wp) = \sum_{n=0}^{\infty} \frac{\wp^n}{\Gamma(1 + n\alpha)}, \quad (8)$$

where, $\alpha > 0$.

3 | PROCESS OF FUZZY LAPLACE ADOMIAN DECOMPOSITION METHOD

For a rough estimate, we'll look at our conceptual model in this part. An Adomian decomposition ^{19,20} techniques is used in conjunction with the fractional Caputo derivative

$$\begin{aligned} \mathfrak{L}\left[{}^c D_{\wp}^{\alpha} \tilde{\mathfrak{N}}(b, \wp; \delta)\right] &= \mathfrak{L}\left[D_b^2 \tilde{\mathfrak{N}}(b, \wp; \delta) + a \tilde{\mathfrak{N}}(b, \wp; \delta) \right. \\ &\quad \left. + b \tilde{\mathfrak{N}}^2(b, \wp; \delta) + c \tilde{\mathfrak{N}}^3(b, \wp; \delta)\right] + \mathfrak{L}\left[\tilde{k}(\delta) \mathfrak{N}(b, \wp; \delta)\right] \end{aligned} \quad (9)$$

where $\alpha \in (\zeta - 1, \zeta]$; therefore the Laplace Transform of equation (9) is

$$\begin{aligned} p^{\alpha} \mathfrak{L}\left[\tilde{\mathfrak{N}}(b, \wp; \delta)\right] - \sum_{q=0}^{\zeta-1} p^{\alpha-q-1} \tilde{\mathfrak{N}}^q(b, 0; \delta) &= \mathfrak{L}\left[D_b^2 \tilde{\mathfrak{N}}(b, \wp; \delta) + a \tilde{\mathfrak{N}}(b, \wp; \delta) + b \tilde{\mathfrak{N}}^2(b, \wp; \delta) \right. \\ &\quad \left. + c \tilde{\mathfrak{N}}^3(b, \wp; \delta)\right] + \mathfrak{L}\left[\tilde{k}(\delta) \mathfrak{N}(b, \wp; \delta)\right] \end{aligned} \quad (10)$$

$$\begin{aligned} \mathfrak{L}\left[\tilde{\mathfrak{N}}(b, \wp; \delta)\right] &= \left\{ \frac{1}{p^{\alpha}} \sum_{q=0}^{\zeta-1} p^{\alpha-q-1} \tilde{\mathfrak{N}}^q(b, 0; \delta) + \frac{1}{p^{\alpha}} \mathfrak{L}\left[D_b^2 \tilde{\mathfrak{N}}(b, \wp; \delta) \right. \right. \\ &\quad \left. \left. + a \tilde{\mathfrak{N}}(b, \wp; \delta) + b \tilde{\mathfrak{N}}^2(b, \wp; \delta) + c \tilde{\mathfrak{N}}^3(b, \wp; \delta)\right] + \frac{1}{p^{\alpha}} \mathfrak{L}\left[\tilde{k}(\delta) \mathfrak{N}(b, \wp; \delta)\right] \right\} \end{aligned} \quad (11)$$

Apply inverse Laplace Transform to the equation (11) we get,

$$\begin{aligned} \tilde{\mathfrak{N}}(b, \wp; \delta) &= \mathfrak{L}^{-1} \left\{ \frac{1}{p^{\alpha}} \sum_{q=0}^{\zeta-1} p^{\alpha-q-1} \tilde{\mathfrak{N}}^q(b, 0; \delta) + \frac{1}{p^{\alpha}} \mathfrak{L}\left[\tilde{k}(\delta) \mathfrak{N}(b, \wp; \delta)\right] \right\} \\ &\quad + \mathfrak{L}^{-1} \left\{ \frac{1}{p^{\alpha}} \mathfrak{L}\left[D_b^2 \tilde{\mathfrak{N}}(b, \wp; \delta) + a \tilde{\mathfrak{N}}(b, \wp; \delta) + b \tilde{\mathfrak{N}}^2(b, \wp; \delta) + c \tilde{\mathfrak{N}}^3(b, \wp; \delta)\right] \right\}. \end{aligned} \quad (12)$$

We suppose series solution of function $\tilde{\mathfrak{N}}(b, \wp; \delta)$ as

$$\tilde{\mathfrak{N}}(b, \wp; \delta) = \sum_{n=0}^{\infty} \tilde{\mathfrak{N}}_n(b, \wp; \delta) \quad (13)$$

Here, note that $\tilde{\mathfrak{N}}(b, \wp; \delta)$ is a fuzzy number-valued function that can be represented in terms of r-cut representations.

$$\tilde{\mathfrak{N}}(b, \wp; \delta) = \left[\underline{\mathfrak{N}}(b, \wp; \delta), \bar{\mathfrak{N}}(b, \wp; \delta) \right] \quad (14)$$

and the operator L is linear;

$$\therefore L\left(\sum_{n=0}^{\infty} \tilde{\mathfrak{N}}_n(b, \wp; \delta)\right) = \sum_{n=0}^{\infty} L\left[\tilde{\mathfrak{N}}_n(b, \wp; \delta)\right] \quad (15)$$

and the operator N is nonlinear; then equation (12) becomes,

$$\sum_{n=0}^{\infty} \tilde{\mathfrak{N}}_n(b, \wp; \delta) = \mathfrak{Z}^{-1} \left\{ \frac{1}{p^\alpha} \sum_{q=0}^{\zeta-1} p^{\alpha-q-1} \tilde{\mathfrak{N}}^q(b, 0; \delta) + \frac{1}{p^\alpha} \mathfrak{Z} \left[\tilde{k}(\delta) \mathfrak{N}(b, \wp; \delta) \right] \right\} \\ + \mathfrak{Z}^{-1} \left\{ \frac{1}{p^\alpha} \mathfrak{Z} \left[D_b^2 \sum_{n=0}^{\infty} \tilde{\mathfrak{N}}_n(b, \wp; \delta) + a \sum_{n=0}^{\infty} \tilde{\mathfrak{N}}_n(b, \wp; \delta) + b \sum_{n=0}^{\infty} \tilde{\mathfrak{N}}_n^2(b, \wp; \delta) + c \sum_{n=0}^{\infty} \tilde{\mathfrak{N}}_n^3(b, \wp; \delta) \right] \right\}. \quad (16)$$

and the fuzzy nonlinear term $N(\tilde{\mathfrak{N}}) = a\tilde{\mathfrak{N}}_n(b, \wp; \delta) + b\tilde{\mathfrak{N}}_n^2(b, \wp; \delta) + c\tilde{\mathfrak{N}}_n^3(b, \wp; \delta)$ is decomposed as follow:

$$N(\tilde{\mathfrak{N}}) = \sum_{n=0}^{\infty} \underline{A}_n \\ N(\bar{\mathfrak{N}}) = \sum_{n=0}^{\infty} \bar{A}_n \quad (17)$$

where $\tilde{A}_n = [\underline{A}_n, \bar{A}_n]$ is the set polynomials of Adomian . The \tilde{A}_n 's has the formula

$$\underline{A}_n = \frac{1}{n!} \frac{\partial^n}{\partial \mathfrak{N}^n} \left[N \left(\sum_{i=0}^n \mathfrak{N}^i \underline{\mathfrak{N}}_i \right) \right]_{\mathfrak{N}=0}, \quad n = 0, 1, 2, 3, \dots \\ \bar{A}_n = \frac{1}{n!} \frac{\partial^n}{\partial \mathfrak{N}^n} \left[N \left(\sum_{i=0}^n \mathfrak{N}^i \bar{\mathfrak{N}}_i \right) \right]_{\mathfrak{N}=0}, \quad n = 0, 1, 2, 3, \dots \quad (18)$$

some $\tilde{A}_i, (i \rightarrow n)$ terms are as follow:

$$\underline{A}_0 = N(\underline{\mathfrak{N}}_0) \\ \underline{A}_1 = \underline{\mathfrak{N}}_1 (N)'(\underline{\mathfrak{N}}_0) \\ \underline{A}_2 = \frac{1}{2} (N)''(\underline{\mathfrak{N}}_0) \underline{\mathfrak{N}}_1^2 + \underline{\mathfrak{N}}_2 (N)'(\underline{\mathfrak{N}}_0) \\ \underline{A}_3 = \frac{1}{6} (N)'''(\underline{\mathfrak{N}}_0) \underline{\mathfrak{N}}_1^3 + \underline{\mathfrak{N}}_2 (N)''(\underline{\mathfrak{N}}_0) \underline{\mathfrak{N}}_1 + \underline{\mathfrak{N}}_3 (N)'(\underline{\mathfrak{N}}_0) \\ \vdots$$

and

$$\bar{A}_0 = N(\bar{\mathfrak{N}}_0) \\ \bar{A}_1 = \bar{\mathfrak{N}}_1 (N)'(\bar{\mathfrak{N}}_0) \\ \bar{A}_2 = \frac{1}{2} (N)''(\bar{\mathfrak{N}}_0) \bar{\mathfrak{N}}_1^2 + \bar{\mathfrak{N}}_2 (N)'(\bar{\mathfrak{N}}_0) \\ \bar{A}_3 = \frac{1}{6} (N)'''(\bar{\mathfrak{N}}_0) \bar{\mathfrak{N}}_1^3 + \bar{\mathfrak{N}}_2 (N)''(\bar{\mathfrak{N}}_0) \bar{\mathfrak{N}}_1 + \bar{\mathfrak{N}}_3 (N)'(\bar{\mathfrak{N}}_0) \\ \vdots$$

substituting equation (17) and equation (18) into (16) we get, parametric form of $\tilde{\mathfrak{N}}(b, \wp; \delta)$ as follows:

$$\underline{\mathfrak{N}}(b, \wp; \delta) = \sum_{n=0}^{\infty} \underline{\mathfrak{N}}_n(b, \wp; \delta) = \left(\mathfrak{Z}^{-1} \left\{ \frac{1}{p^\alpha} \sum_{q=0}^{\zeta-1} p^{\alpha-q-1} \underline{\mathfrak{N}}^q(b, 0; \delta) + \frac{1}{p^\alpha} \mathfrak{Z} \left[\underline{k}(\delta) \mathfrak{N}(b, \wp; \delta) \right] \right\} \right) \\ + \mathfrak{Z}^{-1} \left\{ \frac{1}{p^\alpha} \mathfrak{Z} \left[D_b^2 \sum_{n=0}^{\infty} \underline{\mathfrak{N}}_n(b, \wp; \delta) + \frac{1}{p^\alpha} \mathfrak{Z} \left[\sum_{n=0}^{\infty} \underline{A}_n \right] \right] \right\} \\ \bar{\mathfrak{N}}(b, \wp; \delta) = \sum_{n=0}^{\infty} \bar{\mathfrak{N}}_n(b, \wp; \delta) = \left(\mathfrak{Z}^{-1} \left\{ \frac{1}{p^\alpha} \sum_{q=0}^{\zeta-1} p^{\alpha-q-1} \bar{\mathfrak{N}}^q(b, 0; \delta) + \frac{1}{p^\alpha} \mathfrak{Z} \left[\bar{k}(\delta) \mathfrak{N}(b, \wp; \delta) \right] \right\} \right) \\ + \mathfrak{Z}^{-1} \left\{ \frac{1}{p^\alpha} \mathfrak{Z} \left[D_b^2 \sum_{n=0}^{\infty} \bar{\mathfrak{N}}_n(b, \wp; \delta) + \frac{1}{p^\alpha} \mathfrak{Z} \left[\sum_{n=0}^{\infty} \bar{A}_n \right] \right] \right\} \quad (19)$$

apply the iterative technique we find the recursive equation in the following form as,

$$\begin{aligned}
\underline{\mathfrak{N}}_0(b, \wp; \delta) &= \mathfrak{Z}^{-1} \left\{ \frac{1}{p^\alpha} \sum_{q=0}^{\zeta-1} p^{\alpha-q-1} \underline{\mathfrak{N}}^q(b, 0; \delta) + \frac{1}{p^\alpha} \mathfrak{Z} \left[\bar{k}(\delta) \underline{\mathfrak{N}}(b, \wp; \delta) \right] \right\} \\
\underline{\mathfrak{N}}_1(b, \wp; \delta) &= \mathfrak{Z}^{-1} \left\{ \frac{1}{p^\alpha} \mathfrak{Z} \left[D_b^2 \underline{\mathfrak{N}}_0(b, \wp; \delta) \right] + \frac{1}{p^\alpha} \mathfrak{Z} \left[\underline{A}_0 \right] \right\} \\
\underline{\mathfrak{N}}_2(b, \wp; \delta) &= \mathfrak{Z}^{-1} \left\{ \frac{1}{p^\alpha} \mathfrak{Z} \left[D_b^2 \underline{\mathfrak{N}}_1(b, \wp; \delta) \right] + \frac{1}{p^\alpha} \mathfrak{Z} \left[\underline{A}_1 \right] \right\} \\
&\vdots \\
\underline{\mathfrak{N}}_{n+1}(b, \wp; \delta) &= \mathfrak{Z}^{-1} \left\{ \frac{1}{p^\alpha} \mathfrak{Z} \left[D_b^2 \underline{\mathfrak{N}}_n(b, \wp; \delta) \right] + \frac{1}{p^\alpha} \mathfrak{Z} \left[\underline{A}_n \right] \right\}
\end{aligned} \tag{20}$$

$$\begin{aligned}
\overline{\mathfrak{N}}_0(b, \wp; \delta) &= \mathfrak{Z}^{-1} \left\{ \frac{1}{p^\alpha} \sum_{q=0}^{\zeta-1} p^{\alpha-q-1} \overline{\mathfrak{N}}^q(b, 0; \delta) + \frac{1}{p^\alpha} \mathfrak{Z} \left[\bar{k}(\delta) \overline{\mathfrak{N}}(b, \wp; \delta) \right] \right\} \\
\overline{\mathfrak{N}}_1(b, \wp; \delta) &= \mathfrak{Z}^{-1} \left\{ \frac{1}{p^\alpha} \mathfrak{Z} \left[D_b^2 \overline{\mathfrak{N}}_0(b, \wp; \delta) \right] + \frac{1}{p^\alpha} \mathfrak{Z} \left[\overline{A}_0 \right] \right\} \\
\overline{\mathfrak{N}}_2(b, \wp; \delta) &= \mathfrak{Z}^{-1} \left\{ \frac{1}{p^\alpha} \mathfrak{Z} \left[D_b^2 \overline{\mathfrak{N}}_1(b, \wp; \delta) \right] + \frac{1}{p^\alpha} \mathfrak{Z} \left[\overline{A}_1 \right] \right\} \\
&\vdots \\
\overline{\mathfrak{N}}_{n+1}(b, \wp; \delta) &= \mathfrak{Z}^{-1} \left\{ \frac{1}{p^\alpha} \mathfrak{Z} \left[D_b^2 \overline{\mathfrak{N}}_n(b, \wp; \delta) \right] + \frac{1}{p^\alpha} \mathfrak{Z} \left[\overline{A}_n \right] \right\}
\end{aligned} \tag{21}$$

Thus, the solution becomes

$$\underline{\mathfrak{N}}(b, \wp; \delta) = \underline{\mathfrak{N}}_0(b, \wp; \delta) + \underline{\mathfrak{N}}_1(b, \wp; \delta) + \dots \tag{22}$$

$$\overline{\mathfrak{N}}(b, \wp; \delta) = \overline{\mathfrak{N}}_0(b, \wp; \delta) + \overline{\mathfrak{N}}_1(b, \wp; \delta) + \dots \tag{23}$$

The Solution of the given series is convergent rapidly and the classical approach to converge to this series has been represented by Y. Cherruault and G. Adomian[3]

4 | NUMERICAL EXAMPLES

Example 1. We consider the fuzzy fractional Klein-Gordan equation as

$$\frac{\partial^\alpha \tilde{\mathfrak{N}}(b, \wp; \delta)}{\partial \wp^\alpha} - \frac{\partial^2 \tilde{\mathfrak{N}}(b, \wp; \delta)}{\partial b^2} + \tilde{\mathfrak{N}}(b, \wp; \delta) = 0, \quad 1 < \alpha \leq 2, \tag{24}$$

With the fuzzy initial conditions

$$\tilde{\mathfrak{N}}(b, 0; \delta) = 0 \quad \text{and} \quad \tilde{\mathfrak{N}}_\wp(b, 0; \delta) = \tilde{k}(\delta) \odot b \tag{25}$$

where $\tilde{k}(\delta) = [\delta - 1, 1 - \delta]$

Using the scheme of equation (20) and (21), we obtain

$$\begin{aligned}
 \underline{\mathfrak{N}}_0(b, \wp; \delta) &= b(\delta - 1)\wp \\
 \overline{\mathfrak{N}}_0(b, \wp; \delta) &= b(1 - \delta)\wp \\
 \underline{\mathfrak{N}}_1(b, \wp; \delta) &= -b(\delta - 1)\frac{\wp^{\alpha+1}}{\Gamma(\alpha + 2)} \\
 \overline{\mathfrak{N}}_1(b, \wp; \delta) &= -b(1 - \delta)\frac{\wp^{\alpha+1}}{\Gamma(\alpha + 2)} \\
 \underline{\mathfrak{N}}_2(b, \wp; \delta) &= b(\delta - 1)\frac{\wp^{2\alpha+1}}{\Gamma(2\alpha + 2)} \\
 \overline{\mathfrak{N}}_2(b, \wp; \delta) &= b(1 - \delta)\frac{\wp^{2\alpha+1}}{\Gamma(2\alpha + 2)} \\
 \underline{\mathfrak{N}}_3(b, \wp; \delta) &= -b(\delta - 1)\frac{\wp^{3\alpha+1}}{\Gamma(3\alpha + 2)} \\
 \overline{\mathfrak{N}}_3(b, \wp; \delta) &= -b(1 - \delta)\frac{\wp^{3\alpha+1}}{\Gamma(3\alpha + 2)} \\
 \underline{\mathfrak{N}}_4(b, \wp; \delta) &= b(\delta - 1)\frac{\wp^{4\alpha+1}}{\Gamma(4\alpha + 2)} \\
 \overline{\mathfrak{N}}_4(b, \wp; \delta) &= b(1 - \delta)\frac{\wp^{4\alpha+1}}{\Gamma(4\alpha + 2)} \\
 \underline{\mathfrak{N}}_5(b, \wp; \delta) &= -b(\delta - 1)\frac{\wp^{5\alpha+1}}{\Gamma(5\alpha + 2)} \\
 \overline{\mathfrak{N}}_4(b, \wp; \delta) &= -b(1 - \delta)\frac{\wp^{5\alpha+1}}{\Gamma(5\alpha + 2)} \\
 &\vdots
 \end{aligned}$$

and so on. Thus the lower and upper forms of a solution are given as

$$\underline{\mathfrak{N}}(b, \wp; \delta) = \underline{\mathfrak{N}}_0(b, \wp; \delta) + \underline{\mathfrak{N}}_1(b, \wp; \delta) + \underline{\mathfrak{N}}_2(b, \wp; \delta) + \underline{\mathfrak{N}}_3(b, \wp; \delta) + \dots$$

$$\overline{\mathfrak{N}}(b, \wp; \delta) = \overline{\mathfrak{N}}_0(b, \wp; \delta) + \overline{\mathfrak{N}}_1(b, \wp; \delta) + \overline{\mathfrak{N}}_2(b, \wp; \delta) + \overline{\mathfrak{N}}_3(b, \wp; \delta) + \dots$$

$$\begin{aligned}
 \underline{\mathfrak{N}}(b, \wp; \delta) &= b(\delta - 1)\wp \left[1 - \frac{\wp^\alpha}{\Gamma(\alpha+2)} + \frac{\wp^{2\alpha}}{\Gamma(2\alpha+2)} - \frac{\wp^{3\alpha}}{\Gamma(3\alpha+2)} \right. \\
 &\quad \left. + \frac{\wp^{4\alpha}}{\Gamma(4\alpha+2)} - \frac{\wp^{5\alpha}}{\Gamma(5\alpha+2)} + \dots \right] \\
 \overline{\mathfrak{N}}(b, \wp; \delta) &= b(1 - \delta)\wp \left[1 - \frac{\wp^\alpha}{\Gamma(\alpha+2)} + \frac{\wp^{2\alpha}}{\Gamma(2\alpha+2)} - \frac{\wp^{3\alpha}}{\Gamma(3\alpha+2)} \right. \\
 &\quad \left. + \frac{\wp^{4\alpha}}{\Gamma(4\alpha+2)} - \frac{\wp^{5\alpha}}{\Gamma(5\alpha+2)} + \dots \right]
 \end{aligned} \tag{26}$$




Figure 1 : 3-Dimensional simulation of Example (1) at $\alpha = 0.8, 1.5$ and uncertainty $\delta \in [0, 1]$




Figure 2 : 2-Dimensional simulation of Example (1) at $\alpha = 0.8, 1.5, 2$ and uncertainty $\delta \in [0, 1]$

If we put $\alpha = 2$ in equation (26) then we have

$$\tilde{\mathfrak{N}}(\mathfrak{b}, \wp; \delta) = \tilde{k}(\delta) \mathfrak{b} \sin(\wp)$$

which is exact solution.




Figure 3 : Exact Geometrical representation of equation (26) at $\alpha = 2$

$$\alpha = 2$$

r	Lower			Upper		
	Approximate	Exact	Error	Approximate	Exact	Error
0	-1.81859	-1.81859	2.58173×10^{-6}	1.81859	1.81859	2.58173×10^{-6}
0.2	-1.45487	-1.45488	2.06538×10^{-6}	1.45487	1.45488	2.06538×10^{-6}
0.4	-1.09116	-1.09116	1.54904×10^{-6}	1.09116	1.09116	1.54904×10^{-6}
0.6	-0.727437	-0.727438	1.03269×10^{-6}	7.27437×10^{-1}	7.27438×10^{-1}	1.03269×10^{-6}
0.8	-0.363718	-0.363719	5.16345×10^{-7}	3.63718×10^{-1}	3.63719×10^{-1}	5.16345×10^{-7}
1	0	0	0.00000	0.00000	0.00000	0.00000

Example 2. We consider the non-homogeneous fuzzy fractional Klein-Gordon equation as

$$\frac{\partial^\alpha \tilde{\mathfrak{N}}(b, \wp; \delta)}{\partial \wp^\alpha} - \frac{\partial^2 \tilde{\mathfrak{N}}(b, \wp; \delta)}{\partial b^2} + \tilde{\mathfrak{N}}(b, \wp; \delta) = 2 \sin(b), \quad 1 < \alpha \leq 2, \quad (27)$$

With the fuzzy initial conditions

$$\tilde{\mathfrak{N}}(b, 0; \delta) = \tilde{k}(\delta) \sin(b) \quad \text{and} \quad \tilde{\mathfrak{N}}_\wp(b, 0; \delta) = \tilde{k}(\delta) \quad (28)$$

where $\tilde{k}(\delta) = [\delta - 1, 1 - \delta]$

Using the scheme of equation (20) and (21), we obtain

$$\begin{aligned} \underline{\mathfrak{N}}_0(b, \wp; \delta) &= \frac{2 \sin(b) \wp^\alpha}{\Gamma(\alpha + 1)} - \sin(b) + \sin(b) \delta + \wp \delta - \wp \\ \overline{\mathfrak{N}}_0(b, \wp; \delta) &= \frac{2 \sin(b) \wp^\alpha}{\Gamma(\alpha + 1)} + \sin(b) - \sin(b) \delta - \wp \delta + \wp \\ \underline{\mathfrak{N}}_1(b, \wp; \delta) &= -\frac{4 \sin(b) \wp^{2\alpha}}{\Gamma(2\alpha + 1)} + \frac{2 \wp^\alpha \sin(b)}{\Gamma(\alpha + 1)} + \frac{\wp^{\alpha+1}}{\Gamma(\alpha + 2)} - \frac{2 \sin(b) \delta \wp^\alpha}{\Gamma(\alpha + 1)} - \frac{\delta \wp^{\alpha+1}}{\Gamma(\alpha + 2)} \\ \overline{\mathfrak{N}}_1(b, \wp; \delta) &= -\frac{4 \sin(b) \wp^{2\alpha}}{\Gamma(2\alpha + 1)} - \frac{2 \wp^\alpha \sin(b)}{\Gamma(\alpha + 1)} - \frac{\wp^{\alpha+1}}{\Gamma(\alpha + 2)} + \frac{2 \sin(b) \delta \wp^\alpha}{\Gamma(\alpha + 1)} + \frac{\delta \wp^{\alpha+1}}{\Gamma(\alpha + 2)} \\ \underline{\mathfrak{N}}_2(b, \wp; \delta) &= -\frac{4 \wp^{2\alpha} \sin(b)}{\Gamma(2\alpha + 1)} + \frac{8 \wp^{3\alpha} \sin(b)}{\Gamma(3\alpha + 1)} - \frac{\wp^{2\alpha+1}}{\Gamma(2\alpha + 2)} + \frac{4 \delta \wp^{2\alpha} \sin(b)}{\Gamma(2\alpha + 1)} + \frac{\delta \wp^{2\alpha+1}}{\Gamma(2\alpha + 2)} \\ \overline{\mathfrak{N}}_2(b, \wp; \delta) &= \frac{4 \wp^{2\alpha} \sin(b)}{\Gamma(2\alpha + 1)} + \frac{8 \wp^{3\alpha} \sin(b)}{\Gamma(3\alpha + 1)} + \frac{\wp^{2\alpha+1}}{\Gamma(2\alpha + 2)} - \frac{4 \delta \wp^{2\alpha} \sin(b)}{\Gamma(2\alpha + 1)} - \frac{\delta \wp^{2\alpha+1}}{\Gamma(2\alpha + 2)} \\ \underline{\mathfrak{N}}_3(b, \wp; \delta) &= \frac{8 \wp^{3\alpha} \sin(b)}{\Gamma(3\alpha + 1)} - \frac{16 \wp^{4\alpha} \sin(b)}{\Gamma(4\alpha + 1)} + \frac{\wp^{3\alpha+1}}{\Gamma(3\alpha + 2)} - \frac{8 \delta \wp^{3\alpha} \sin(b)}{\Gamma(3\alpha + 1)} - \frac{\delta \wp^{3\alpha+1}}{\Gamma(3\alpha + 2)} \\ \overline{\mathfrak{N}}_3(b, \wp; \delta) &= -\frac{8 \wp^{3\alpha} \sin(b)}{\Gamma(3\alpha + 1)} + \frac{16 \wp^{4\alpha} \sin(b)}{\Gamma(4\alpha + 1)} - \frac{\wp^{3\alpha+1}}{\Gamma(3\alpha + 2)} + \frac{8 \delta \wp^{3\alpha} \sin(b)}{\Gamma(3\alpha + 1)} + \frac{\delta \wp^{3\alpha+1}}{\Gamma(3\alpha + 2)} \\ \underline{\mathfrak{N}}_4(b, \wp; \delta) &= \left[-\frac{4 \wp^{2\alpha} \sin(b)}{\Gamma(2\alpha + 1)} + \frac{8 \wp^{3\alpha} \sin(b)}{\Gamma(3\alpha + 1)} - \frac{32 \wp^{4\alpha} \sin(b)}{\Gamma(4\alpha + 1)} + \frac{2 \wp^\alpha \sin(b)}{\Gamma(\alpha + 1)} - \frac{\wp^{4\alpha+1}}{\Gamma(4\alpha + 2)} \right. \\ &\quad \left. + \frac{16 \delta \wp^{4\alpha} \sin(b)}{\Gamma(4\alpha + 1)} + \frac{\delta \wp^{4\alpha+1}}{\Gamma(4\alpha + 2)} \right] \\ \overline{\mathfrak{N}}_4(b, \wp; \delta) &= \left[-\frac{4 \wp^{2\alpha} \sin(b)}{\Gamma(2\alpha + 1)} + \frac{8 \wp^{3\alpha} \sin(b)}{\Gamma(3\alpha + 1)} - \frac{32 \wp^{4\alpha} \sin(b)}{\Gamma(4\alpha + 1)} + \frac{2 \wp^\alpha \sin(b)}{\Gamma(\alpha + 1)} - \frac{\wp^{4\alpha+1}}{\Gamma(4\alpha + 2)} \right. \\ &\quad \left. + \frac{16 \delta \wp^{4\alpha} \sin(b)}{\Gamma(4\alpha + 1)} + \frac{\delta \wp^{4\alpha+1}}{\Gamma(4\alpha + 2)} \right] \\ &\vdots \end{aligned}$$

and so on. Thus the lower and upper forms of a solution are given as

$$\begin{aligned} \underline{\mathfrak{N}}(b, \wp; \delta) &= \underline{\mathfrak{N}}_0(b, \wp; \delta) + \underline{\mathfrak{N}}_1(b, \wp; \delta) + \underline{\mathfrak{N}}_2(b, \wp; \delta) + \underline{\mathfrak{N}}_3(b, \wp; \delta) + \dots \\ \overline{\mathfrak{N}}(b, \wp; \delta) &= \overline{\mathfrak{N}}_0(b, \wp; \delta) + \overline{\mathfrak{N}}_1(b, \wp; \delta) + \overline{\mathfrak{N}}_2(b, \wp; \delta) + \overline{\mathfrak{N}}_3(b, \wp; \delta) + \dots \end{aligned}$$

$$\begin{aligned}
\underline{\mathfrak{N}}(b, \wp; \delta) &= \left[\begin{aligned} &-\frac{12\wp^{2\alpha}\sin(b)}{\Gamma(2\alpha+1)} + \frac{24\wp^{3\alpha}\sin(b)}{\Gamma(3\alpha+1)} - \frac{48\wp^{4\alpha}\sin(b)}{\Gamma(4\alpha+1)} + \frac{6\wp^{\alpha}\sin(b)}{\Gamma(\alpha+1)} + \frac{\wp^{\alpha+1}}{\Gamma(\alpha+2)} - \frac{\wp^{2\alpha+1}}{\Gamma(2\alpha+2)} \\ &+ \frac{\wp^{3\alpha+1}}{\Gamma(3\alpha+2)} - \frac{\wp^{4\alpha+1}}{\Gamma(4\alpha+2)} - \sin(b) + \frac{4r\wp^{2\alpha}\sin(b)}{\Gamma(2\alpha+1)} - \frac{8r\wp^{3\alpha}\sin(b)}{\Gamma(3\alpha+1)} + \frac{16r\wp^{4\alpha}\sin(b)}{\Gamma(4\alpha+1)} \\ &- \frac{2r\wp^{\alpha}\sin(b)}{\Gamma(\alpha+1)} - \frac{r\wp^{\alpha+1}}{\Gamma(\alpha+2)} + \frac{r\wp^{2\alpha+1}}{\Gamma(2\alpha+2)} - \frac{r\wp^{3\alpha+1}}{\Gamma(3\alpha+2)} + \frac{r\wp^{4\alpha+1}}{\Gamma(4\alpha+2)} + r\sin(b) + r\wp - \wp \end{aligned} \right] \\
\bar{\mathfrak{N}}(b, \wp; \delta) &= \left[\begin{aligned} &-\frac{4\wp^{2\alpha}\sin(b)}{\Gamma(2\alpha+1)} + \frac{8\wp^{3\alpha}\sin(b)}{\Gamma(3\alpha+1)} - \frac{48\wp^{4\alpha}\sin(b)}{\Gamma(4\alpha+1)} + \frac{2\wp^{\alpha}\sin(b)}{\Gamma(\alpha+1)} - \frac{\wp^{\alpha+1}}{\Gamma(\alpha+2)} + \frac{\wp^{2\alpha+1}}{\Gamma(2\alpha+2)} \\ &+ \frac{\wp^{3\alpha+1}}{\Gamma(3\alpha+2)} - \frac{\wp^{4\alpha+1}}{\Gamma(4\alpha+2)} + \sin(b) - \frac{4r\wp^{2\alpha}\sin(b)}{\Gamma(2\alpha+1)} + \frac{8r\wp^{3\alpha}\sin(b)}{\Gamma(3\alpha+1)} - \frac{16r\wp^{4\alpha}\sin(b)}{\Gamma(4\alpha+1)} \\ &+ \frac{2r\wp^{\alpha}\sin(b)}{\Gamma(\alpha+1)} + \frac{r\wp^{\alpha+1}}{\Gamma(\alpha+2)} - \frac{r\wp^{2\alpha+1}}{\Gamma(2\alpha+2)} + \frac{r\wp^{3\alpha+1}}{\Gamma(3\alpha+2)} - \frac{r\wp^{4\alpha+1}}{\Gamma(4\alpha+2)} - r\sin(b) - r\wp + \wp \end{aligned} \right]
\end{aligned} \tag{29}$$




Figure 4 : 3-Dimensional simulation of (2) at $\alpha = 0.8, 1.5$ and uncertainty $\delta \in [0, 1]$




Figure 5 : 2-Dimensional simulation of (2) at $\alpha = 0.8, 1.5, 2$ and uncertainty $\delta \in [0, 1]$

If we put $\alpha = 2$ in equation (29) then we have

$$\tilde{\mathfrak{N}}(b, \wp; \delta) = \tilde{k}(\delta) \left(\sin(b) + \sin(\wp) \right)$$

which is exact solution.




Figure 6 : Exact Geometrical representation of equation (26) at $\alpha = 2$

$\alpha = 2$						
		Lower		Upper		
r	Approximate	Exact	Error	Approimate	Exact	Error
0	3.4699×10^{-1}	-1.81859	2.16559	4.5101×10^{-1}	1.81859	1.36758
0.2	3.57392×10^{-1}	-1.45488	1.81227	4.40608×10^{-1}	1.45488	1.01427
0.4	3.67794×10^{-1}	-1.09116	1.45895	4.30206×10^{-1}	1.09116	6.60951×10^{-1}
0.6	3.78196×10^{-1}	-0.727438	1.10563	4.19804×10^{-1}	7.27438×10^{-1}	3.07634×10^{-1}
0.8	3.88598×10^{-1}	-0.363719	7.52317×10^{-1}	4.09402×10^{-1}	3.63719×10^{-1}	4.56834×10^{-2}
1	3.99×10^{-1}	0	3.99×10^{-1}	3.99×10^{-1}	0	3.99×10^{-1}

Example 3. We consider the fuzzy fractional Klein-Gordan equation as

$$\frac{\partial^\alpha \tilde{\mathfrak{N}}(b, \wp; \delta)}{\partial \wp^\alpha} - \frac{\partial^2 \tilde{\mathfrak{N}}(b, \wp; \delta)}{\partial b^2} - \left(\frac{\partial \tilde{\mathfrak{N}}(b, \wp; \delta)}{\partial b} \right)^2 - \tilde{\mathfrak{N}}^2(b, \wp; \delta) = 0, \quad 0 < \alpha \leq 1, \quad (30)$$

With the fuzzy initial condition

$$\tilde{\mathfrak{N}}(b, 0; \delta) = 0 \quad \text{and} \quad \tilde{\mathfrak{N}}_\wp(b, 0; \delta) = \tilde{k}(\delta) e^b \quad (31)$$

where $\tilde{k}(\delta) = [\delta - 1, 1 - \delta]$

Using the scheme of equation(18), (20) and (21), we get

$$\begin{aligned}\underline{\mathfrak{N}}_0(b, \wp; \delta) &= \wp(\delta - 1)e^b \\ \overline{\mathfrak{N}}_0(b, \wp; \delta) &= \wp(1 - \delta)e^b \\ \underline{\mathfrak{N}}_1(b, \wp; \delta) &= e^b(\delta - 1)\frac{\wp^{\alpha+1}}{\Gamma(\alpha + 2)} \\ \overline{\mathfrak{N}}_1(b, \wp; \delta) &= e^b(1 - \delta)\frac{\wp^{\alpha+1}}{\Gamma(\alpha + 2)} \\ \underline{\mathfrak{N}}_2(b, \wp; \delta) &= e^b(\delta - 1)\frac{\wp^{2\alpha+1}}{\Gamma(2\alpha + 2)} \\ \overline{\mathfrak{N}}_2(b, \wp; \delta) &= e^b(1 - \delta)\frac{\wp^{2\alpha+1}}{\Gamma(2\alpha + 2)} \\ \underline{\mathfrak{N}}_3(b, \wp; \delta) &= e^b(\delta - 1)\frac{\wp^{3\alpha+1}}{\Gamma(3\alpha + 2)} \\ \overline{\mathfrak{N}}_3(b, \wp; \delta) &= e^b(1 - \delta)\frac{\wp^{3\alpha+1}}{\Gamma(3\alpha + 2)} \\ &\vdots\end{aligned}$$

and so on. Thus the lower and upper forms of a solution are given as

$$\begin{aligned}\underline{\mathfrak{N}}(b, \wp; \delta) &= e^b(\delta - 1)\wp \left(1 + \frac{\wp^\alpha}{\Gamma(\alpha + 2)} + \frac{\wp^{2\alpha}}{\Gamma(2\alpha + 2)} + \frac{\wp^{3\alpha}}{\Gamma(3\alpha + 2)} + \frac{\wp^{4\alpha}}{\Gamma(4\alpha + 2)} + \dots \right) \\ \overline{\mathfrak{N}}(b, \wp; \delta) &= e^b(1 - \delta)\wp \left(1 + \frac{\wp^\alpha}{\Gamma(\alpha + 2)} + \frac{\wp^{2\alpha}}{\Gamma(2\alpha + 2)} + \frac{\wp^{3\alpha}}{\Gamma(3\alpha + 2)} + \frac{\wp^{4\alpha}}{\Gamma(4\alpha + 2)} + \dots \right)\end{aligned}\quad (32)$$

If we put $\alpha = 2$ in equation (32) then we have

$$\tilde{\mathfrak{N}}(b, \wp; \delta) = \tilde{k}(\delta) \left(e^b \sinh(\wp) \right)$$

which is exact solution.

$\alpha = 2$						
Lower			Upper			
r	Approximate	Exact	Error	Approimate	Exact	Error
0	-26.7987	-26.7991	3.89017×10^{-4}	2.67987×10^1	2.67991×10^1	3.89017×10^{-4}
0.2	-21.4389	-21.4393	3.11213×10^{-4}	2.14389×10^1	2.14393×10^1	3.11213×10^{-4}
0.4	-16.0792	-16.0794	2.3341×10^{-4}	1.60792×10^1	1.60794×10^1	2.3341×10^{-4}
0.6	-10.7195	-10.7196	1.55607×10^{-4}	1.07195×10^1	1.07196×10^1	1.55607×10^{-4}
0.8	-5.35974	-5.35982	7.78034×10^{-5}	5.35974	5.35982	7.78034×10^{-5}
1	0	0	0	0	0	0




Figure 7 : 3-Dimensional simulation of Example (3) at $\alpha = 0.6, 0.8$ and uncertainty $r \in [0, 1]$




Figure 8 : 2-Dimensional simulation of Example (3) at $\alpha = 0.6, 0.8, 1$ and uncertainty $r \in [0, 1]$




Figure 9 : Exact Geometrical representation of equation (26) at $\alpha = 2$

CONCLUSION

An approximation to the solution of the fuzzy time fractional telegraph equations was the primary focus of this article's efforts. The authors were successful in accomplishing their mission thanks to the application of the Fuzzy Laplace-Adomian decomposition approach. The Adomian decomposition method is a powerful method that has presented a possibility for effective efficiency in the solution of nonlinear differential equations for physical applications. The authors conducted their computations with the help of the Wolfram Mathematica 11.3 programme.

ACKNOWLEDGMENTS

I'm very grateful to our co-authors and mentors, Professor Vasant R. Nikam, Professor S.B.Gaikwad, and Professor Shivaji Tarate, among others. They gave me a lot of help when I was writing this research article.

Author contributions

Shivaji Tarate was helping in Typography.

Financial disclosure

There is no funding of any agency for this article.

Conflict of interest

The authors declare no potential conflict of interests.

References

1. Abbod MF, Keyserlingk vDG, Linkens DA, Mahfouf M. Survey of utilisation of fuzzy technology in medicine and healthcare. *Fuzzy Sets and Systems* 2001; 120(2): 331–349.
2. Datta DP. The golden mean, scale free extension of real number system, fuzzy sets and 1/f spectrum in physics and biology. *Chaos, Solitons & Fractals* 2003; 17(4): 781–788.
3. El Naschie M. From experimental quantum optics to quantum gravity via a fuzzy Kähler manifold. *Chaos, Solitons & Fractals* 2005; 25(5): 969–977.
4. El Naschie M. On a fuzzy Kähler-like manifold which is consistent with the two slit experiment. *International Journal of Nonlinear Sciences and Numerical Simulation* 2005; 6(2): 95–98.
5. Zhang H, Liao X, Yu J. Fuzzy modeling and synchronization of hyperchaotic systems. *Chaos, Solitons & Fractals* 2005; 26(3): 835–843.
6. Fei W. Existence and uniqueness of solution for fuzzy random differential equations with non-Lipschitz coefficients. *Information Sciences* 2007; 177(20): 4329–4337.
7. Kaleva O. Fuzzy differential equations. *Fuzzy sets and systems* 1987; 24(3): 301–317.
8. Kaleva O. The Cauchy problem for fuzzy differential equations. *Fuzzy sets and systems* 1990; 35(3): 389–396.
9. Rodríguez-López R. Comparison results for fuzzy differential equations. *Information Sciences* 2008; 178(6): 1756–1779.
10. Ma M, Friedman M, Kandel A. Numerical solutions of fuzzy differential equations. *Fuzzy sets and systems* 1999; 105(1): 133–138.

11. Mizukoshi MT, Barros LD, Chalco-Cano Y, Román-Flores H, Bassanezi RC. Fuzzy differential equations and the extension principle. *Information Sciences* 2007; 177(17): 3627–3635.
12. Chang SS, Zadeh LA. On fuzzy mapping and control. In: World Scientific. 1996 (pp. 180–184).
13. Dubois D, Prade H. Towards fuzzy differential calculus part 3: Differentiation. *Fuzzy sets and systems* 1982; 8(3): 225–233.
14. Abbasbandy S. Application of He's homotopy perturbation method for Laplace transform. *Chaos, Solitons & Fractals* 2006; 30(5): 1206–1212.
15. Yusufoglu E. Numerical solution of Duffing equation by the Laplace decomposition algorithm. *Applied Mathematics and Computation* 2006; 177(2): 572–580.
16. Babolian E, Biazar J, Vahidi A. A new computational method for Laplace transforms by decomposition method. *Applied Mathematics and Computation* 2004; 150(3): 841–846.
17. Kiymaz O. An algorithm for solving initial value problems using Laplace Adomian decomposition method. 2009.
18. Khuri SA. A Laplace decomposition algorithm applied to a class of nonlinear differential equations. *Journal of applied mathematics* 2001; 1(4): 141–155.
19. Adomian G. A review of the decomposition method and some recent results for nonlinear equations. *Mathematical and Computer Modelling* 1990; 13(7): 17–43.
20. Adomian G. *Solving frontier problems of physics: the decomposition method*. 60. Springer Science & Business Media . 2013.
21. Nourazar SS, Soori M, Nazari-Golshan A, Nourazar SS, Soori M, Nazari-Golshan A. On the Exact Solution of Burgers-Huxley Equation Using the Homotopy Perturbation Method. *Journal of Applied Mathematics and Physics* 2015; 3(3): 285–294. doi: 10.4236/JAMP.2015.33042
22. Shang D. Exact solutions of coupled nonlinear Klein-Gordon equation. *Applied Mathematics and Computation* 2010; 217(4): 1577–1583. doi: 10.1016/J.AMC.2009.06.072
23. Allahviranloo T, Taheri N. An analytic approximation to the solution of fuzzy heat equation by Adomian decomposition method. *Int. J. Contemp. Math. Sci* 2009; 4(3): 105–114.
24. Alikhani R, Bahrami F. Fuzzy partial differential equations under the cross product of fuzzy numbers. *Information Sciences* 2019; 494: 80–99. doi: 10.1016/j.ins.2019.04.030
25. Jameel A, Anakira N, Alomari A, Hashim I, Momani S. A new approximation method for solving fuzzy heat equations. *Journal of Computational and Theoretical Nanoscience* 2016; 13(11): 7825–7832.
26. Stepnicka M, Valasek R. Numerical solution of partial differential equations with help of fuzzy transform. In: IEEE. ; 2005: 1104–1109.
27. Jameel A. Semi-analytical solution of heat equation in fuzzy environment. *International Journal of Applied Physics and Mathematics* 2014; 4(6): 371–378.
28. Chakraverty S, Tapaswini S, Behera D. *Fuzzy arbitrary order system: fuzzy fractional differential equations and applications*. John Wiley & Sons . 2016.
29. Kochubei A, Luchko Y. *Fractional differential equations*. New York: Mathematics in Science and Engineering . 2019
30. Mazandarani M, Kamyad AV. Modified fractional Euler method for solving fuzzy fractional initial value problem. *Communications in Nonlinear Science and Numerical Simulation* 2013; 18(1): 12–21.
31. Allahviranloo T. Fuzzy fractional differential equations. In: Springer. 2021 (pp. 127–192).
32. Allahviranloo T, Ahmadi MB. Fuzzy laplace transforms. *Soft Computing* 2010; 14(3): 235–243.

AUTHOR BIOGRAPHY

	Kishor A. Kshirsagar Working as Assistance Prof. and research fellow in New Arts,Commerce and Science college Ahmednagar,Maharastra,India.
shiv.pdf	Prof. Vasant R. Nikam Working as Professor and Reaserch Guide in Loknete Vyankatrao Hiray Arts, Science and Commerce College, Nashik,Maharastra,India.
ap.pdf	Prof. S.B.Gaikwad Working as Professor and Reaserch Guide in New Arts,Commerce and Science college Ahmednagar,Maharastra,India.
sbk.pdf	Shivaji A. Tarate Working as Assistance Prof. and research fellow in New Arts,Commerce and Science college Ahmednagar,Maharastra,India.
kk.pdf	How to cite this article: Kishor K., S. Tarate, V. Nikam, and S. Gaikwad (2022), Fuzzy Laplace-Adomian Decomposition method for solving Fuzzy Klein-Gordan equations, <i>Q.J.R. Meteorol. Soc.</i> , 2017;00:1–6.