

The Combined and Individual Effects of the North Atlantic Oscillation and the Atlantic Meridional Mode on Early Rainfall Season Precipitation in the Insular Caribbean

Flavia D. S. Moraes*, Gabriel Kooperman, and Thomas L. Mote

Climate Research Laboratory, Department of Geography, University of Georgia, Athens, GA, USA

Introduction

The insular Caribbean is a region influenced by North Atlantic climate variability. Effects of low-frequency atmospheric circulation patterns on the precipitation of the Caribbean have been well documented. However, individual modes of variability are usually only considered in isolation. Here we **analyse the combined and individual effects of the North Atlantic Oscillation (NAO) and the Atlantic Meridional Mode (AMM) on insular Caribbean precipitation**. This work focuses on the Early Rainfall Season (ERS, April-July), which explains much of the interannual variability in precipitation for this region, from 1960-2016.

Data and Methods

- Used Climate Research Unit (CRU, V4.1) precipitation dataset to calculate monthly precipitation anomaly (%), from 1960-2016.
- Correlated monthly precipitation anomaly (%) from each country (19) with monthly NAO and AMM indices (NOAA Earth System Research Laboratory).
- Analyzed which months presented more spatially distributed and statistically significant correlation (p -value < 0.05) between NAO/AMM and precipitation anomalies (%) (**Figure 1**).




Figure 1. Correlation between NAO/AMM and the insular Caribbean precipitation anomalies. Each relationship is represented by one circle sector and colored in red (positive) or blue (negative).

- Created sea surface temperature (SST) and sea level pressure (SLP) anomalies composites based on the same statistically significant correlated months (NOAA Monthly Climate Composites).

References

Charlery, J., Nurse, L., & Whitehall, K. (2006). Exploring the relationship between the North Atlantic Oscillation and rainfall patterns in Barbados. *International Journal of Climatology*, 26, 819–827.
George, S. E., & Saunders, M. A. (2001). North Atlantic Oscillation impact on tropical North Atlantic winter atmospheric variability. *Geophysical Research Letters*, 28, 1015–1018.
Jury, M., Malmgren, B. A., & Winter, A. (2007). Subregional precipitation climate of the Caribbean and relationships with ENSO and NAO. *Journal of Geophysical Research: Atmospheres*, 112, D16107.
Malmgren, B. A., Winter, A., & Chen, D. (1998). El Niño–Southern Oscillation and North Atlantic Oscillation control of climate in Puerto Rico. *Journal of Climate*, 11, 2713–2717.

Results

The NAO and AMM (Feb-Mar) presented a **correlation of opposite signs** and affected the Eastern Caribbean during the ERS months of May-Jun, resulting in precipitation anomalies of at least $\pm 10\%$. For both **positive and negative precipitation anomalies distribution**, the more frequent (19 years) and consistent regional effects were registered over the **Eastern Caribbean** when **NAO and AMM occurred together** in the previous winter (**Figure 2,a**). However, greater values of precipitation anomalies(+) were related to wintertime NAO(-) alone, while lower values of precipitation anomalies(-) were related to wintertime AMM(-) alone (**Figure 2,b**).




Figure 2. Distribution of precipitation anomalies (a) and precipitation anomalies extremes (b) in the Eastern Caribbean (May-Jun) when NAO and AMM occurred together or isolated (Feb-Mar).

The consistent regional effect on precipitation anomalies during ERS related to the combination of wintertime NAO/AMM was evident during the **severe drought of 2015 (Figure 3)**, when the Eastern Caribbean registered between -15% and -30% of annual precipitation anomaly compared to 1979-2014 (Mote et al. 2017).




Figure 3. Reservoir in Toa Alta (right) and dry banks of La Plata River in Comerio (left), Puerto Rico.

SST and SLP anomalies composites indicated that when wintertime NAO and AMM are combined, both SST and SLP anomalies persist over the Eastern Caribbean during the months of May-Jun (**Figure 4**). The persistent anomalies in SST and SLP do not happen over the entire Eastern Caribbean when NAO or AMM occurred alone in the previous winter (not shown).




Figure 4. SST and SLP anomalies composites during Feb-Mar and May-Jun when NAO(-) and AMM(+) are combined and precipitation anomaly(+) is registered in Eastern Caribbean (a-d), and when NAO(+) and AMM(-) are combined and precipitation anomaly(-) is registered in Eastern Caribbean (e-h).

Conclusions

The combined and individual effects of NAO and AMM indicate that the **Eastern Caribbean** is where precipitation anomalies (May-Jun) and NAO/AMM (Feb-Mar) are correlated. More frequent and consistent **regional effects on precipitation anomalies**, and more regionally spread and persistent SLP and SST were registered when **both NAO and AMM occurred together in the previous winter**. These results could help in seasonal forecasting, for regional drought or precipitation prediction.

*Contact information: flavia.moraes@uga.edu

Mote, T. L., Ramseier, C. A., & Miller, P. W. (2017). The Saharan Air Layer as an early rainfall season suppressant in the eastern Caribbean: The 2015 Puerto Rico drought. *Journal of Geophysical Research: Atmospheres*, 122, 10,966–10,982.
Vimont, D. J., and Kossin, J. P. (2007). The Atlantic Meridional Mode and hurricane activity. *Geophysical Research Letters*, 34, 1–5.
Rugg, A., Foltz, G. R., and Perez, R. C. (2016). Role of Mixed Layer Dynamics in Tropical North Atlantic Interannual Sea Surface Temperature Variability. *Journal of Climate*, 29, 8083–8101.