


DEVELOPING NITROGEN BUDGETS USING AN INTEGRATED BIOPHYSICAL MODEL TO INVESTIGATE CURRENT AND FUTURE PHYTOPLANKTON DYNAMICS IN A RAPIDLY CHANGING SUBTROPICAL ESTUARY, BARATARIA BASIN (GC23H-1305)




THE WATER INSTITUTE
OF THE GULF®

Hoonshin Jung^{1*}, Melissa Baustian¹, Ehab Meselhe², Tim Carruthers¹,

¹The Water Institute of the Gulf, Baton Rouge, LA, USA, ²Department of River-Coastal Science and Engineering, Tulane University

*hjung@thewaterinstitute.org

Background



Location of study area in the Barataria Basin, LA.


- Sediment diversions on the lower Mississippi River are proposed as a large-scale restoration strategy to create new wetlands and sustain existing wetland areas in Barataria and Breton Sound basins, Louisiana.
- Introduction of significant amount of nutrient-rich freshwater along with sediment to the receiving basins will likely drive changes in light and nutrient dynamics as well as phytoplankton composition, resulting in ecosystem-level changes.
- In order to understand nitrogen dynamics in the Barataria Basin due to large scale coastal restoration practices, quantification of nitrogen inputs, outputs and processes is essential because it is the limiting nutrient for most estuarine primary producers (e.g., phytoplankton and emergent macrophytes).
- Creating nitrogen budgets in the system will allow for better understanding of the major sources, sinks, inputs and exports across the system, increasing understanding of the amount of nitrogen available to drive estuarine primary production.

Objectives

- Develop a water quality model using an Integrated Biophysical Model
- Develop a nitrogen budget for Barataria Basin and quantify nitrogen inputs, outputs and processes
- Investigate potential impacts on nitrogen and phytoplankton dynamics due to the large-scale restoration project (proposed sediment diversion)


Methods

Integrated Biophysical Model




Conceptual overview of the Integrated Biophysical Model that includes the feedbacks between components (Baustian et al., 2018)

- Water quality variables
 - TOC, POC, DOC, PON, DON, NH₄, NO₃, POP, DOP, PO₄, Si, Silt, Clay, Sand, DO
- Seven sediment/soil layers
- 8 Phytoplankton groups
 - Freshwater Diatoms, Freshwater Flagellates, Green Algae, *Microcystis*, *Anabaena*
 - Marine Diatoms, Marine Flagellates, Dinoflagellates
- Seven vegetation taxa
 - Saline:** *Spartina alterniflora*
 - Brackish:** *Spartina patens*
 - Intermediate:** *Sagittaria lancifolia*, *Phragmites* spp., *Typha* spp.
 - Fresh:** *Sagittaria latifolia*, *Zizaniopsis miliacea*




D-WAQ model segments and vegetated area

Model Calibration and Validation



Taylor diagrams, showing standard deviation, correlation coefficient, and centered RMSD



Operation Plan of Mid-Barataria Sediment Diversion Coastal Changes and Phytoplankton Response



Flow conditions of the Mississippi River and proposed Mid-Barataria Sediment Diversion

The diversion diverts freshwater to a maximum capacity 2,123 cms (= 75,000 cfs) when the Mississippi River discharge is more than 16,900 cms (= 600,000 cfs) between February and August 2009, 2010, 2011 related to nitrogen budget figures below.

Nitrogen Budget



* Point sources including the proposed diversion

Conclusions

Introduction of freshwater into the Barataria Basin

- In case of "No Diversion"
 - Major source of DIN was the bottom sediment flux
 - Phytoplankton and wetland vegetation is a major consumer of DIN in water column and sediment/soil, respectively
 - In sediment/soil, settling of TON was major source of nitrogen
- In case of "With Diversion"
 - The operation of the diversion was one of major sources of DIN.
 - Increase in TON supply from vegetation might be related to increase in inundation mortality
 - Diversion operation increased the flushing rate of the system and thus increased TN export rate through the boundary
- Changes in Phytoplankton Dynamics
 - Dominant phytoplankton taxa in April changed from cyanobacteria of *Anabaena* and *Microcystis* to more freshwater diatoms in the mid and upper basin due to input of cold freshwater and silicate from the Mississippi River when proposed sediment diversion is operating.

Acknowledgements

This project was supported by the Science and Engineering Program of The Water Institute of the Gulf with funds from the Louisiana Coastal Protection and Restoration Authority and the Baton Rouge Area Foundation. We thank R. Eugene Turner and Erick Swenson at Louisiana State University for providing the water quality data to help calibrate the model.

The Water Institute of the Gulf is a not-for-profit, independent research institute dedicated to advancing the understanding of coastal, deltaic, river and water resource systems, both within the Gulf Coast and around the world. This mission supports the practical application of innovative science and engineering, providing solutions that benefit society.

1110 S. River Road., Ste. 200
Baton Rouge, Louisiana 70802

(225) 448-2813
WWW.THEWATERINSTITUTE.ORG



@theH2Oinstitute



The Water Institute of the Gulf