

Stable oxygen isotope composition is biased by shell calcification intensity in planktonic Foraminifera

Manuel F. G. Weinkauf^{1,a}, Jeroen Groeneveld², Joanna J. Waniek³, Torsten Vennemann⁴, & Rossana Martini¹

¹Department of Earth Sciences, University of Geneva, Rue des Maraîchers 13, 1205 Geneva, Switzerland

²Alfred Wegener Institute, Helmholtz-Center for Polar and Marine Sciences, Telegrafenberg A45, 14473 Potsdam, Germany

³Marine Chemistry Section, Leibniz Institute for Baltic Sea Research Warnemünde, Seestraße 15, 18199 Rostock, Germany

⁴Faculty of Geosciences and Environment, University of Lausanne, Bâtiment Geopolis, 1015 Lausanne, Switzerland

^aCurrent address: Institute of Geology and Palaeontology, Charles University, Albertov 2038/6, 128 43 Prague, Czech Republic

Mar 31st, 2020

ABSTRACT

Planktonic Foraminifera are widely used for environmental reconstructions through measurements of their shell's geochemical characteristics, including its stable oxygen and carbon isotope composition. Using these parameters as unbiased proxies requires a firm knowledge of all potential confounding factors influencing foraminiferal shell geochemistry. One such parameter is the shell calcification intensity (shell weight normalized for shell size) that may influence the shell $\delta^{18}\text{O}$ value either bioenergetically (by reducing energy available and required for equilibrium isotope fractionation during faster calcification) or kinetically (by influencing calcification depth through the shell's density contrast with seawater). Specimens from

the *Globigerinoides ruber/elongatus* compound from a sediment trap in the North Atlantic have been used to quantify the influence of shell calcification intensity on shell $\delta^{18}\text{O}$ values. Shell calcification intensity was found to have a significant effect on the shell stable oxygen isotope composition in all species. Through model fitting, it is suggested that the effect size may be in a range of 1 to 2‰ (depending on species, depth migration, and local oceanographic conditions). We show that the confounding effect of shell calcification intensity on stable oxygen isotope composition can be of importance, depending on the anticipated precision of the derived reconstructions. A framework is provided to quantify this effect in future studies.

INTRODUCTION

Planktonic Foraminifera are an abundant protist group in the world's oceans and their calcitic shells are commonly preserved in the sediment (Schiebel, 2002; Schiebel & Hemleben, 2017). For this reason, planktonic foraminiferal shells are abundantly used for paleoenvironmental reconstructions and biomonitoring purposes (e.g., Consolaro et al., 2018; Gorbarenko & Sounthor, 2000; Kučera, 2007; Weinkauf et al., 2016). A major application of planktonic foraminiferal shells is the use of their geochemical compositions for reconstructions of past environmental conditions (Lea, 2002; Ravelo & Hillaire-Marcel, 2007; Rohling & Cooke, 2002;

Rosenthal, 2007), which also provides the basis for climate models (e.g., Ezat et al., 2016; G. Ganssen & Troelstra, 1987; G. M. Ganssen et al., 2011; Groeneveld et al., 2006; Hemleben et al., 1996; Henehan et al., 2016; Ivanova, 1985; Milker et al., 2013; Steph et al., 2009).

The stable isotope composition of planktonic foraminiferal shells is usually not in equilibrium with seawater. For instance, for the shell's stable oxygen isotope composition, a vital effect often results in a different isotopic composition compared to a composition in equilibrium with the ambient seawater (compare Spero et al., 1991; Zeebe et al.,

2008). For different species, the size of this vital effect varies, necessitating species-specific correction functions to allow reliable paleoenvironmental interpretations (Carter et al., 2017; Spero, 1998; Steinke et al., 2005; Wit et al., 2013). For the application of stable oxygen isotope compositions of planktonic foraminiferal shells, for instance for paleoclimate (e.g., G. Ganssen & Troelstra, 1987; Milker et al., 2013) and depth habitat reconstructions (e.g., Mortyn & Charles, 2003; van Eijden, 1995), it is necessary to understand further factors that may bias the values measured within the shells. This is especially important when applying

single-shell geochemical analyses, as the geochemical variation between individual shells precipitated under comparable conditions is mostly larger than expected and beyond the analytical error of the measurement (Groeneveld et al., 2019; Killingley et al., 1981; Schiffelbein & Hills, 1984). For instance, the seasonality of species may play an important role in the observed stable isotope composition of a temporally integrated sediment sample (G. M. Ganssen et al., 2011; King & Howard, 2005), as does the intermixing of sedimented shells due to bioturbation (Billups & Spero, 1996; Stott & Tang, 1996). The interpretation of geochemical compositions of shells may even be further complicated by intra-shell variations of stable isotope compositions (Vetter et al., 2013) and ontogenetic changes in shell geochemistry (Rohling et al., 2004; Shackleton et al., 1985). While progress is being made in the understanding of the individual variation of the geochemical signal in planktonic Foraminifera, a larger part of the observable variation in single-shell analyses is still unaccounted for (compare Groeneveld et al., 2019, and citations therein). Incidentally, there have been few studies so far investigating to what degree the stable oxygen isotope composition of individual planktonic foraminiferal shells is influenced by their shell calcification intensity (the amount of calcite in the shell, normalized for shell size; Billups & Spero, 1995; Ezard et al., 2015).

Shell calcification intensity varies widely across planktonic foraminiferal species, and can in its own right be used as an environmental proxy (compare Marshall et al., 2013; Weinkauf et al., 2016). In this regard, shell calcification intensity has been mostly used as a proxy for dissolution (Broecker & Clark, 2001; Lohmann, 1995; Schiebel et al., 2007) and carbonate saturation of the seawater (Bijma et al., 2002; Bijma et al., 1999; Davis et al., 2019; de Moel et al., 2009; Osborne et al., 2016; Spero et al., 1997; Weinkauf et al., 2013). However, other studies could show that the picture is more complicated, and shell calcification intensity is indeed influenced by a variety of environmental factors (Aldridge et al., 2012; Marshall et al., 2013; Mohan

et al., 2015; Weinkauf et al., 2016), which may also have an influence on the shell's geochemical composition (e.g., water temperature, salinity).

The mechanisms of shell calcification in Foraminifera are still a matter of discussion, with seawater vacuolization, transmembrane ion transport, calcification via organic matrices, and active pH manipulation all touted as possibly valid calcification models (de Nooijer et al., 2014; de Nooijer et al., 2009; Ohno et al., 2016; Toyofuku et al., 2017). Since shell calcification in Foraminifera is so complicated and hardly understood, only comparatively little work has been performed so far to investigate its influence on the shell geochemical composition (e.g., Raitzsch et al., 2010; Steinhardt et al., 2015; van Dijk et al., 2017). Nevertheless, it is very feasible that calcification intensity of the shell can affect the foraminiferal shell's geochemical composition of the Foraminifera via two potential pathways.

The first possible influence stems from the calcification process itself. Normally, organisms preferentially use lighter isotopes of elements for inclusion in their organic and inorganic structural parts when compared to inorganic calcite (De La Rocha, 2003; Gabitov et al., 2012; Kendall & Caldwell, 1998). This is because inclusion of lighter isotopes is energetically favored, since bonds are weaker with the lighter isotopes (Kendall & Caldwell, 1998). In this regard, the calcification process involves an active selection of isotopes from the seawater, which is normally considered to be the so-called vital effect (fractionation). This process itself is time- and energy consuming, especially in those calcification models that require active membrane transport of components at the time of calcification (de Nooijer et al., 2014; Toyofuku et al., 2017). A higher calcification intensity in the shell can either result from a faster calcification rate for the same amount of time, or calcification at a constant rate for a longer time-interval. Under the assumption that a higher calcification intensity in the adult foraminiferal shell was derived from a faster instead of longer calcification duration during chamber formation (Hemleben et al., 1987; Spero,

1988; Spero et al., 1991), we may assume that the faster calcification rate must increase the size of the vital effect. This scenario leaves the foraminifer with less time to perform the calcification process, necessitating a more frequent inclusion of the much more abundant lighter isotope ^{16}O and thus lower $\delta^{18}\text{O}$ values of the shell when the shell's calcification intensity increases (Norris, 1998). Such a process has for instance been observed in corals in relation to their growth rate, where faster growing parts of the skeletal structure were enriched in ^{16}O (McConaughey, 1989).

The second influence stems from the effect of a heavier shell on the depth habitat of the Foraminifera. While planktonic foraminiferal species in general prefer a certain, species-specific habitat (Bé, 1977; Bé & Tolderlund, 1971; Numberger et al., 2009; Schiebel & Hemleben, 2017; Weiner et al., 2012), this depth habitat has been shown to be rather wide and reasonably flexible (compare Meiland et al., 2019; Rebotim et al., 2017). Moreover, there are signs for an ontogenetic change of the depth habitat in planktonic Foraminifera (Eggins et al., 2003; Schiebel & Hemleben, 2017; van Eijden, 1995). It can be reasonably assumed that a planktonic foraminifer with a higher calcification intensity lives in a deeper habitat than a foraminifer of a similar species with a lighter calcified shell simply due to gravitational pull and reduced buoyancy (Haenel, 1987; Weinkauf et al., in press; Zarkogiannis et al., 2019). Since the stable isotope composition of the seawater varies as a function of density, which in turn is largely controlled by the salinity and temperature of water through changes in evaporation and/or freshwater contributions, this could influence the stable isotope composition of all subsequently built chambers in correlation with the shell calcification intensity. If higher density shells, hence, find themselves at larger depths compared to lower density shells, this could influence the $\delta^{18}\text{O}$ values of planktonic foraminiferal shells. For example, foraminiferal shells with a higher shell calcification intensity would have formed within higher-salinity or lower-temperature waters at larger water depths and, thus, show increased

$\delta^{18}\text{O}$ values.

In this study, we use sediment trap material from the subtropical North Atlantic to study the correlation between shell calcification intensity and the $\delta^{18}\text{O}$

values of individual shells of planktonic Foraminifera, to quantify the influence and deduce the mechanisms leading to potential correlations with external crystallization factors. Investigating such con-

founding factors is of great importance to unambiguously use foraminiferal shell geochemical data as environmental proxies.

MATERIAL AND METHODS

SAMPLING MATERIAL

We used material from the 2000 m deep sediment trap Kiel 276–25 in the North Atlantic (33° N, 22° W; Fig. 1), spanning the time between May 2005 and April 2006 with a sampling resolution of 1–2 months (Table S1). The $>63\text{ }\mu\text{m}$ fraction of the dried and de-poisoned sediment trap samples were desalinated using MilliQ water and then dried overnight at 50 °C. Desalinated samples were then split into the 63–150 μm and $>150\text{ }\mu\text{m}$ size fraction using a Retsch AS 200 sieve tower (10 min. at intensity 40). Only the $>150\text{ }\mu\text{m}$ size fraction was used for further analyses to avoid ontogenetic effects on calcification intensity (Peeters et al., 1999). All planktonic Foraminifera from all samples were separated from the residue. Planktonic Foraminifera for shell calcification analyses were picked from representative samples of those concentrates, split with an ASC Scientific MS-1 microsplitter, such that approximately 100 specimens per sample were available for analyses. In samples with fewer specimens, 100 % of the sample was used (Table S2).

All specimens for analyses were cleaned with 2% ethanol and brushes and then rinsed in distilled water and on wet filter paper. They were then transferred into 10-hole cardboard slides and left to dry at room temperature for at least 24 hours.

Choice of species

The species were chosen to fulfill certain criteria. (1) They had to have a relatively restrictive depth habitat, so that our environmental reconstructions were effective in eliminating confounding factors, but still large enough to observe habitat depth changes. (2) They needed to be abundant enough for a reliable analysis. (3) They had to be species which are




Fig. 1: Sampling area including position of the mooring Kiel 276 and the catchment area of the 2000 m-sediment trap (J. J. Waniek et al., 2005). The Azores Front separates the North Atlantic temperate water from the North Atlantic subtropical gyre (NAST). Bathymetry from ETOPO1 (Amante & Eakins, 2009), temperature isolines from the “averaged decades” annual dataset of the World Ocean Atlas 2018 (Locarnini et al., 2018).

often used in geochemical analyses to maximize the practical value of the study and stay comparable with the literature.

We thus selected the three species *Globigerinoides ruber* (pink), *Globigerinoides ruber* (white), and *Globigerinoides elongatus* (Fig. 2). They all have photosymbionts (Schiebel & Hemleben, 2017) and are, thus, restricted to the photic zone, as is also confirmed by *in situ* studies (Meiland et al., 2019; Rebotim et al., 2017; Steinhardt et al., 2015). They are extensively used in geochemical analyses and especially *G. ruber* (white)

and *G. elongatus* were studied in detail concerning their differences in geochemical shell composition (Carter et al., 2017; Kawahata, 2005; Numberger et al., 2009; Steinke et al., 2005; Wang, 2000). We here follow the suggestions by Aurahs et al. (2011) to distinguish *G. ruber* (white) and *G. elongatus*: *Globigerinoides elongatus* has a flat last and penultimate chamber and especially the last chamber is very asymmetrical, while these chambers are inflated and symmetrical in *G. ruber* (white). The separation into *G. ruber* (white) and *G. elongatus* was




Fig. 2: Depiction of the three planktonic foraminiferal species used in this study (upper row: scanning electron micrographs, lower row: light microscopy images of the same specimens). *Globigerinoides ruber* (pink) is characterized by a reddish color of the chambers (often missing in the last and sometimes the penultimate chamber). *Globigerinoides ruber* (white) and *Globigerinoides elongatus* can be distinguished by the former having round and inflated chambers while in the latter the final and penultimate chamber are distinctly flattened and asymmetrical (Aurahs et al., 2011).

entirely done by the same scientists (principal author) to avoid any influence of competing species concepts.

Calcification intensity analysis

All specimens were weighed individually on a Mettler Toledo UMX2 microbalance (nominal precision 0.1 μg). The empty weighing boats (Elemental Microanalysis D5007) were first weighed ten times to reliably determine their weight. Afterwards, individual Foraminifera were transferred into the weighing boats and weighed four times each, and the difference between the empty and filled weighing boat was noted as the weight (W) of the foraminifer. The replication of weighing per individual allows to determine

the measurement error of the weighing procedure.

The same specimens were then photographed in apertural standard view at 27.5× magnification on a Leica M420 light microscope using a Leica DFC420 camera. The cross-sectional area (A) of each foraminifer was extracted in Fiji (Schindelin et al., 2012) running ImageJ v. 1.52o (Schneider et al., 2012), and the calcification intensity ρ_A sensu Marshall et al. (2013) was calculated as $\rho_A = \frac{W}{A}$ (compare Fig. S1).

Stable isotope analyses

Approximately 50 specimens per species for stable isotope analysis were selected using a stratified random sampling ap-

proach. For this, the sample was divided into 4 strata based on the range of observed specimen calcification intensity, and an equal number of specimens was randomly selected from each stratum. Thus, it could be made sure that all ρ_A classes were equally represented in the analyses. All geochemical measurements were performed at the Faculty of Geosciences and Environment of the University of Lausanne.

Selected specimens were individually cleaned via sonication (3 min.) in Krantz cells filled with tap water to remove any potential contaminants from the handling during the calcification intensity analysis. Specimens were then transferred into glass vials, closed with screw-on septum caps, and then analyzed for their $\delta^{13}\text{C}$

and $\delta^{18}\text{O}$ values on a Gas Bench II He-carrier gas system connected to a ThermoFisher DeltaPlus XL mass spectrometer (method by Spötl & Vennemann, 2003, adapted for small vials of 4 ml volume: acidifying with four drops of orthophosphoric acid of a specific gravity of 1.90, and measuring only four instead of ten peaks for each sample). Carrara marble was used as a standard and was measured alongside the foraminiferal shells every six samples and additionally two times at the start and at the end of each sequence. Standards and samples had the same weights and, hence, yields for the carbonate. All values are expressed in the familiar δ -notation and normalized against NBS-19 on the VPDB-scale. Repeated measurements for the standards after a size correction are within 0.1‰ for both oxygen and carbon isotope compositions.

Environmental setting

The sediment trap is located close to the Azores Front, which separates the temperate North Atlantic water from the North Atlantic subtropical gyre and is influenced by both water regimes (Figs. 1 and 3; Fründt & Wanek, 2012). Environmental data were calculated as averages for the respective sample duration and the estimated catchment area of the sediment trap (based on J. Wanek et al., 2000; J. J. Wanek et al., 2005) (Table S1). Sea surface temperature (SST) is based on the NOAA Optimum Interpolation (OI) Sea Surface Temperature (SST) V2 provided by the NOAA/OAR/ESRL PSD, Boulder, Colorado, USA (<https://www.esrl.noaa.gov/psd/>; Reynolds et al., 2002). Sea surface salinity (SSS) was calculated from the EN4 quality controlled ocean data (v. EN4.2.0; Good et al., 2013). The carbonate system of the seawater was estimated using the MS Excel program CO2Sys v. 2.1 (Lewis et al., 1998) with K1 and K2 constants from Mehrbach et al. (1973) (refit by Dickson & Millero, 1987) and KSO4 constant after Dickson (1990).

DATA ANALYSIS

All data analyses were performed in R v. 3.6.2 (R Core Team, 2019) and interpreted following Murthaugh (2014). Differences between parameters across species were tested using pairwise Mann–Whitney U tests (Mann & Whitney, 1947) with p -values corrected for the false discovery rate after Benjamini and Yekutieli (2001). The data for regression analyses were cleaned from potential outliers using bagplots of $\delta^{18}\text{O}$ against calcification intensity (Rousseeuw et al., 1999) as implemented in the R-package “aplypack” v. 1.3.3. Problems with multicollinearity in the data were eliminated following suggestions by Dormann et al. (2013): (1) The presence of a considerable degree of multicollinearity in the data was tested based on the condition number (Dormann et al., 2013). (2) Multicollinearity-clusters in the predictor variables were identified using principal component analysis (PCA; Booth et al., 1994) and cluster analysis on Hoeffding’s (1948) D -similarity index (Kauffmann & Rousseeuw, 2009). (3) The ranking of collinear parameters was estab-

lished based on the deviance in bivariate regressions with $\delta^{18}\text{O}$ values. (4) Sequential regression (Graham, 2003) was used to calculate the residuals of lower-ranked parameters and we used these as new parameters for the ensuing analyses (Dormann et al., 2013). We further applied general additive models (Friedman & Stuetzle, 1981) to identify all factors influencing the $\delta^{18}\text{O}$ values of the foraminiferal shells in the R-package “gam” v. 1.16.1, and tested the significance of the influence of shell calcification intensity via analysis of variances (ANOVA; Fisher, 1919) of the additive model against a null-model without calcification intensity. When the shell calcification intensity had a significant effect when accounting for all other parameters, linear and non-linear regression was used to fit different models to the data. Because most of the equations of the fitted models can only describe relationships in the first quadrant (to avoid e.g., division by zero), 10‰ had been added to all $\delta^{18}\text{O}$ values to transform them into positive values for this step. The corrected Akaike information criterion (AIC_c; Akaike, 1974) in combination with Akaike weights (Wagenmakers & Farrell, 2004) was used to determine the best fitting model from a set of mechanistically feasible candidate models and estimate the effect size. The relationship between $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ was investigated using a Deming regression (Deming, 1943) that took the associated measurement error in both values into account, using the R-package “deming” v. 1.4.

RESULTS

We analyzed the shell calcification intensity and shell geochemical composition of 136 specimens of the *Globigerinoides ruber/elongatus* compound from four samples covering an entire year from sediment trap Kiel 276-25 (Fig. S2, Table S1). In total, 42 specimens of *Globigerinoides ruber* (pink), 42 specimens of *Globigerinoides ruber* (white), and 52 specimens of *Globigerinoides elongatus* were successfully analyzed. The results have been

summarized in Fig. 4 and Table S3.

In terms of raw shell weight, we note that specimens of *G. ruber* (pink) are on average more than twice as heavy as specimens of *G. ruber* (white) and more than 25% heavier than *G. elongatus* specimens. The difference between all species is moderately significant at $p < 0.02$, indicating a general inherent difference in net shell weight. A different picture emerges when examining the

shell calcification intensities. *Globigerinoides elongatus* is most heavily calcified, followed by *G. ruber* (pink) and *G. ruber* (white). However, these differences are all very small and insignificant at $p > 0.3$, implying a comparable size-weight ratio between the species, which suits well with the fact that they are closely related and inhabit roughly the same portion of the water column, which requires a comparable density contrast




Fig. 3: Oceanography of the sampling area during summer and winter. Maps showing sea surface temperature and salinity as average of the upper 50 m of the water column. Depth profiles showing average values for the catchment area of the trap (indicated by the dotted rectangle in the maps). Temperature and salinity data from the 2005–2017 “decadal period” and oxygen as “average over all decades” of the World Ocean Atlas 2018 (Garcia et al., 2018; Locarnini et al., 2018; Zweng et al., 2018).

with the surrounding seawater.

For the stable isotope composition, we also see considerable differences between the species. The $\delta^{13}\text{C}$ values of *G. ruber* (pink) are positive, while they are negative for the other species, with *G. ruber* (white) having the lowest values. The difference in average $\delta^{13}\text{C}$ values is moderately significant for all pairwise comparisons ($p < 0.02$). For the stable oxygen isotopes, we see another signal. Values are highest for *G. elongatus* and lowest for *G. ruber* (pink), but the differences between neither *G. ruber* (pink) and *G. ruber* (white) ($p = 0.131$) nor *G. ruber* (white) and *G. elongatus* ($p = 1.000$) are significant. Only be-

tween *G. elongatus* and *G. ruber* (pink) do differences in $\delta^{18}\text{O}$ values of the shell accumulate to a convincingly significant signal ($p = 0.002$).

We applied general additive models to evaluate the influence of shell calcification intensity on the stable oxygen isotope composition of shells within the *Globigerinoides ruber/elongatus* compound. To disentangle this influence from the effects of water temperature, salinity, and symbiont-photosynthetic activity (approximated by shell $\delta^{13}\text{C}$ values), which all influence the $\delta^{18}\text{O}$ values of foraminiferal shells, we used the model described in Eq. (1).

$$\delta^{18}\text{O} = \ell(T) + \ell(S) + \ell(a_P) + \ell(\rho_A) + \epsilon \quad (1)$$

with T as sea surface temperature, S as sea surface salinity, a_P as photosynthetic activity, ρ_A as shell calcification intensity, ϵ as error term, and ℓ indicating a LOESS smooth.

To facilitate the analysis of the impact of environmental factors and shell calcification on the isotopic composition of the foraminiferal shells, it was necessary to establish the multicollinearity between the different parameters included in Eq. (1) in order to remove confounding factors from the analyses. We tested the




Fig. 4: Comparison of shell weight (a), shell calcification intensity (b), and shell $\delta^{13}\text{C}$ (c) and $\delta^{18}\text{O}$ (d) values of shells of *Globigerinoides ruber* (pink), *Globigerinoides ruber* (white), and *Globigerinoides elongatus* from sediment trap Kiel 276–25. Thick lines indicate median, boxes cover interquartile range (IQR), whiskers extend to $1.5 \times \text{IQR}$, outliers are marked by black diamonds. Small letters above the boxes denote groups according to pairwise Mann–Whitney *U* tests.

presence of multicollinearity between any of the parameters by calculating the condition number of the correlation matrix as $\kappa = 11.11$, far in excess of the suggested threshold of $\kappa \geq 5$ that would imply the presence of collinearity. An ensuing PCA, where elements which show a loading $L > 0.3$ on the first principal component (PC) are supposed to be collinear, suggests a noteworthy collinearity between temperature ($L_{\text{PC}1} = 0.68$), salinity ($L_{\text{PC}1} = 0.65$), and photosynthetic activity ($L_{\text{PC}1} = 0.33$). A cluster analysis (complete linkage) on the Hoeffding D -similarity index supports this collinear-

ity, with only shell calcification intensity showing a similarity below 0 with the other parameters (Fig. S3).

To eliminate the collinearity between temperature, salinity, and photosynthetic activity before calculation of the general additive models, we used general linear regression to regress collinear parameters against $\delta^{18}\text{O}$ as dependent variable. Through the deviance d of these regressions, we could establish the relative importance of the parameters on the $\delta^{18}\text{O}$ of the foraminiferal shells as (1) temperature ($d = 45.36$), (2) photosynthetic activity ($d = 42.81$), and (3) salinity ($d = 42.35$).

Through sequential regression using general linear models, we then calculated the residuals for photosynthetic activity and salinity for use in the additive models instead of the raw values of these parameters. Based on these corrections for multicollinearity, the additive models we calculated (based on Eq. 1) are described in Eq. (2).

$$\delta^{18}\text{O} = \ell(T) + \ell(e_{T+ap}(S)) + \ell(e_T(ap)) + \ell(\rho_A) + e \quad (2)$$

with e as residuals and other parameters as for Eq. (1).

Results of the additive models according to Eq. (2) are summarized in Table 1. For *G. ruber* (pink), no outliers were detected, but we encountered the problem that because one sample did not contain any specimens, the data set was not suitable for an additive model approach because it had only three distinct temperature and salinity values. We thus limited the analyses to the photosynthetic activity and calcification intensity parameters in this species. The model suggests an influence of shell calcification intensity on shell $\delta^{18}\text{O}$ values and a significant increase in model fit when adding shell calcification intensity in comparison with the model that ignores it ($p = 0.031$).

Ultimately, we can use the results from the other species to get a clearer picture of the influence of shell calcification intensity on the stable oxygen isotope composition of the foraminiferal shells considering all other parameters, even though this was impossible in *G. ruber* (pink). In *G. ruber* (white), no outliers were detected. Using an additive model, we could identify that all included parameters have a moderately to convincingly significant influence on the shell $\delta^{18}\text{O}$ values. An ANOVA against the null-model confirms the significant increase in model fit when shell calcification intensity is included at $p < 0.001$. In *G. elongatus*, two specimens (Specimen 77 from Sample 3 and

Specimen 88 from Sample 5) were identified as outliers and removed from further analyses. The additive model confirms a convincingly significant influence of temperature, photosynthetic activity, and calcification intensity, but not salinity, on the stable oxygen isotope composition of the shells. Shell calcification intensity improves the model significantly in comparison to the null-model ($p = 0.049$). In all species analyzed here, we thus find a significant influence of shell calcification intensity on the shell $\delta^{18}\text{O}$ values when accounting for all other relevant parameters.

DISCUSSION

The average $\delta^{18}\text{O}$ values are negative in *Globigerinoides ruber* (pink), *Globigerinoides ruber* (white), and *Globigerinoides elongatus* in our study, and are compatible with a typical foraminiferal vital effect and with a preference for lighter ^{16}O during calcification in all species. The general trend in all species is an increase in $\delta^{18}\text{O}$ (toward zero) with increasing shell calcification intensity. This trend indicates that the influence of shell calcification intensity on $\delta^{18}\text{O}$ is linked to depth habitat changes when the density contrast between the shell and ambient seawater changes. If calcification intensity influences $\delta^{18}\text{O}$ values through a modification of living depths via changing density contrasts with the surrounding seawater (Weinkauf et al., in press; Zarkogiannis et al., 2019), we would expect lighter individuals that live in shallower depths and thus warmer water (especially in summer, we could expect temperature changes of ca. 6 °C within the upper 100 m of the water column; compare Fig. 3), to have lower $\delta^{18}\text{O}$ values; this is indeed noted by the present data. In contrast, if faster calcification rates would limit the time or energy available for fractionation, heavier calcified individuals would show reduced $\delta^{18}\text{O}$ values because of unselected uptake of the more abundant ^{16}O (Norris, 1998).

To better understand the observed relationship between shell calcification inten-

sity and shell $\delta^{18}\text{O}$ values and estimate the effect size, we fitted linear and non-linear regressions to the data (Bolker, 2008) and determined, which would best describe the observations using the corrected Akaike information criterion (AIC_c). We tried the following regression functions: (1) A Michaelis–Menten (Michaelis & Menten, 1913) function and (2) a power-law function, which through their bioenergetic applicability would both be suitable candidate functions (Raitzsch et al., 2010; Russell et al., 2004). (3) A monomolecular function and (4) a second-degree polynomial function, which are functionally suitable for fractionation processes during calcification (Groeneveld et al., 2018; Pearson, 2012; Steinhardt et al., 2015). The results of this model assessment are summarized in Fig. 5 and Table 2. In *G. ruber* (pink), the data are best described by a polynomial function and second-best by a monomolecular function. The estimated effect size using the two best-fitting models is 0.63–1.01 ‰. A similar signal is obtained for *G. ruber* (white), where a monomolecular function describes the data much better than the second-best model (Michaelis–Menten function; normalized probability: 0.587). The effect size in *G. ruber* (white) is approximately in the range of 1.37–1.42 ‰. In *G. elongatus*, the power-law and Michaelis–Menten models outperform both alternative models, indicating

a smaller effect size of ca. 0.56 ‰.

We consider the measured shell geochemical composition in our data largely free from constraining influences. It is known that photosynthetic activity influences the $\delta^{18}\text{O}$ value measured in foraminiferal shells (Ezard et al., 2015; Spero, 1998; Spero & Lea, 1993), but $\delta^{13}\text{C}$ is directly scaled to photosynthetic activity or bioproductivity, and hence also to depth (e.g., Norris, 1998; Spero, 1998; Spero & Lea, 1993; Spero et al., 1991). Therefore, by including shell $\delta^{13}\text{C}$ values in our general additive models, we can test for the effect of calcification intensity on shell $\delta^{18}\text{O}$ values, regardless. A shell size effect on $\delta^{13}\text{C}$ has been noted, which does not exist for $\delta^{18}\text{O}$ to any noteworthy extent in the species used here (Ezard et al., 2015; Franco-Fraguas et al., 2011; Norris, 1998; Shackleton et al., 1985), meaning that our analyses are robust against the shell size range used in this study. Carbonate ion concentration can have an effect on stable isotope compositions of planktonic Foraminifera, especially $\delta^{13}\text{C}$ values (Spero, 1998), but the sampling region is considerably stable concerning its carbonate system. We used data from the ESTOC time series (González-Dávila, 2016a, 2016b), which is situated close to sediment trap Kiel 276, and the MS Excel program CO2Sys to estimate the carbonate system parameters. The maximum annual range of

Tab. 1: Results of an additive model analysis of stable oxygen isotope composition against temperature (T), salinity (S), photosynthetic activity (a_P), and shell calcification intensity (ρ_A) in foraminiferal shells from sediment trap Kiel 276–25.

Parameter	Parametric terms			Nonparametric terms	
	Mean squares	F-value	p-value	F-value	p-value
<i>Globigerinoides ruber</i> (pink)					
a_P	4.10	44.58	<0.001	3.48	0.023
ρ_A	0.25	2.69	0.111	5.03	0.006
<i>Globigerinoides ruber</i> (white)					
T	61.36	371.03	<0.001	20.90	<0.001
S	1.59	9.61	0.004	21.00	<0.001
a_P	26.83	162.22	<0.001	0.81	0.487
ρ_A	5.66	34.25	<0.001	2.12	0.130
<i>Globigerinoides elongatus</i>					
T	48.59	536.71	<0.001	200.39	<0.001
S	0.00	0.00	0.989	97.95	<0.001
a_P	1.57	17.34	<0.001	1.84	0.144
ρ_A	0.75	8.32	0.007	0.98	0.414

Tab. 2: Results from a fitting of four model functions to the influence of shell calcification intensity on shell $\delta^{18}\text{O}$ in foraminiferal shells from sediment trap Kiel 276–25. The corrected Akaike information criterion (AIC_c ; Akaike, 1974) and model weights (Wagenmakers & Farrell, 2004) are reported.

Parameter	Michaelis–Menten function	Power-law function	Monomolecular function	2 nd -degree polynomial function
<i>Globigerinoides ruber</i> (pink)				
AIC_c	55.175	56.953	52.577	51.526
ΔAIC_c	3.649	5.427	1.051	0.000
Weight	0.089	0.036	0.325	0.550
<i>Globigerinoides ruber</i> (white)				
AIC_c	94.469	95.207	93.767	96.263
ΔAIC_c	0.702	1.440	0.000	2.496
Weight	0.284	0.196	0.404	0.116
<i>Globigerinoides elongatus</i>				
AIC_c	41.277	41.093	42.097	43.340
ΔAIC_c	0.184	0.000	1.003	2.247
Weight	0.321	0.352	0.213	0.114

CO_3^{2-} in surface waters in the region is $<50 \mu\text{mol kg}^{-1}$, and the average range is as small as $<10 \mu\text{mol kg}^{-1}$ throughout spring–autumn (Table S4), ruling out a large effect of this parameter. We therefore believe that a better interpretation of the data is required.

One possible solution to explain the observations would be dissolution of the shells (Broecker & Clark, 2001; Lohmann, 1995; Schiebel et al., 2007), as selective dissolution of poorly crystallized, small crystallites of calcite with a particular stable oxygen isotope composition could result in a spurious $\delta^{18}\text{O}$ –calcification intensity–correlation. This is unlikely,

however, for the following reasons. (1) We used the same stratified random sampling applied to pick the specimens for isotope analyses to choose a second random sample per species to be investigated under the scanning electron microscope. The vast majority of the shells shows good to excellent preservation, with minor recrystallization observable in only a small number of shells (Figs. S4–S6). (2) This is supported by a reconstruction of the carbonate system using the ESTOC time series (González-Dávila, 2016a, 2016b). Throughout the entire sampling interval, the calcite saturation state at the surface is >4 , and even at the sediment

trap depth of 2000 m it never decreases below 1.38 at any time (Table S4). The latter value is near the lower boundary below which dissolution starts to affect shell geochemistry (Dekens et al., 2002; Regenberg et al., 2006), but not yet problematic. (3) We further note that, should dissolution play a major role, we would expect to see the inverse effect. This is because calcite that was precipitated in warmer waters has a higher Mg/Ca ratio, and is generally dissolved more readily than calcite precipitated in colder waters (de Villiers et al., 2002; Rosenthal et al., 2003). Since calcite that was formed in warmer water would also have lower




Fig. 5: Models for the shell $\delta^{18}\text{O}$ values depending on the shell calcification intensity in the planktonic foraminifers (a) *Globigerinoides ruber* (pink), (b) *Globigerinoides ruber* (white), and (c) *Globigerinoides elongatus* from sediment trap Kiel 276–25. (d) Relationship between shell $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ in the same samples. Error bars indicate standard error of the measurements.

$\delta^{18}\text{O}$ values, this process of selective dissolution would increase the $\delta^{18}\text{O}$ values of the lighter shells, such that the pattern we observe here would have been even more pronounced if such a selective dissolution would have taken place.

The other possibility is an environmental effect. On the one hand, this could for instance be a salinity effect. Although salinity was determined as least important and non-significant parameter, and is strongly collinear with temperature, this is only true for the remote and averaged environmental data available here. It was shown that such data often underestimate the true short-term variation in oceanic settings (Laepple

& Huybers, 2014a, 2014b). *Globigerinoides ruber* is very resistant against salinity changes (Bijma et al., 1990) and planktonic Foraminifera can build new chambers within hours (Bé et al., 1979; Spero, 1988), which would make it theoretically feasible that the local salinity could have changed sufficiently on smaller time-scales (e.g. via north–south-fluctuation of the Azores front; Fründt & Waniek, 2012) without influencing chamber formation in the Foraminifera, thus explaining part of the measured range of compositions. However, this is unlikely to be responsible for the majority of the correlation due to the inconceivable amplitude of salinity changes in an open-ocean

setting this assumption would require. In fact, even through fluctuation of the Azores front, salinities beyond 36.2 are practically never observed in this oceanographic setting (Fründt & Waniek, 2012; Pérez et al., 2003). Temperature changes due to vertical migration, on the other hand, could explain a major part of the measured differences in isotopic compositions for shells with different calcification intensities (Rebotim et al., 2019). The depth habitat of species in the *Globigerinoides ruber/elongatus* compound is generally shallow, partly due to its dependence on sunlight for photosynthesis, but both living- and calcification-depths were shown to be reasonably variable

within the uppermost 50–100 m of the water column (Meilland et al., 2019; Rebotim et al., 2017; Steinhardt et al., 2015). Within the studied area, the mixed layer covers this depth in winter, with an average temperature change of only 0.3 °C over the top 100 m. During summer, however, the average temperature variation across the first 100 m is ca. 6 °C, and thus large enough to explain perhaps 1.5–2 % of the observed variation in shell $\delta^{18}\text{O}$ (compare Fig. 3). Even in winter, changes beyond the recorded mean values can occur due to the fluctuation of the Azores front across the sediment trap mooring (Fründt & Waniek, 2012).

Interestingly, $\delta^{13}\text{C}$ values can be interpreted as a water depth indicator, especially pronounced in photosymbiont-bearing Foraminifera (Spero & Williams, 1988). For this reason, one would expect to note a positive correlation between $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ in such foraminiferal assemblages, where the recorded temperature changes are at least partly related to depth migration (Norris, 1998; Spero & Williams, 1988). We see a pronounced pattern of this type in *G. ruber* (pink) and *G. ruber* (white), but a much weaker signal in *G. elongatus* for our dataset (Fig. 5d). Coincidentally, *G. elongatus* is also the species that shows the smallest effect size of shell calcification intensity on shell $\delta^{18}\text{O}$ values. It was shown by Steinke et al. (2005) that *G. elongatus* lives generally deeper than *G. ruber*, but because the species is still dependent on light for its photosymbionts this implies that extensive depth migration is less of a possibility for this species, which could explain why a more stable depth habitat in *G. elongatus* is compatible with our

data. We hypothesize here that the measured compositions in all species are the result of a depth migration-related temperature change (directly scaled to shell calcification intensity via density contrast with the seawater). We further hypothesize that *G. elongatus* shows the smallest effect size due to its limited depth migration. If this interpretation is true, then shell calcification intensity can be a viable proxy to reduce the bias in geochemical shell measurements of planktonic Foraminifera that is associated with habitat depth. Our study can consequently be used in the implementation of bias-modelling of climate archive proxies as for instance through "Sedproxy" (Dolman & Laepple, 2018), and can help to further improve the reliability of climate reconstructions (Dolman et al., 2020). Such an interpretation is also in line with earlier observations that larger shells within some species have higher $\delta^{13}\text{C}$ values (e.g., Spero & Lea, 1993). Larger shells contain more cytoplasm that has a negative density contrast to the surrounding seawater. Unless the foraminifer increases its relative shell thickness (to keep the shell calcification intensity constant), it will be subject to greater buoyant forces, reducing its depth habitat as shown by the higher $\delta^{13}\text{C}$ values of its shell. This is also implied by observations in Zarkogiannis et al. (2019), where it is shown that larger shells tend to be more buoyant because their weight does not increase at the same rate as their volume if the shell thickness remains constant.

Further studies are suggested in this field, ideally involving controlled laboratory experiments, especially regarding

if this effect also exists for asymbiotic Foraminifera – as is implied by Billups and Spero (1995) at least for small specimens – or species which tend to develop thick crusts (Schiebel & Hemleben, 2017, e.g., *Globoconella inflata*, *Globorotalia crassaformis*, *Globorotalia truncatulinoides*). If our hypothesis is true, the effect of shell calcification intensity on the stable oxygen isotope composition of foraminiferal shells through depth habitat changes is rather variable between locations and species. This means that the effect on temperature reconstructions can be small enough that for most purposes a correction for shell calcification intensity does not need to be applied, but a bias of around 1 °C seems feasible under certain oceanographic conditions. Such effects may be important for some studies (e.g., Dolman et al., 2020), especially in regions or species where shell calcification intensity varies more strongly within a species and in environments with strong vertical changes in the ambient seawater environment. We presented in this study a framework to deal with this problem in the future, by quantifying shell calcification intensity of specimens intended for geochemical analyses and quantifying its effect size for data correction. Since especially the portion of the influence that is caused by depth migration will be dependent on the investigated species (theoretical range of depth habitat), observed range of calcification intensity (practical range of depth habitat), and the local oceanography, no generally applicable corrections can be suggested here. We thus suggest an implementation of this factor in bias-modelling protocols in the future (e.g., Dolman & Laepple, 2018).

CONCLUSIONS

Our analyses of foraminiferal species of the *Globigerinoides ruber/elongatus* compound imply that shell calcification intensity has an impact on shell $\delta^{18}\text{O}$ values and therefore has the potential to be a confounding factor in environmental reconstructions derived from stable oxygen isotope measurements of planktonic foraminiferal shells.

Our data suggest that the observed

effect is mainly an environmental effect, where heavier calcified shells would occupy a deeper habitat of the water column and are thus exposed to lower temperatures of their ambient seawater. The effect size will depend on the species (range of depth habitat; compare Rebotim et al., 2019), the range of shell calcification intensity, and the local oceanography.

We suggest using shell calcification intensity studies to evaluate and eliminate this effect on geochemical analyses if high-precision reconstructions are needed, but argue that the effect is probably small enough to not interfere with the majority of practical applications. We suggest further studies in this field and an implementation of this bias in standard bias-modelling protocols.

ACKNOWLEDGMENTS

We thank Howard J. Spero (University of California – Davis, USA) for discussions about vital effects in planktonic Foraminifera and his support in securing funding by the Palaeontological Association. Agathe Martignier (University of Geneva, Switzerland) is thanked for her help with scanning electron microscope work. Geochemical analyses of the foraminiferal shells were partly funded by a Sylvester–Bradley Award of the Palaeontological Association (granted to MFGW, reference-no. PA-SB201705), which is thankfully recognized. JW is thankful for receiving funding for the work on the mooring Kiel 276 from the Deutsche Forschungsgemeinschaft (grant numbers WA2175/1-1 to WA2175/3-1 and WA2157/4).

AUTHOR CONTRIBUTIONS

Conceptualization:	MFGW, JC
Methodology:	MFGW, TV
Software:	MFGW
Validation:	MFGW, JC, JJW, TV, RM
Formal analysis:	MFGW, JC
Investigation:	MFGW, JJW, TV
Resources:	JJW, RM
Data curation:	MFGW
Writing – original draft preparation:	MFGW, JC, JJW, TV, RM
Visualization:	MFGW
Project administration:	RM
Funding acquisition:	MFGW, JJW

CITED LITERATURE

- Akaike, H. (1974). A new look at the statistical model identification. *IEEE Transactions on Automatic Control*, 19, 716–723. <https://doi.org/10.1109/TAC.1974.1100705>
- Aldridge, D., Beer, Ch. J., & Purdie, D. A. (2012). Calcification in the planktonic Foraminifera *Globigerina bulloides* linked to phosphate concentrations in surface waters of the North Atlantic Ocean. *Biogeosciences*, 9, 1725–1739. <https://doi.org/10.5194/bg-9-1725-2012>
- Amante, C., & Eakins, B. W. (2009). *ETOPO1 1 arc-minute global relief model: Procedures, data sources and analysis* (NOAA Technical Memorandum NESDIS NGDC-24). National Geophysical Data Center. <http://www.ngdc.noaa.gov/mgg/global/relief/ETOPO1/docs/ETOPO1.pdf>
- Aurahs, R., Treis, Y., Darling, K., & Kučera, M. (2011). A revised taxonomic and phylogenetic concept for the planktonic foraminifer species *Globigerinoides ruber* based on molecular and morphometric evidence. *Marine Micropaleontology*, 79, 1–14. <https://doi.org/10.1016/j.marmicro.2010.12.001>
- Bé, A. W. H. (1977). An ecological, zoogeographic and taxonomic review of recent planktonic Foraminifera. In A. T. S. Ramsay (Ed.), *Oceanic micropalaeontology* (pp. 1–100). London, New York, San Francisco, Academic Press.
- Bé, A. W. H., Hemleben, Ch., Anderson, O. R., & Spindler, M. (1979). Chamber formation in planktonic Foraminifera. *Micropaleontology*, 25, 294–307. <http://www.jstor.org/stable/1485304>
- Bé, A. W. H., & Tolderlund, D. S. (1971). Distribution and ecology of living planktonic Foraminifera in surface waters of the Atlantic and Indian Oceans. In B. M. Funnell & W. R. Riedel (Eds.), *The micropalaeontology of oceans* (pp. 105–149). London, Cambridge University Press.
- Benjamini, Y., & Yekutieli, D. (2001). The control of the false discovery rate in multiple testing under dependency. *The Annals of Statistics*, 29, 1165–1188. <https://doi.org/10.1214/aos/1013699998>
- Bijma, J., Faber, W. W., Jr., & Hemleben, Ch. (1990). Temperature and salinity limits for growth and survival of some planktonic foraminifers in laboratory cultures. *Journal of Foraminiferal Research*, 20, 95–116. <https://doi.org/10.2113/gsjfr.20.2.95>
- Bijma, J., Hönnisch, B., & Zeebe, R. E. (2002). Impact of the ocean carbonate chemistry on living foraminiferal shell weight: Comment on “Carbonate ion concentration in glacial-age deep waters of the Caribbean Sea” by W. S. Broecker and E. Clark. *Geochemistry, Geophysics, Geosystems*, 3, Article 1064. <https://doi.org/10.1029/2002GC000388>
- Bijma, J., Spero, H. J., & Lea, D. W. (1999). Reassessing foraminiferal stable isotope geochemistry: Impact of the oceanic carbonate system (experimental results). In G. Fischer & G. Wefer (Eds.), *Use of proxies in paleoceanography: Examples from the South Atlantic* (pp. 489–512). Berlin, Heidelberg, Springer-Verlag.

- Billups, K., & Spero, H. J. (1995). Relationships between shell size, thickness and stable isotopes in individual planktonic Foraminifera from 2 Equatorial Atlantic cores. *Journal of Foraminiferal Research*, 25, 24–37. <https://doi.org/10.2113/gsjfr.25.1.24>
- Billups, K., & Spero, H. J. (1996). Reconstructing the stable isotope geochemistry and paleotemperatures of the equatorial Atlantic during the last 150,000 years: Results from individual Foraminifera. *Paleoceanography*, 11, 217–238. <https://doi.org/10.1029/95PA03773>
- Bolker, B. M. (2008). *Ecological models and data in R*. New Jersey, Woodstock, Princeton University Press.
- Booth, G. D., Niccolucci, M. J., & Schuster, E. G. (1994). *Identifying proxy sets in multiple linear regression: An aid to better coefficient interpretation* (tech. rep.). United States Department of Agriculture. Washington, DC.
- Broecker, W., & Clark, E. (2001). An evaluation of Lohmann's Foraminifera weight dissolution index. *Paleoceanography*, 16, 531–534. <https://doi.org/10.1029/2000PA000600>
- Carter, A., Clemens, S., Kubota, Y., Holbourn, A., & Martin, A. (2017). Differing oxygen isotopic signals of two *Globigerinoides ruber* (white) morphotypes in the East China Sea: Implications for paleoenvironmental reconstructions. *Marine Micropaleontology*, 131, 1–9. <https://doi.org/10.1016/j.marmicro.2017.01.001>
- Consolaro, Ch., Rasmussen, T. L., & Panieri, G. (2018). Palaeoceanographic and environmental changes in the eastern Fram Strait during the last 14,000 years based on benthic and planktonic Foraminifera. *Marine Micropaleontology*, 139, 84–101. <https://doi.org/10.1016/j.marmicro.2017.11.001>
- Davis, A. N., Davis, C. V., Thunell, R. C., Osborne, E. B., Black, D. E., & Benitez-Nelson, C. R. (2019). Reconstructing 800 years of carbonate ion concentration in the Cariaco basin using the area density of planktonic Foraminifera shells. *Paleoceanography and Paleoceanography*, 34, 2129–2140. <https://doi.org/10.1029/2019PA003698>
- De La Rocha, Ch. L. (2003). Silicon isotope fractionation by marine sponges and the reconstruction of the silicon isotope composition of ancient deep water. *Geology*, 31, 423–426. [https://doi.org/10.1130/0091-7613\(2003\)031<0423:SIFBMS>2.0.CO;2](https://doi.org/10.1130/0091-7613(2003)031<0423:SIFBMS>2.0.CO;2)
- Dekens, P. S., Lea, D. W., Pak, D. K., & Spero, H. J. (2002). Core top calibration of Mg/Ca in tropical Foraminifera: Refining paleotemperature estimation. *Geochemistry, Geophysics, Geosystems*, 3, 1–29. <https://doi.org/10.1029/2001GC000200>
- Deming, W. E. (1943). *Statistical adjustment of data*. New York, John Wiley & Sons, Ltd.
- de Moel, H., Ganssen, G. M., Peeters, F. J. C., Jung, S. J. A., Kroon, D., Brummer, G.-J. A., & Zeebe, R. E. (2009). Planktic foraminiferal shell thinning in the Arabian Sea due to anthropogenic ocean acidification? *Biogeosciences*, 6, 1917–1925. <https://doi.org/10.5194/bg-6-1917-2009>
- de Nooijer, L. J., Spero, H. J., Erez, J., Bijma, J., & Reichart, G.-J. (2014). Biomineralization in perforate Foraminifera. *Earth-Science Reviews*, 135, 48–58. <https://doi.org/10.1016/j.earscirev.2014.03.013>
- de Nooijer, L. J., Toyofuku, T., Kitazato, H., & Stanley, S. M. (2009). Foraminifera promote calcification by elevating their intracellular pH. *Proceedings of the National Academy of Sciences of the United States of America*, 106, 15374–15378. <https://doi.org/10.1073/pnas.0904306106>
- de Villiers, S., Greaves, M., & Elderfield, H. (2002). An intensity ratio calibration method for the accurate determination of Mg/Ca and Sr/Ca of marine carbonates by ICP-AES. *Geochemistry, Geophysics, Geosystems*, 3, 2001GC000169. <https://doi.org/10.1029/2001GC000169>
- Dickson, A. G., & Millero, F. J. (1987). A comparison of the equilibrium constants for the dissociation of carbonic acid in seawater media. *Deep Sea Research, Part A: Oceanographic Research Papers*, 34, 1733–1743. [https://doi.org/10.1016/0198-0149\(87\)90021-5](https://doi.org/10.1016/0198-0149(87)90021-5)
- Dickson, A. G. (1990). Standard potential of the reaction: $\text{AgCl}(\text{s}) + 12\text{H}_2(\text{g}) = \text{Ag}(\text{s}) + \text{HCl}(\text{aq})$, and the standard acidity constant of the ion HSO_4^- in synthetic sea water from 273.15 to 318.15 K. *The Journal of Chemical Thermodynamics*, 22, 113–127. [https://doi.org/10.1016/0021-9614\(90\)90074-Z](https://doi.org/10.1016/0021-9614(90)90074-Z)
- Dolman, A. M., Kunz, T., Groeneveld, J., & Laepple, Th. (2020). Estimating the timescale-dependent uncertainty of paleoclimate records – a spectral approach. Part II: Application and interpretation. *Climate of the Past Discussions*. <https://doi.org/10.5194/cp-2019-153>
- Dolman, A. M., & Laepple, Th. (2018). Sedproxy: A forward model for sediment-archived climate proxies. *Climate of the Past*, 14, 1851–1868. <https://doi.org/10.5194/cp-14-1851-2018>
- Dormann, C. F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carré, G., Marquéz, J. R. G., Gruber, B., Lafourcade, B., Leitão, P. J., Münkemüller, T., McClean, C., Osborne, P. E., Reineking, B., Schröder, B., Skidmore, A. K., Zurell, D., & Lautenbach, S. (2013). Collinearity: A review of methods to deal with it and a simulation study evaluating their performance. *Ecography*, 36, 27–46. <https://doi.org/10.1111/j.1600-0587.2012.07348.x>
- Eggins, S., De Deckker, P., & Marshall, J. (2003). Mg/Ca variation in planktonic Foraminifera tests: Implications for reconstructing palaeo-seawater temperature and habitat migration. *Earth and Planetary Science Letters*, 212, 291–306. [https://doi.org/10.1016/S0012-821X\(03\)00283-8](https://doi.org/10.1016/S0012-821X(03)00283-8)
- Ezard, Th. H. G., Edgar, K. M., & Hull, P. M. (2015). Environmental and biological controls on size-specific $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ in recent planktonic Foraminifera. *Paleoceanography*, 30, 151–173. <https://doi.org/10.1002/2014PA002735>
- Ezat, M. M., Rasmussen, T. L., & Groeneveld, J. (2016). Reconstruction of hydrographic changes in the southern

- Norwegian Sea during the past 135 kyr and the impact of different foraminiferal Mg/Ca cleaning protocols. *Geochemistry, Geophysics, Geosystems*, 17, 3420–3436. <https://doi.org/10.1002/2016GC006325>
- Fisher, R. A. (1919). The correlation between relatives on the supposition of Mendelian inheritance. *Transactions of the Royal Society of Edinburgh: Earth Sciences*, 52, 399–433. <https://doi.org/10.1017/S0080456800012163>
- Franco-Fraguas, P., Costa, K. B., & de Lima Toledo, F. A. (2011). Stable isotope/test size relationship in *Cibicidoides wuellerstorfi*. *Brazilian Journal of Oceanography*, 59, 287–291. <https://doi.org/10.1590/S1679-87592011000300010>
- Friedman, J. H., & Stuetzle, W. (1981). Projection pursuit regression. *Journal of the American Statistical Association*, 76, 817–823. <https://doi.org/10.1080/01621459.1981.10477729>
- Fründt, B., & Waniek, J. J. (2012). Impact of the Azores Front propagation on deep ocean particle flux. *Central European Journal of Geosciences*, 4, 531–544. <https://doi.org/10.2478/s13533-012-0102-2>
- Gabitov, R. I., Watson, E. B., & Sadekov, A. (2012). Oxygen isotope fractionation between calcite and fluid as a function of growth rate and temperature: An in situ study. *Chemical Geology*, 306–307, 92–102. <https://doi.org/10.1016/j.chemgeo.2012.02.021>
- Ganssen, G., & Troelstra, S. R. (1987). Paleoenvironmental changes from stable isotopes in planktonic Foraminifera from Eastern Mediterranean sapropels. *Marine Geology*, 75, 221–230. [https://doi.org/10.1016/0025-3227\(87\)90105-8](https://doi.org/10.1016/0025-3227(87)90105-8)
- Ganssen, G. M., Peeters, F. J. C., Metcalfe, B., Anand, P., Jung, S. J. A., Kroon, D., & Brummer, G.-J. A. (2011). Quantifying sea surface temperature ranges of the Arabian Sea for the past 20 000 years. *Climate of the Past*, 7, 1337–1349. <https://doi.org/10.5194/cp-7-1337-2011>
- Garcia, H. E., Weathers, K. W., Paver, Ch. R., Smolyar, I. V., Boyer, T. P., Locarnini, R. A., Zweng, M. M., Mishonov, A. V., Baranova, O. K., Seidov, D., & Reagan, J. R. (2018). Dissolved oxygen, apparent oxygen utilization, and dissolved oxygen saturation. In S. Levitus & A. Mishonov (Eds.), *World Ocean Atlas 2018* (pp. 1–27). Silver Spring, National Oceanic; Atmospheric Administration. http://data.nodc.noaa.gov/woa/WOA13/DOC/woa13_vol3.pdf
- González-Dávila, M. (2016a). *Physical oceanography measured on water bottle samples at Site ESTOC in 2005*. PANGAEA. <https://doi.org/10.1594/PANGAEA.856610>
- González-Dávila, M. (2016b). *Physical oceanography measured on water bottle samples at Site ESTOC in 2006*. PANGAEA. <https://doi.org/10.1594/PANGAEA.856611>
- Good, S. A., Martin, M. J., & Rayner, N. A. (2013). EN4: Quality controlled ocean temperature and salinity profiles and monthly objective analyses with uncertainty estimates. *Journal of Geophysical Research, Oceans*, 118, 6704–6716. <https://doi.org/10.1002/2013JC009067>
- Gorbarenko, S. A., & Southon, J. R. (2000). Detailed Japan Sea paleoceanography during the last 25 kyr: Constraints from AMS dating and $\delta^{18}\text{O}$ of planktonic Foraminifera. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 156, 177–193. [https://doi.org/10.1016/S0031-0182\(99\)00137-6](https://doi.org/10.1016/S0031-0182(99)00137-6)
- Graham, M. H. (2003). Confronting multicollinearity in ecological multiple regression. *Ecology*, 84, 2809–2815. <https://doi.org/10.1890/02-3114>
- Groeneveld, J., Filipsson, H. L., Austin, W. E. N., Darling, K., McCarthy, D., Quintana Krupinski, N. B., Bird, C., & Schweizer, M. (2018). Assessing proxy signatures of temperature, salinity, and hypoxia in the Baltic Sea through Foraminifera-based geochemistry and faunal assemblages. *Journal of Micropalaeontology*, 37, 403–429. <https://doi.org/10.5194/jm-37-403-2018>
- Groeneveld, J., Ho, S. L., Mackensen, A., Mohtadi, M., & Laepple, Th. (2019). Deciphering the variability in Mg/Ca and stable oxygen isotopes of individual Foraminifera. *Paleoceanography and Paleoclimatology*, 34, 755–773. <https://doi.org/10.1029/2018PA003533>
- Groeneveld, J., Steph, S., Tiedemann, R., Garbe-Schönberg, C.-D., Nürnberg, D., & Sturm, A. (2006). Pliocene mixed-layer oceanography for Site 1241, using combined Mg/Ca and $\delta^{18}\text{O}$ analyses of *Globigerinoides sacculifer*. *Proceedings of the Ocean Drilling Program: Scientific Results*, 202, 1–27. <https://doi.org/10.2973/odp.proc.sr.202.209.2006>
- Haenel, P. (1987). Intérêt paléoocéanographique d'*Orbulina universa* d'Orbigny (foraminifère). *Oceanologica Acta*, 10, 15–25.
- Hemleben, Ch., Meischner, D., Zahn, R., Almogi-Labin, A., Erlenkeuser, H., & Hiller, B. (1996). Three hundred eighty thousand year long stable isotope and faunal records from the Red Sea: Influence of global sea level change on hydrography. *Paleoceanography*, 11, 147–156. <https://doi.org/10.1029/95PA03838>
- Hemleben, Ch., Spindler, M., Breitinger, I., & Ott, R. (1987). Morphological and physiological responses of *Globigerinoides sacculifer* (Brady) under varying laboratory conditions. *Marine Micropaleontology*, 12, 305–324. [https://doi.org/10.1016/0377-8398\(87\)90025-9](https://doi.org/10.1016/0377-8398(87)90025-9)
- Henehan, M. J., Foster, G. L., Bostock, H. C., Greenop, R., Marshall, B. J., & Wilson, P. A. (2016). A new boron isotope-pH calibration for *Orbulina universa*, with implications for understanding and accounting for 'vital effects'. *Earth and Planetary Science Letters*, 454, 282–292. <https://doi.org/10.1016/j.epsl.2016.09.024>
- Hoeffding, W. (1948). A non-parametric test of independence. *The Annals of Mathematical Statistics*, 19, 546–557. <https://www.jstor.org/stable/2236021>
- Ivanova, E. V. (1985). Late Quaternary biostratigraphy and paleotemperatures of the Red Sea and the Gulf of Aden based on planktonic Foraminifera and pteropods. *Marine Micropaleontology*, 9, 335–364. [https://doi.org/10.1016/0377-8398\(85\)90003-9](https://doi.org/10.1016/0377-8398(85)90003-9)

- Kauffmann, L., & Rousseeuw, P. J. (2009). *Finding groups in data: An introduction to cluster analysis*. Hoboken, New Jersey, John Wiley & Sons, Ltd.
- Kawahata, H. (2005). Stable isotopic composition of two morphotypes of *Globigerinoides ruber* (white) in the subtropical gyre in the North Pacific. *Paleontological Research*, 9, 27–35. <https://doi.org/10.2517/prpsj.9.27>
- Kendall, C., & Caldwell, E. A. (1998). Fundamentals of isotope geochemistry. In C. Kendall & J. J. McDonnell (Eds.), *Isotope tracers in catchment hydrology* (pp. 51–86). Amsterdam, Elsevier. <https://wwwrcamnl.wr.usgs.gov/isoig/isopubs/itchch2.html>
- Killingley, J. S., Johnson, R. F., & Berger, W. H. (1981). Oxygen and carbon isotopes of individual shells of planktonic Foraminifera from Ontong-Java plateau, equatorial Pacific. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 33, 193–204. [https://doi.org/10.1016/0031-0182\(81\)90038-9](https://doi.org/10.1016/0031-0182(81)90038-9)
- King, A. L., & Howard, W. R. (2005). $\delta^{18}\text{O}$ seasonality of planktonic Foraminifera from Southern Ocean sediment traps: Latitudinal gradients and implications for paleoclimate reconstructions. *Marine Micropaleontology*, 56, 1–24. <https://doi.org/10.1016/j.marmicro.2005.02.008>
- Kučera, M. (2007). Planktonic Foraminifera as tracers of past oceanic environments. In C. Hillaire-Marcel, A. de Vernal, & H. Chamley (Eds.), *Proxies in late Cenozoic paleoceanography* (pp. 213–262). Amsterdam, Elsevier. [https://doi.org/10.1016/S1572-5480\(07\)01011-1](https://doi.org/10.1016/S1572-5480(07)01011-1)
- Laepple, Th., & Huybers, P. (2014a). Global and regional variability in marine surface temperatures. *Geophysical Research Letters*, 41, 2528–2534. <https://doi.org/10.1002/2014GL059345>
- Laepple, Th., & Huybers, P. (2014b). Ocean surface temperature variability: Large model–data differences at decadal and longer periods. *Proceedings of the National Academy of Sciences of the United States of America*, 111, 16682–16687. <https://doi.org/10.1073/pnas.1412077111>
- Lea, D. W. (2002). Trace elements in foraminiferal calcite. In B. K. Sen Gupta (Ed.), *Modern foraminifera* (pp. 259–277). Dordrecht, Kluwer Academic Publishers.
- Lewis, E., Wallace, D., & Allison, L. J. (1998). *Program developed for CO₂ system calculations* (tech. rep. No. 4735). Carbon Dioxide Information Analysis Center. Oak Ridge. <http://cdiac.ornl.gov/oceans/co2rprt.html>
- Locarnini, R. A., Mishonov, A. V., Baranova, O. K., Boyer, T. P., Zweng, M. M., Garcia, H. E., Reagan, J. R., Seidov, D., Weathers, K. W., Paver, Ch. R., & Smolyar, I. V. (2018). Temperature. In A. Mishonov (Ed.), *World Ocean Atlas 2018* (pp. 1–45). Silver Spring, National Oceanic; Atmospheric Administration. https://data.nodc.noaa.gov/woa/WOA18/DOC/woa18_vol1.pdf
- Lohmann, G. P. (1995). A model for variation in the chemistry of planktonic Foraminifera due to secondary calcification selective dissolution. *Paleoceanography*, 10, 445–457. <https://doi.org/10.1029/95PA00059>
- Mann, H. B., & Whitney, D. R. (1947). On a test of whether one of two random variables is stochastically larger than the other. *The Annals of Mathematical Statistics*, 18, 50–60. <https://doi.org/10.1214/aoms/1177730491>
- Marshall, B. J., Thunell, R. C., Henehan, M. J., Astor, Y., & Wejnert, K. R. (2013). Planktonic foraminiferal area density as a proxy for carbonate ion concentration: A calibration study using the Cariaco Basin Ocean Time Series. *Paleoceanography*, 28, 1–14. <https://doi.org/10.1002/palo.20034>
- McConaughey, T. A. (1989). Biomineralization mechanisms (R. E. Crick, Ed.). In R. E. Crick (Ed.), *Origin, evolution, and modern aspects of biomineralization in plants and animals*, Arlington, Plenum Press.
- Mehrbach, C., Culberson, C. H., Hawley, J. E., & Pytcowicz, R. M. (1973). Measurement of the apparent dissociation constants of carbonic acid in seawater at atmospheric pressure. *Limnology and Oceanography*, 18, 897–907. <https://doi.org/10.4319/lo.1973.18.6.0897>
- Meiland, J., Siccha, M., Weinkauf, M. F. G., Jonkers, L., Morard, R., Baranowski, U., Baumeister, A., Bertlich, J., Brummer, G.-J., Debray, P., Fritz-Endres, Th., Groeneveld, J., Magerl, L., Munz, P., Rillo, M. C., Schmidt, Ch., Takagi, H., Theara, G., & Kučera, M. (2019). Highly replicated sampling reveals no diurnal vertical migration but stable species-specific vertical habitats in planktonic Foraminifera. *Journal of Plankton Research*, 41, 127–141. <https://doi.org/10.1093/plankt/fbz002>
- Michaelis, L., & Menten, M. L. (1913). Die Kinetik der Invertinwirkung [The kinetics of invertase action]. *Biochemische Zeitschrift*, 49, 333–369. https://path.upmc.edu/divisions/chp/PDF/Michaelis-Menten_Kinetik.pdf
- Milker, Y., Rachmayani, R., Weinkauf, M. F. G., Prange, M., Raitzsch, M., Schulz, M., & Kučera, M. (2013). Global and regional sea surface temperature trends during marine isotope stage 11. *Climate of the Past*, 9, 2231–2252. <https://doi.org/10.5194/cp-9-2231-2013>
- Mohan, R., Shetye, S. S., Tiwari, M., & Anilkumar, N. (2015). Secondary calcification of planktic Foraminifera from the Indian sector of southern ocean. *Acta Geologica Sinica (English Edition)*, 89, 27–37. <https://doi.org/10.1111/1755-6724.12392>
- Mortyn, P. G., & Charles, Ch. D. (2003). Planktonic foraminiferal depth habitat and $\delta^{18}\text{O}$ calibrations: Plankton tow results from the Atlantic sector of the Southern Ocean. *Paleoceanography*, 18, Article 1037. <https://doi.org/10.1029/2001PA000637>
- Murthaugh, P. A. (2014). In defense of *p* values. *Ecology*, 95, 611–617. <https://doi.org/10.1890/13-0590.1>
- Norris, R. D. (1998). Recognition and macroevolutionary significance of photosymbiosis in molluscs, corals, and Foraminifera. In R. D. Norris & R. M. Corfield (Eds.), *Isotope paleobiology and paleoecology*. Lawrence, The Paleontological Society.
- Numberger, L., Hemleben, Ch., Hoffmann, R., Mackensen, A., Schulz, H., Wunderlich, J.-M., & Kučera, M. (2009).

- Habitats, abundance patterns and isotopic signals of morphotypes of the planktonic foraminifer *Globigerinoides ruber* (d'Orbigny) in the Eastern Mediterranean Sea since the marine isotopic stage 12. *Marine Micropaleontology*, 73, 90–104. <https://doi.org/10.1016/j.marmicro.2009.07.004>
- Ohno, Y., Fujita, K., Toyofuku, T., & Nakamura, T. (2016). Cytological observations of the large symbiotic foraminifer *Amphisorus kudakajimensis* using calcein acetoxymethyl ester. *PLoS ONE*, 11, e0165844. <https://doi.org/10.1371/journal.pone.0165844>
- Osborne, E. B., Thunell, R. C., Marshall, B. J., Holm, J. A., Tappa, E. J., Benitez-Nelson, C., Cai, W.-J., & Chen, B. (2016). Calcification of the planktonic Foraminifera *Globigerina bulloides* and carbonate ion concentration: Results from the Santa Barbara Basin. *Paleoceanography*, 31, 1083–1102. <https://doi.org/10.1002/2016PA002933>
- Pearson, P. N. (2012). Oxygen isotopes in Foraminifera: Overview and historical review. In L. C. Ivany & B. T. Huber (Eds.), *Reconstructing Earth's deep-time climate—the state of the art in 2012* (pp. 1–38). Cambridge, Cambridge University Press. <https://doi.org/10.1017/S1089332600002539>
- Peeters, F., Ivanova, E., Conan, S., Brummer, G.-J., Ganssen, G., Troelstra, S., & van Hinte, J. (1999). A size analysis of planktic Foraminifera from the Arabian Sea. *Marine Micropaleontology*, 36, 31–63. [https://doi.org/10.1016/S0377-8398\(98\)00026-7](https://doi.org/10.1016/S0377-8398(98)00026-7)
- Pérez, F. F., Gilcoto, M., & Ríos, A. F. (2003). Large and mesoscale variability of the water masses and the deep chlorophyll maximum in the Azores front. *Journal of Geophysical Research*, 108, Article 3215. <https://doi.org/10.1029/2000JC000360>
- R Core Team. (2019). *R: A language and environment for statistical computing*. Version 3.6.2. R Foundation for Statistical Computing. Vienna. <http://www.R-project.org/>
- Raitzsch, M., Dueñas-Bohórquez, A., Reichart, G.-J., de Nooyer, L. J., & Bickert, T. (2010). Incorporation of Mg and Sr in calcite of cultured benthic Foraminifera: Impact of calcium concentration and associated calcite saturation state. *Biogeosciences*, 7, 869–881. <https://doi.org/10.5194/bg-7-869-2010>
- Ravelo, A. Ch., & Hillaire-Marcel, C. (2007). The use of oxygen and carbon isotopes of Foraminifera in paleoceanography. In C. Hillaire-Marcel, A. de Vernal, & H. Chamley (Eds.), *Proxies in late Cenozoic paleoceanography* (pp. 735–764). Amsterdam, Elsevier. [https://doi.org/10.1016/S1572-5480\(07\)01023-8](https://doi.org/10.1016/S1572-5480(07)01023-8)
- Rebotim, A., Voelker, A. H. L., Jonkers, L., Wanek, J. J., Meggers, H., Schiebel, R., Fraile, I., Schulz, M., & Kučera, M. (2017). Factors controlling the depth habitat of planktonic Foraminifera in the subtropical eastern North Atlantic. *Biogeosciences*, 14, 827–859. <https://doi.org/10.5194/bg-14-827-2017>
- Rebotim, A., Voelker, A. H. L., Jonkers, L., Wanek, J. J., Schulz, M., & Kučera, M. (2019). Calcification depth of deep-dwelling planktonic Foraminifera from the eastern North Atlantic constrained by stable oxygen isotope ratios of shells from stratified plankton tows. *Journal of Micropalaeontology*, 38, 113–131. <https://doi.org/10.5194/jm-38-113-2019>
- Regenberg, M., Nürnberg, D., Steph, S., Groeneveld, J., Garbe-Schönberg, D., Tiedemann, R., & Dullo, W.-C. (2006). Assessing the effect of dissolution on planktonic foraminiferal Mg/Ca ratios: Evidence from Caribbean core tops. *Geochemistry, Geophysics, Geosystems*, 7, Q07P15. <https://doi.org/10.1029/2005GC001019>
- Reynolds, R. W., Rayner, N. A., Smith, Th. M., Stokes, D. C., & Wang, W. (2002). An improved in situ and satellite SST analysis for climate. *Journal of Climate*, 15, 1609–1625. [https://doi.org/10.1175/1520-0442\(2002\)015<1609:AIISAS>2.0.CO;2](https://doi.org/10.1175/1520-0442(2002)015<1609:AIISAS>2.0.CO;2)
- Rohling, E. J., & Cooke, S. (2002). Stable oxygen and carbon isotopes in foraminiferal carbonate shells. In B. K. Sen Gupta (Ed.), *Modern foraminifera* (pp. 240–258). Dordrecht, Kluwer Academic Publishers.
- Rohling, E. J., Sprovieri, M., Cane, T., Casford, J. S. L., Cooke, S., Bouloubassi, I., Emeis, K. – Ch., Schiebel, R., Rogerson, M., Hayes, A., Jorissen, F. J., & Kroon, D. (2004). Reconstructing past planktic foraminiferal habitats using stable isotope data: A case history for Mediterranean Sapropel S5. *Marine Micropaleontology*, 50, 89–123. [https://doi.org/10.1016/S0377-8398\(03\)00068-9](https://doi.org/10.1016/S0377-8398(03)00068-9)
- Rosenthal, Y. (2007). Elemental proxies for reconstructing Cenozoic seawater paleotemperatures from calcareous fossils. In C. Hillaire-Marcel, A. de Vernal, & H. Chamley (Eds.), *Proxies in late Cenozoic paleoceanography* (pp. 765–797). Amsterdam, Elsevier. [https://doi.org/10.1016/S1572-5480\(07\)01024-X](https://doi.org/10.1016/S1572-5480(07)01024-X)
- Rosenthal, Y., Oppo, D. W., & Lindsey, B. K. (2003). The amplitude and phasing of climate change during the last deglaciation in the Sulu Sea, western equatorial Pacific. *Geophysical Research Letters*, 30, Article 1428. <https://doi.org/10.1029/2002GL016612>
- Rousseeuw, P. J., Ruts, I., & Tukey, J. W. (1999). The bagplot: A bivariate boxplot. *The American Statistician*, 53, 382–387. <https://doi.org/10.1080/00031305.1999.10474494>
- Russell, A. D., Hönnisch, B., Spero, H. J., & Lea, D. W. (2004). Effects of seawater carbonate ion concentration and temperature on shell U, Mg, and Sr in cultured planktonic Foraminifera. *Geochimica et Cosmochimica Acta*, 68, 4347–4361. <https://doi.org/10.1016/j.gca.2004.03.013>
- Schiebel, R. (2002). Planktic foraminiferal sedimentation and the marine calcite budget. *Global Biogeochemical Cycles*, 16, Article 13. <https://doi.org/10.1029/2001GB001459>
- Schiebel, R., Barker, S., Lendt, R., Thomas, H., & Bollmann, J. (2007). Planktic foraminiferal dissolution in the twilight zone. *Deep-Sea Research, Part II: Topical Studies in Oceanography*, 54, 676–686. <https://doi.org/10.1016/j.dsr2.2007.01.009>

- Schiebel, R., & Hemleben, Ch. (2017). *Planktic foraminifers in the modern ocean*. Berlin, Heidelberg, Springer-Verlag. <https://doi.org/10.1007/978-3-662-50297-6>
- Schiffelbein, P., & Hills, S. (1984). Direct assessment of stable isotope variability in planktonic Foraminifera populations. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 48, 197–213. [https://doi.org/10.1016/0031-0182\(84\)90044-0](https://doi.org/10.1016/0031-0182(84)90044-0)
- Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D. J., Hartenstein, V., Eliceiri, K., Tomancak, P., & Cardona, A. (2012). Fiji: An open-source platform for biological-image analysis. *Nature Methods*, 9, 676–682. <https://doi.org/10.1038/nmeth.2019>
- Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of image analysis. *Nature Methods*, 9, 671–675. <https://doi.org/10.1038/nmeth.2089>
- Shackleton, N. J., Corfield, R. M., & Hall, M. A. (1985). Stable isotope data and the ontogeny of Paleocene planktonic Foraminifera. *Journal of Foraminiferal Research*, 15, 321–336. <https://doi.org/10.2113/gsjfr.15.4.321>
- Spero, H. J. (1988). Ultrastructural examination of chamber morphogenesis and biomineralization in the planktonic foraminifer *Orbulina universa*. *Marine Biology*, 99, 9–20. <https://doi.org/10.1007/BF00644972>
- Spero, H. J. (1998). Life history and stable isotope geochemistry of planktonic Foraminifera. In R. D. Norris & R. M. Corfield (Eds.), *Isotope paleobiology and paleoecology* (pp. 7–36). Pittsburgh, The Paleontological Society.
- Spero, H. J., Bijma, J., Lea, D. W., & Bemis, B. E. (1997). Effect of seawater carbonate concentration on foraminiferal carbon and oxygen isotopes. *Nature*, 390, 497–500. <https://doi.org/10.1038/37333>
- Spero, H. J., & Lea, D. W. (1993). Intraspecific stable isotope variability in the planktic Foraminifera *Globigerinoides sacculifer*: Results from laboratory experiments. *Marine Micropaleontology*, 22, 221–234. [https://doi.org/10.1016/0377-8398\(93\)90045-Y](https://doi.org/10.1016/0377-8398(93)90045-Y)
- Spero, H. J., Lerche, I., & Williams, D. F. (1991). Opening the carbon isotope “vital effect” black box, 2, quantitative model for interpreting foraminiferal carbon isotope data. *Paleoceanography*, 6, 639–655. <https://doi.org/10.1029/91PA02022>
- Spero, H. J., & Williams, D. F. (1988). Extracting environmental information from planktonic foraminiferal $\delta^{13}\text{C}$ data. *Nature*, 335, 717–719. <https://doi.org/10.1038/335717a0>
- Spötl, C., & Vennemann, T. W. (2003). Continuous-flow isotope ratio mass spectrometric analysis of carbonate minerals. *Rapid Communications in Mass Spectrometry*, 17, 1004–1006. <https://doi.org/10.1002/rcm.1010>
- Steinhardt, J., Cléroux, C., de Nooijer, L. J., Brummer, G.-J., Zahn, R., Ganssen, G., & Reichart, G.-J. (2015). Reconciling single-chamber Mg/Ca with whole shell $\delta^{18}\text{O}$ in surface to deep-dwelling planktonic Foraminifera from the Mozambique Channel. *Biogeosciences*, 12, 2411–2429. <https://doi.org/10.5194/bg-12-2411-2015>
- Steinke, S., Chiu, H.-Y., Yu, P.-S., Shen, C.-C., Löwemark, L., Mii, H.-S., & Chen, M.-T. (2005). Mg/Ca ratios of two *Globigerinoides ruber* (white) morphotypes: Implications for reconstructing past tropical/subtropical surface water conditions. *Geochemistry, Geophysics, Geosystems*, 6, Q11005. <https://doi.org/10.1029/2005GC000926>
- Steph, S., Regenberg, M., Tiedemann, R., Mulitza, S., & Nürnberg, D. (2009). Stable isotopes of planktonic Foraminifera from tropical Atlantic/Caribbean core-tops: Implications for reconstructing upper ocean stratification. *Marine Micropaleontology*, 71, 1–19. <https://doi.org/10.1016/j.marmicro.2008.12.004>
- Stott, L. D., & Tang, C. M. (1996). Reassessment of foraminiferal-based tropical sea surface $\delta^{18}\text{O}$ paleotemperatures. *Paleoceanography*, 11, 37–56. <https://doi.org/10.1029/95PA03344>
- Toyofuku, T., Matsuo, M. Y., de Nooijer, L. J., Nagai, Y., Kawada, S., Fujita, K., Reichart, G.-J., Nomaki, H., Tsuchiya, M., Sakaguchi, H., & Kitazato, H. (2017). Proton pumping accompanies calcification in Foraminifera. *Nature Communications*, 8, Article 14145. <https://doi.org/10.1038/ncomms14145>
- van Dijk, I., de Nooijer, L. J., Wolthers, M., & Reichart, G.-J. (2017). Impacts of pH and $[\text{CO}_3^{2-}]$ on the incorporation of Zn in foraminiferal calcite. *Geochimica et Cosmochimica Acta*, 197, 263–277. <https://doi.org/10.1016/j.gca.2016.10.031>
- van Eijden, A. J. M. (1995). Morphology and relative frequency of planktic foraminiferal species in relation to oxygen isotopically inferred depth habitats. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 113, 267–301. [https://doi.org/10.1016/0031-0182\(95\)00057-S](https://doi.org/10.1016/0031-0182(95)00057-S)
- Vetter, L., Kozdon, R., Mora, C. I., Eggin, S. M., Valley, J. W., Hönsch, B., & Spero, H. J. (2013). Micron-scale intrashell oxygen isotope variation in cultured planktic foraminifers. *Geochimica et Cosmochimica Acta*, 107, 267–278. <https://doi.org/10.1016/j.gca.2012.12.046>
- Wagenmakers, E.-J., & Farrell, S. (2004). AIC model selection using Akaike weights. *Psychonomic Bulletin & Review*, 11, 192–196. <https://doi.org/10.3758/BF03206482>
- Wang, L. (2000). Isotopic signals in two morphotypes of *Globigerinoides ruber* (white) from the South China Sea: Implications for monsoon climate change during the last glacial cycle. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 161, 381–394. [https://doi.org/10.1016/S0031-0182\(00\)00094-8](https://doi.org/10.1016/S0031-0182(00)00094-8)
- Waniek, J., Koeve, W., & Prien, R. D. (2000). Trajectories of sinking particles and the catchment areas above sediment traps in the northeast Atlantic. *Journal of Marine Research*, 58, 983–1006. <https://doi.org/10.1357/002224000763485773>
- Waniek, J. J., Schulz-Bull, D. E., Blanz, Th., Prien, R. D., Oschlies, A., & Müller, Th. J. (2005). Interannual variability of deep water particle flux in relation to production and

- lateral sources in the northeast Atlantic. *Deep-Sea Research, Part I: Oceanographic Research Papers*, 52, 33–50. <https://doi.org/10.1016/j.dsr.2004.08.008>
- Weiner, A., Aurahs, R., Kurasawa, A., Kitazato, H., & Kučera, M. (2012). Vertical niche partitioning between cryptic sibling species of a cosmopolitan marine planktonic protist. *Molecular Ecology*, 21, 4063–4073. <https://doi.org/10.1111/j.1365-294X.2012.05686.x>
- Weinkauf, M. F. G., Kunze, J. G., Waniek, J. J., & Kučera, M. (2016). Seasonal variation in shell calcification of planktonic Foraminifera in the NE Atlantic reveals species-specific response to temperature, productivity, and optimum growth conditions. *PLoS ONE*, 11, e148363. <https://doi.org/10.1371/journal.pone.0148363>
- Weinkauf, M. F. G., Moller, T., Koch, M. C., & Kučera, M. (2013). Calcification intensity in planktonic Foraminifera reflects ambient conditions irrespective of environmental stress. *Biogeosciences*, 10, 6639–6655. <https://doi.org/10.5194/bg-10-6639-2013>
- Weinkauf, M. F. G., Zwick, M. M., & Kučera, M. (in press). Constraining the role of shell porosity in the regulation of shell calcification intensity in the modern planktonic foraminifer *Orbulina universa* d'Orbigny. *Journal of Foraminiferal Research*.
- Wit, J. C., Reichart, G.-J., & Ganssen, G. M. (2013). Unmixing of stable isotope signals using single specimen $\delta^{18}\text{O}$ analyses. *Geochemistry, Geophysics, Geosystems*, 14, 1312–1320. <https://doi.org/10.1002/ggge.20101>
- Zarkogiannis, S. D., Antonarakou, A., Tripati, A., Kontakiotis, G., Mortyn, P. G., Drinia, H., & Greaves, M. (2019). Influence of surface ocean density on planktonic Foraminifera calcification. *Scientific Reports*, 9, Article 533. <https://doi.org/10.1038/s41598-018-36935-7>
- Zeebe, R. E., Bijma, J., Hönnisch, B., Sanyal, A., Spero, H. J., & Wolf-Gladrow, D. A. (2008). Vital effects and beyond: A modelling perspective on developing palaeoceanographical proxy relationships in Foraminifera. In W. E. N. Austin & R. H. James (Eds.), *Biogeochemical controls on palaeoceanographic environmental proxies* (pp. 45–58). London, The Geological Society.
- Zweng, M. M., Reagan, J. R., Seidov, D., Boyer, T. P., Locarnini, R. A., Garcia, H. E., Mishonov, A. V., Baranova, O. K., Weathers, K. W., Paver, Ch. R., & Smolyar, I. V. (2018). Salinity. In A. Mishonov (Ed.), *World Ocean Atlas 2018* (pp. 1–40). Silver Spring, National Oceanic; Atmospheric Administration. https://data.nodc.noaa.gov/woa/WOA18/DOC/woa18_vol2.pdf