
Study On the Mechanism of Atmospheric Electric Field Anomalies Before Earthquakes

Su Jianfeng^{1,3,*}, Chen Hai^{1,2}

(1) National Space Science Center, CAS, Beijing, 100190, China

(2) Nation Space Science Data Center, National Space Science Center, Beijing, 101407, China

(3) State Key Laboratory of Space Weather, National Space Science Center, CAS, Beijing, 100190, China

Abstract. Negative abnormal changes of atmospheric electric field have been monitored many times before earthquakes, especially before large earthquakes. However, its mechanism has not been clearly concluded yet. Based on the comparative analysis of the positive anomaly characteristics of atmospheric electric field before lightning and geoelectric field before earthquakes in earthquake areas, this paper concludes that the negative anomalies of atmospheric electric field before earthquakes are caused by the rise of the geoelectric potential in earthquake areas, and the areas with the abnormal rise of the geoelectric potential are potential earthquake areas. Based on the inference above, this paper further proposes a comprehensive earthquake early warning method with station network layout by geoelectric field instruments in combination with the atmospheric electric field data monitored by Electric Field Mills (EFMs). The research results will provide an possible reference for exploring the mechanism of earthquake generation, and will be help to explore earthquake early warning.

Key Words. Atmospheric Electric Field, Geoelectric Field, Earthquake

1. Positive anomalies of atmospheric electric field before lightning

Since the 17th century, scientists have been studying the phenomenon of atmospheric lightning. In 1920, British scientist Wilson [1,2] established a relatively complete concept of atmospheric circuit. It is found that when the sky is clear, there is always an electric field pointing to the ground in the atmosphere, and the electric

potential gradient is about 100V/m. Although the scientific community has never had a clear conclusion on the formation mechanism of thunderstorm clouds, there is a clear consensus on the occurrence mechanism of lightning: if the electric potential difference between the bottom of the cloud layers and the ground reaches a certain degree, lightning will occur, and when the first discharge is completed, the electric potential difference between the bottom of the cloud layers and the ground becomes smaller, then the cloud layers gather charges again until the next lightning process.

In order to study the electrical characteristics of the atmosphere, scientists invented Electric Field Mill (EFM) to monitor the absolute electric field of the atmosphere relative to the ground, as shown in the left side of Figure 1. EFM is also used for early warning of lightning, and the basic principle of which is to measure the electric field of atmospheric cloud layers pointing to the ground. Taking the ground as zero potential, when the positively charged cloud layers approach, the atmospheric electric field will increase and a positive anomaly will occur. When the electric field increases to a certain extent, a lightning phenomenon will occur, as shown in the right side of Figure 1.

2. Negative anomalies of atmospheric electric field before earthquakes

At the same time, negative abnormal changes of atmospheric electric field have been monitored many times before earthquakes [5,6,7,8,9,10,11,12]. Some people try to find the cause, For example, Professor Harrison et.al mentioned the radon emanation [13]. Moreover, Freund et.al also given one the possibility from the stresses rock [14]. Especially before large earthquakes, the negative abnormal changes of atmospheric electric field are obvious, such as the 2008 Wenchuan earthquake in Sichuan province, as shown in Figure 2 [7]. The left side of Figure 2 shows the locations of EFMs in Wenjiang and Pixian counties, as well as the epicenter of the earthquake. The right side of Figure 2 shows the data of atmospheric electric field before the earthquake, from which it can be seen that the atmospheric electric field presented obvious negative anomaly within a few hours before the earthquake and

58 quickly recovered to normal after the earthquake.

59

60 The meteorological data shows that the weather condition before the
61 Wenchuan earthquake was sunny, and there were no large number of clouds. Then
62 according to the law of conservation of electric charge, the negative anomaly of
63 atmospheric electric field could only be caused by a positive charge of the ground, i.e.,
64 the geoelectric field was at a high positive potential before the earthquake.

65 **3. Positive anomalies of geoelectric field before earthquakes**

66 In order to further confirm the result of the analysis above, we can analyze it
67 directly from the perspective of geoelectric field. Reference [16] gives the data
68 recorded by the geoelectric field meter at Pixian station, which is about 50km away
69 from the epicenter of the Wenchuan earthquake. It can be seen from Figure 3 (a) (b)
70 that an obvious discharge occurred during the earthquake. Figure 3 (a) (b) does not
71 specifically analyze the direction of the geoelectric field, but we can roughly deduce
72 the direction change of the geoelectric field before and after the earthquake from the
73 data.

74 The epicenter of the Wenchuan earthquake is located at 31.0° latitude and 103.4°
75 longitude, and the location of the Pixian geoelectric field meter is about 30.6°
76 latitude and 104.0° longitude. According to the data graph, before the earthquake, the
77 NS electric field vector was about -20V/m , the EW electric field vector was about
78 40V/m , and after the earthquake, the NS electric field vector was about -40V/m , and
79 the EW electric field vector was about -40V/m , as shown in Figure 3 (d). It is obvious
80 that the electric potential at the epicenter before the earthquake was high, pointing
81 from the Wenchuan epicenter to Pixian station (red arrow), and after the earthquake, it
82 became pointing from the northeast to the southwest (black arrow).

83 Similarly, let's look at the 6.2 magnitude earthquake occurred in Zhangbei area,
84 Hebei Province, China on January 10, 1998. The geoelectric field equipment located
85 in Baodi, Tianjin recorded that the direction of the geoelectric field \mathbf{E} vector
86 ($\mathbf{E}=\mathbf{E1}+\mathbf{E2}$, where $\mathbf{E1}$ is the SN component and $\mathbf{E2}$ is the EW component) changed

significantly. Before the earthquake, E vector was pointing from the Zhangbei epicenter to Baodi station, as shown in Figure 4 [6].

According to the changes in the direction of the geoelectric field before and after the earthquake, the whole process can be divided into four stages:

I:1997-11-21~1997-12-05 (before the earthquake);

II:1997-12-06~1998-01-10 (from the negative anomaly before the earthquake to the earthquake);

III:1998-01-11~1998-01-23 (from the earthquake to normal);

IV:1998-01-07~1998-01-10 (4 days before the earthquake).

The left side of Figure 4 shows the locations of the Zhangbei epicenter and Baodi geoelectric field equipment, and the right side of Figure 4 shows the direction of geoelectric field E in each period. It is obvious that the direction of the geoelectric field changed significantly before and after the earthquake. Before the earthquake, the geoelectric field pointed from the Zhangbei epicenter to Baodi station. This shows that an obvious positive anomaly of geoelectric field occurred in the Zhangbei earthquake area before the earthquake, that is, the geoelectric field in the earthquake area was at high potential before the earthquake.

4. Analysis of negative anomalies of atmospheric electric field before earthquakes

The interior of the earth is equivalent to a complex and dynamic capacitor. With the analysis above, it can be determined that the geoelectric field in an earthquake area is at high potential before the earthquake, and a discharge phenomenon occurred during the earthquake, which has the same characteristics as lightning phenomenon in the atmosphere. From this, we can draw the following possible conclusions:

a) Earthquakes may be "underground lightning" phenomena;

b) There may be an "underground lightning" earthquake type;

c) High geoelectric potential occurs before some earthquakes.

In case of a) b), it is the process of underground electric energy release -

116 earthquake is the process of underground lightning. Electric energy is converted into
 117 other forms of energy, including mechanical energy and light energy. The mechanical
 118 energy can produce ground vibration and generate ground sound, which explains the
 119 occurrence of earthquake phenomenons (ground thundering); The light energy can
 120 produce ground light, which explains the ground light phenomenons that can be
 121 observed during a large earthquake at night (ground flashing [17]).

122 In any of the cases above, the atmospheric lightning early warning principle can
 123 be used for earthquake early warning, and monitoring high potential of the geoelectric
 124 field can do help for earthquake early warning.

125 **5. Suggestions and Prospects - Earthquake warning**

126 Based on the mechanism research above, we can draw a conclusion that the
 127 principle of atmospheric lightning early warning can also be used for earthquake early
 128 warning, at least for some earthquakes. Special design can be made for EFM, such as
 129 avoiding the influence of meteorological factors. In the way of station network layout,
 130 multiple geoelectric field instruments can be used to monitor the direction of the
 131 geoelectric field in order to locate high geoelectric potential areas, i.e, the potential
 132 earthquake areas, and make a comprehensive judgment based on the monitoring data
 133 of EFMs.

134 The underground geological structure is far more complex than the atmosphere,
 135 so further in-depth research is needed. The research results above are just references
 136 for the scientific community.

137 **References**

- 138 [1] D.R. MacCorman, and W.D. Rust. (1998).The Electrical Nature of Storm, Oxford University Press, New York,p,185-187.
 139 [2] Joseph R.Dwyer, and Martin A.Uman. 2014. The physics of lightning. Physics Report,534,pp.147-241.
 140 doi.org/10.1016/j.physrep.2013.09.004
 141 [3] EFM-100_Manual_03102015.p37.
 142 [4] Jennifer G. Wilson a , Kenneth L. Cummins b,c.2021.Thunderstorm and fair-weather quasi-static electric fields over land
 143 and ocean. Atmospheric Research 257 (2021) 105618, doi.org/10.1016/j.atmosres.2021.105618
 144 [5] Hao Jianguo, Tang Tianming, Li Derui, 1998. A kind of short-term and imminent precursory information of earthquakes:
 145 study on the anomaly of atmospheric electric field before earthquakes. Acta Seismologica Sinica, 20 (1): 101-108 ,
 146 doi:/10.1007/BF02650462

- [6] Abhijit Choudhury etc.2013. A statistical study on precursory effects of earthquakes observed through the atmospheric vertical electric field in northeast India.. *Annals of geophysics*, Vol.56, No. 3. 2013, R0331; doi:10.4401/ag-6235.
- [7] Chen Tao, Zhang Xiaoxin, et al.,2021. Imminent estimation of earthquake hazard by regional network monitoring the near surface vertical atmospheric electrostatic field. *Chinese Journal of Geophysics* Vol.64. No.4. Apr.,2021
- [8] Xiaobing Jin,Liang Zhang , Junwei Bu, Guilan Qiu, Li Ma, Chang Liu, Yiding Li, 2020.Discussion on anomaly of atmospheric electrostatic field in Wenchuan Ms8.0 earthquake.. *Journal of Electrostatics* 104(2020)103423.doi.org/10.1016/j.elstat.2020.103423
- [9] Tong Xu,Yani Hu, Jian.Wu, Zhensen Wu, Chunbin Li, Zhenwen Xu' Yucheng Suo.2011. Anomalous enhancement of electric field derived from ionosonde data before the great Wenchuan earthquake. *Advances in space research* 47(2011)1001-1005. doi.org/10.1016/j.asr.2010.11.006
- [10] H.g.Silva,M.Bezzeghoud,A.H.Reis,R.N.Rosa,M.Tlemcani,A.A.Araujo,C.Serrano,J.F.Borges,B.Caldeira,and P.F.Biagi. 2011.Atmospheric electrical field decrease during the M=4.1 sousel earthquake(Portugal) , *Nat, Hazards earth Syst,Sci*,11,987-991,doi:10.5194/nhess-11-987-2011.2011.
- [11] Liperovsky, V. A, Meister, C.-V., Liperovskaya, E. V., and Bogdanov, V.V.:2008. on the generation of electric field and infrared radiation in aerosol clouds due to radon emanation in the atmosphere before earthquakes, *Nat, Hazards earth Syst,Sci*,8,1199-1205,doi:10.5194/nhess-8-745-2008.2008.
- [12] Smirnov, S:2008.Association of the negative anomalies of the quasistatic field in atmosphere with Kamchatka seismicity, *Nat, Hazards earth Syst,Sci*,8,745-749,doi:10.5194/nhess-8-745-2008.2008.
- [13] R.G. Harrison, K.L. Aplin and M.J. Rycroft. 2010.Atmospheric electricity coupling between earthquake regions and the ionosphere. *Journal of Atmospheric and Solar-Terrestrial Physics J. Atmos Solar Terr Physics* 72, 376-381 (2010) .doi.org/10.1016/j.jastp.2009.12.004
- [14] Friedemann T.Freund,Ipek G.Kulahci,Gary Cyr, Julia Ling, Matthew Winnick, Jeremy Tregloan-Reed, Minoru M.Freund. 2009.Air ionization at rock surface and pre-earthquake signals. *Journal of Atmospheric and Solar-terrestrial Physics* 71(2009)1824-1834. doi.org/10.1016/j.jastp.2009.07.013
- [15] Mao Tongen, Xi Jilou, Wang Yanqiong, Yin Shuzhi .1999.The variation characteristics of the telluric field in the process of earthquake. *Chinese Journal of Geophysics* Vol.42. No. 4. July., 1999
- [16] Private communication.
- [17] Yuji Enomoto,Tsuneaki Yamabe, Kazuyuki Mizuhara , Shigeki Sugiura, Hitoshi Kondo. 2020.Laboratory investigation of earthquake lighting due to landslide. *Earth,Planet and Space* (2020)72:108. doi.org/10.1186/s40623-020-01237-8.




Fig 1 Schematic diagram of EFM and Increasing field due to approaching thunderclouds




Fig. 2 The locations of EFMs in Wenjiang and Pixian counties, as well as the epicenter of the earthquake and The data of EFMs before the earthquake broke out




Fig. 3 The goelectric field orientation diagram before and after the Wenchuan earthquake based on Pixian station data[18]




Fig.4 The relative position between the epicenter and Baodi station and The regnant direction

276 of the geoelectric field vector \mathbf{E} in each stage in the process of the Zhangbei earthquake Ms 6.2
277 [17]

278

279

280 Advising experts :

281 K.L. Cummins e-mail: kcummins@arizona.edu

282 Giles Harrison e-mail: r.g.harrison@reading.ac.uk

283 Moffett Field e-mail: friedemann.t.freund@nasa.gov

284 Anirban Guha e-mail: anirban1001@yahoo.com

285 H.G. Silva e-mail: hgsilva@uevora.pt

286 S. Smirnov e-mail: sergey@ikir.ru

287 C.-V. Meister e-mail: cvmeister@aip.de

288 Yuji Enomoto e-mail: enomoto@shinshu-u.ac.jp

289

Fig. 1 Schematic diagram of EFM and Increasing field due to approaching thunderclouds.








Fig. 2 The locations of EFMs in Wenjiang and Pixian counties, as well as the epicenter of the earthquake and The data of EFMs before the earthquake broke out.



$E/(KV/m)$








Fig. 3 The geoelectric field orientation diagram before and after the Wenchuan earthquake based on Pixian station data[18].





(c)

(d)

Fig.4 The relative position between the epicenter and Baodi station and The regnant direction of the geoelectric field vector E in each stage in the process of the Zhangbei earthquake Ms 6.2 [17].

